

SIMULATION OF RADIO TELEMETRY FOR HOME RANGE PREDICTING OF WREATHED HORNBILL (RHYTICEROS UDULATUS) ON MOUNT UNGARAN

by Siti Alimah

Submission date: 03-Jul-2020 09:58AM (UTC+0700)

Submission ID: 1352856975

File name: ding_SIMULATION_OF_RADIO_TELEMETRY_FOR_HOME_RANGE_PREDICTING.pdf (180.96K)

Word count: 1793

Character count: 9841



ICMSE 2015

INTERNATIONAL CONFERENCE ON MATHEMATICS, SCIENCE, AND EDUCATION
Faculty of Mathematics and Natural Science Semarang State University
phone: +62 248508112, +62 85876267947, +62 81805844994
email: icmse2015@gmail.com
website: icmse.unnes.com

SIMULATION OF RADIO TELEMETRY FOR HOME RANGE PREDICTING OF WREATHED HORNBILL (*RHYTICEROS UDULATUS*) ON MOUNT UNGARAN

Margareta R, Siti Alimah, Misbahul Munir¹

¹Department of Biology, Faculty of Mathematics and Natural Science, Semarang State University
Email : elak_sigid@mail.unnes.ac.id

ABSTRACT

Dynamics data of home range and utilization of the Wreathed Hornbill (*Rhyticeros undulatus*) habitat in Mount Ungaran be required. The objective of the study was to determine the models of transmitter and receiver to be used in determining the home range area of Wreathed Hornbill in Mount Ungaran Central Java. Time simulation implemented starting in April-August 2015. Simulations step include the identification of specifications and models of radio transmitter, antennas and receivers type are used, matching the wave, manufacture coat (jacket) for telemetry and try the ability, endurance telemetry which is attachment on the specimen in the Mangkang zoo. The result of identification of specifications and models of radio telemetry used the product Holohil System Ltd RI-2C transmitter, the active battery 3 (three) years with the 150000-151999 MHz frequency range. Yagi antenna (Yagi Antenna VHF Perdix 151 MHz) used type whip the product of Perdix Wildlife supplies, while the receiver used Yaesu FT 1802 M E.

Keyword : Wreathed Hornbill (*Rhyticeros undulatus*), Radio telemetry, simulation, home range

INTRODUCTION

Habitat structure and resource availability are important factors that influence bird communities in both naturally occurring and man-made environments (Chiang et al 2012). The conversion of natural habitats to agriculture, plantation, industry, grazing, illegal logging and urbanization has been cited as causes of declines in bird species.

Wreathed hornbills (*Rhyticeros undulatus*) is the bird species of Bucerotidae family that we can found in Mount Ungaran and using of natural forest for their habitat. The conservation status of Wreathed Hornbill has been category in the Least Concern (LC) (IUCN 2014) and based on CITES (Convention on International Trade of Endangered Species of Wild Fauna and Flora), the status belongs to the second appendix (Soehartono and Mardiasiti 2003). Hornbills species including Wreathed hornbill have a very important role in the Mount Ungaran forest as seed dispersers. Population of Wreathed Hornbill on Mount Ungaran is tendency threatened by habitat loss because of illegal logging, fragmentation, local hunting pressure, and also trading of bird. Previous study showed that the natural forest of Mount Ungaran a suitable habitat for Wreathed hornbill (Rahayuningsih and Nugroho 2013, Rahayuningsih and Nugroho 2015). Existence of Wreathed Hornbill in the habitat will support the balance of forest ecosystems and must be

support by conservation of forest environment. One of the efforts to provide a foundation of conservation strategy is through the study of ecology and their habitat. However, no empirical studies of Wreathed Hornbills home ranges and habitat use in Mount Ungaran exist. To estimate the home range and habitat use of Wreathed hornbills, we using radio telemetry method. Radio telemetry is a technique as the transmission of information from a transmitter transmitter on a free-ranging wild animal to a receiver. Radio telemetry it possible for determining bird movements over home range in size, breeding area, territories of resident bird species, habitat use, also to the movement patterns of international migratory species. Despite its popularity, radio-telemetry is inappropriate under many circumstances. It is an expensive and time-consuming technique which has proven to be unsuitable for use in some species (Andrusiak et al 1998). In Indonesia Radio telemetry methods have been used on Javan Hawk Eagle, (Afianto et al 1999, Gjersha et al 2004). We have planning Radio telemetry study for three years (2015-2017) with Wreathed Hornbills as a target species. Before the study begin in Mount Ungaran on Oktober-2015, we need simulation the method to ensure which the type of Radio telemetry that we can use. The objective of the study was to determine and simulation the type of transmitter and receiver to be used in the home range area of Wreathed Hornbill in Mount Ungaran Central Java.

METHOD

The early study for simulation start on April-August 2015. The location for simulation at Semarang state University (SSU), Mangkang Zoo, and Mount Ungaran. First step, we have been done to apply letter permission to arrest Wreathed Hornbills to BKSDA (Balai Konservasi Sumberdaya Alam) Central Java, cooperation with LIPI (Lembaga Ilmu Pengetahuan Indonesia) and IBBS (Indonesia Bird Banding Scheme). The data that we need are : identifying specifications and types radiotelemetry, antenna and receiver is used, matching the wave, manufacture coat (jacket) for telemetry, and try ability, endurance telemetry which is mounted on the specimen at the zoo Mangkang and Mount Ungaran..

RESULT AND DISCUSSION

To find the transmitter most suitable for Wreathed Hornbill we are looking for some products through internet sites, one of which was issued by HSL (Holohill Systems Ltd) "Transmitter". The first step to find the appropriate transmitter is the body size and weight of the bird. Because the body size and weight of Wreathed hornbill estimated to be the type of raptor, then we search the type of transmitter on raptor group. Conventional transmitters consist of an antenna, a power source and a transmitter unit. The transmitter model for raptor group on HSL are BD2, PD2, R1-2B, R2-2C, R12A, S1-2B, and A1-2B (Table 1 Figure 1). Then we look for the type of transmitter that is more specific and tailored to the purpose of research.

Table 1. Transmitter model for Raptor group (HSL)

Model	Attachment method	Weight (gram)	Life (weeks)	Antenna type
BD2	Backpack/tailmount	0.62-1.95	1-20	Whip
PD2	Backpack/tailmount	2.0-4.0	14-26	Whip
R1-2B	Backpack	5-14	26-100	Whip
R1-2C	Backpack/tailmount	5-21	25-250	Whip
R1-2A	Backpack	10-20	40-100	Whip
S1-2B	Backpack	12-16	26-100	Whip
A1-2B	Backpack	19-13	50-250	Whip



Frequency range : 138 to 235 MHz
Transmitter : Crystal controlle two stage design, pulse by CMOS multivibrator
pulse widyh & Rate: 20 to 24 ms, nominal 0.6 p/s (36 p/m)

Figure 1. R1-2C transmitter model

The result of identification specifications and radio telemetri model from raptor goup used is the product Holohil System Ltd Ri2C transmitter models, with about 10 grams and the active period of the battery 3 (three) years grams. We used 6 R1-C2 with the different number (Table 2, Figure 2). This transmitter was designed for tail mount attachment on raptors and other birds with robust tails. It can also be configured for backpack attachment to birds, herptiles and small mammals (HTL).

Table 2. R1-2C Transmitter model for Wreathed Hornbill

Model	Serial number	Freq. (MHz)	Pulse	Attachment Methode	Antena type
RI-2C	201043	150.019	0.63	Backpack,	Whip
RI-2C	201044	150.239	0.63	Backpack	Whip
RI-2C	201045	150.319	0.67	Backpack,	Whip
RI-2C	201046	150.470	0.65	Backpack,	Whip
RI-2C	201047	150.661	0.63	Backpack,	Whip
RI-2C	201048	150.880	0.62	Backpack,	Whip



Figure 2 R1-2C transmitter models by HSL with the different number

The antennas type that we used also Whip type, while receivers are used Yaetsu mkl1 the FT 690 mhz frequency 150000-151999 (Figure 3 and 4).



Figure 3 Antenna Yagi 151 MHz



Figure 4 Radio receiver Yaesu FT-1802 M/E

The next step was to coordinate the member of team researchers, also with LIPI and RCS (Raptor Conservation Society) to conduct stabilization team through theoretical training and simulation in the Semarang state University area (Figure 5), mangkang Zoo, and then directly in the field study.



Figure 5 Simulation radio transmitter in SSU

The training have been done primarily to familiarize team members for distinguish types captured signal receiver, the other team members are required to recognize individuals who have been given a trial installation of radio transmitter.

Trial installation of the transmitter have been done on Wreathed Hornbill specimens in Mangkang Zoo. The transmitter placed in the backpack of the bird for one week (Figure 6 and 7).



Figure 5 To setting radio transmitter on backpack




Figure 6 Radio telemetri backpack attachment method

The test results showed that a transmitter attachment on the backpack has a strong resistance. So that even the birds had several times tried to pull, strong enough transmitter attached and not damaged. After one week simulation, originally a bird disturbed so that it always seeks probe backs and wings to try to take it off, but the next day had not demonstrated such behavior.

The results of trials conducted in the wave reception Mount Ungaran also showed good results, the transmitter can be detected and monitored through a radio receiver that has been configured radio waves.

CONCLUSION

The model and specification radio telemetry are RI-2C transmitters, receivers Yaesu FT 690 mkl with 150000-151999 mhz frequency and Whip antenna type can be mounted and used on Wreathed Hornbill Mount Ungaran.

ACKNOWLEDGMENT

The authors would like to thank to the Directorate General of Higher Education and Culture for funding the research. The expression of thanks also addressed to the Association of Natural Conservation and Indonesian Institute Science and Perhutani for giving the research license. Mr. Hidayat Ashari and Mr Usep Suparman for helped during the training.

REFERENCES

- Afianto A, Jarwadi H, Dewi M. Aplikasi penggunaan radiotelemetry pada pendugaan karakteristik wilayah jelajah Elang Jawa (*Spizaetus bartelsi*) di Gunung Salak Jawa Barat. disampaikan pada seminar sehari Penerapan Sistem Informasi Geografi dan RadioTracking Untuk Pengelolaan Keanekaragaman Hayati, IPB-Darmaga, 26 Oktober 1999
- Andrusiak, Eric Walters, Eliot Terry and Keith Simpson. 1998. *Wildlife Radio-telemetry*. Standards for Components of British Columbia's Biodiversity

- No. 5. The Province of British Columbia Published by the Resources Inventory Committee.
- Chiang, S. N., P. H. Bloom, A. M. Bartuszevige, and S. E. Thomas. 2012. Home range and habitat use of Cooper's Hawks in urban and natural areas. Online (www.ucpress.edu/go/sab) in C. A. Lepczyk and P. S. Warren (editors). Urban bird ecology and conservation. *Studies in Avian Biology* (no. 45), University of California Press, Berkeley, CA.
- Gjersha et al. 2004, Home range size of the Javan Hawk Eagle (*Spizaetus bartlesi*) estimated from direct observation and radiotelemetry . *J. Raptor Res.* 38(4):343-349
- IUCN. 2014. The IUCN Red List of Threatened Species. www.iucnredlist.org. [12 April 2014]
- Rahayuningsih M and Nugroho Edi K. The Distribution and Population of Whreathed Hornbill in Mount Ungaran Central Java, *International Journal of Environmental Science and Development*, Vol 5, pp 492-495, Oktober 2013
- Rahayuningsih M and Nugroho Edi K.. Spatial modelling of Wreathed Hornbill (*Aceros undulatus*) habitat in Mount Ungaran Cntral Java. *International Journal of Environmental Science and Development*, Vol 6 No 6, pp 474-477, Oktober 2013
- Soehartono, T, Mardiasuti A. Pelaksanaan konvensi CITES di Indonesia. Jakarta : Japan International Cooperation Agency (JICA, 2003.

SIMULATION OF RADIO TELEMTRY FOR HOME RANGE PREDICTING OF WREATHED HORNBILL (RHYTICEROS UDULATUS) ON MOUNT UNGARAN

ORIGINALITY REPORT

11%

SIMILARITY INDEX

6%

INTERNET SOURCES

6%

PUBLICATIONS

4%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

5%
★ M Rahayuningsih, NE Kartijomo, A Retnaningsih, M
Munir, J Dahlan. " Nest Records of Wreathed Hornbill (
) in Gunung Gentong Station, Mount Ungaran Central
Java ", Journal of Physics: Conference Series, 2017

Publication

Exclude quotes Off
Exclude bibliography On

Exclude matches < 15 words