

Determination of Oviposition, pH, and Salinity of *Aedes aegypti*'s Breeding Places in Semarang Regency.pdf

by Widya Hary Cahyati

Submission date: 08-Feb-2021 12:46PM (UTC+0700)

Submission ID: 1504264503

File name: Determination of Oviposition, pH, and Salinity of *Aedes aegypti*'s Breeding Places in Semarang Regency.pdf (201.34K)

Word count: 5217

Character count: 24390



Determination of Oviposition, pH, and Salinity of *Aedes aegypti*'s Breeding Places in Semarang Regency

Widya Hary Cahyati[✉], Nur Siyam

Jurusan Ilmu Kesehatan Masyarakat, FIK, Universitas Negeri Semarang

Article Info

Article History:
Submitted March 2019
Accepted September 2019
Published November 2019

Keywords:
Oviposition,
Salinity, *Aedes aegypti*

DOI
<https://doi.org/10.15294/kemas.v15i2.21844>

Abstract

Dengue Hemorrhagic Fever (DHF) is still a health problem in Indonesia. This is because DHF can cause decease, and the incidence is always increasing. Vector control efforts in Semarang Regency can be done by knowing mosquito bionomics, so one of the actions can be taken is to modify the environment, so that the *Ae. aegypti* mosquito feels uncomfortable about our environment. This type of research is pure experimental. In this research, mosquito larva will be the subject to be breded using water with various degrees of acidity (pH) and salinity, then the ability to reproduce in various water with the pH and salinity levels is observed. The sample used was F1 larva originating from breeding places using ovitrap in Semarang Regency. Based on the results of the study showed that the most preferred container of the *Ae. Aegypti* mosquito for oviposition is a container made of plastic and can. The most optimal water pH for instar 2 larva breeding is water with a pH of 9, followed by water with a pH of 8 and 7. The most optimal water salinity for instar 2 larva breeding is water with a 0-6gr / 1 salinity.

Introduction

Dengue Hemorrhagic Fever (DHF) is still a health problem in Indonesia. This is because DHF can cause decease, and the incidence is always increasing. DHF is one of the diseases caused by dengue virus. This virus can be transmitted from patient to others through mosquito bites from the genus *Aedes*. The main vector of DHF is *Ae. Aegypti* mosquito, while *Ae. Albopictus* is a secondary vector. According to the Republic of Indonesia Health Office, starting in 2014, the number of DHF sufferers in Indonesia had increased. This can be seen from the Indonesian Health Profile which stated that in 2015 there were 129,650 dengue cases in Indonesia, with a total of 1,071 deceases. The increase in dengue cases cannot be separated from the effect of increasing population mobility, which is also accompanied by the spread of mosquitoes which become dengue vectors (infectious vectors) that are

increasingly spread in Indonesia (Depkes RI, 2005).

Prevention of DHF can be done by increasing public knowledge about DHF, for example through simple counseling activities or by avoiding contact with the dengue vector, namely *Ae. Aegypti* mosquito, including by mosquito nets application, closing house vents with wire mesh, and using mosquito repellent (Sayono et al., 2011). Vector control efforts in Semarang Regency can be done by knowing mosquito bionomics, so actions can be done by modifying the environment, so that the mosquito feels uncomfortable about our environment. Thus, gradually it will leave our environment, and we will be free from the risk of several diseases that can be transmitted by *Ae. Aegypti* mosquito. Some research also informs that insecticide resistance has occurred in several countries.

One of the efforts of the Semarang

[✉] Correspondence Address:
Jurusan Ilmu Kesehatan Masyarakat, FIK, Universitas Negeri Semarang
Email: widyahary27@mail.unnes.ac.id

Regency Health Office in controlling DHF is to use fogging. The active ingredient used in the DHF control fogging in Semarang Regency is sipermetrin 19: 1 since 2013 until now. The use of insecticides with the same type in a long time continuously will run the risk of resistance. This encourages the promotion of environmental modification measures that are considered safest in controlling mosquito population (Yotopranto et al., 1998). But the weakness in this activity is that it requires quite high community participation to carry it out. It is expected that the results of this research can be used as references in environmental modification activities to control *Ae. aegypti* mosquito population.

Method

This research type was pure experimental. In this research, mosquito larva would be the subject of treatment using water with various degrees of acidity (pH) and salinity, then the ability to reproduce is observed. The sample used was F1 larva originating from breeding places taken by larva capture using ovitrap in Semarang Regency. During the treatment, the room temperature was controlled by recording the temperature (morning, afternoon, evening), and keeping it in optimal condition for mosquito.

The stages of this research began by preparing the *Aedes aegypti* mosquito. Researchers installed ovitrap at the study site for a week. Next, the mosquito eggs that had been attached to the filter paper were then taken. Eggs taken from the location to the laboratory for rearing. The eggs would turn into larva and then turned into pupae in about 7 days. The larva were fed liquid sugar, pet food or fish food, and occasionally blood. Eggs that hatched into larva were transferred to a tray that contained 2 liters of water. Larva density in trays was estimated to range from 0.5 to 1 larva / cm³ with 2.5 cm water depth. Living larva were fed 0.5 gram of chicken liver from day 1 to 5 and the next day larva were fed as much as 1 gram. The pupa would turn into an adult

mosquito for about 3 days. After becoming an adult mosquito, the male and female mosquito will mate, so the female mosquito would lay eggs. The eggs from the mating would then be hatched, the mosquitoes that hatch were F1 offspring and would be tested for their ability to breed under various conditions of pH and water salinity. This research used *Aedes aegypti* mosquitoes from Semarang Regency. The sample in this study was a number of *Aedes aegypti* eggs taken randomly from the *Aedes aegypti* mosquito population from 18 villages / sub-districts in Semarang Regency.

Sampling was done by installing an egg trap (ovitrap) and larva examination because the number of *Aedes aegypti* mosquitoes in each village cannot be known certainly. The number of ovitraps placed were 100 ovitraps and done randomly. Ovitrap installation distance was 100-600 meters to avoid eggs from the same mosquito and placed in every randomly chosen house. The collection of *Aedes aegypti* mosquito eggs used ovitraps installed in every home, inside or outside the house. After the sample was bred to F1, and the age of the mosquito reaches 2-3 days stepping on the adult stage can fly, then the sample was taken using an aspirator. The data collected were used as primary data and obtained from the calculation of the number of *Aedes aegypti* mosquito deaths during the research. Data analysis was performed descriptively to see how the ability of 2nd instar larva to become mosquitoes under various conditions of pH and salinity of water.

Result and Discussion

The sample in this study used *Ae. aegypti* larva instar 3, each as many as 25 larvae for each treatment. Previously, researchers conducted mosquito rearing starting from capturing larva in nature using ovitrap, then bred in a laboratory. After the larva develop into adult mosquitoes, then both female and male *Ae. aegypti* who have almost the same age were

Table 1. Room Temperature During Experiment of Mosquito Breedstock Place Tendency

Measurement Time	Temperatur on Experiment Day (°C)													
	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Morning	26	26	26	25	25	26	26	26	26	26	25	26	26	26
Noon	28	28	28	28	27	28	28	28	28	28	28	28	28	28
Afternoon	27	27	27	27	26	27	27	27	27	27	27	27	27	28

Source : Primary Data, 2019

Table 2. Distribution of *Ae. Aegypti* Breeding Place Tendency

Breeding Container	Data taking	Number of Pupa found on Day-													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
Plastic	1	-	-	5	43	21	5	4	-	-	-	-	-	-	
	2	-	-	3	38	60	9	8	1	-	-	-	-	-	
	3	-	-	4	52	43	14	4	1	1	-	-	-	-	
	4	-	-	-	34	57	15	8	-	1	-	-	-	-	
	Avg	-	-	3	41,8	45,3	10,8	6	0,5	0,5	-	-	-	-	
Tin	1	-	-	3	35	15	5	5	-	-	-	-	-	-	
	2	-	-	-	42	17	9	2	1	-	-	-	-	-	
	3	-	-	2	21	23	7	12	-	2	-	-	-	-	
	4	-	-	-	26	24	11	5	-	1	-	-	-	-	
	Avg	-	-	1,3	31	19,8	8	6	0,3	0,8	-	-	-	-	
Soil	1	-	-	-	-	-	1	1	-	-	-	-	-	-	
	2	-	-	-	-	-	1	-	-	-	-	-	-	-	
	3	-	-	-	-	-	-	1	-	-	-	-	-	-	
	4	-	-	-	-	-	-	-	-	-	-	-	-	-	
	Avg	-	-	-	-	-	0,5	0,5	-	-	-	-	-	-	
Plastic with soil bed	1	-	-	5	20	11	2	3	-	-	-	-	-	-	
	2	-	-	-	13	8	4	-	2	1	-	-	-	-	
	3	-	-	-	16	13	9	4	-	2	-	-	-	-	
	4	-	-	-	7	11	12	3	1	-	-	-	-	-	
	Avg	-	-	1,3	14	10,8	6,8	2,5	0,8	0,8	-	-	-	-	
Tin with soil bed	1	-	-	-	24	9	2	-	-	-	-	-	-	-	
	2	-	-	-	12	8	-	2	-	-	-	-	-	-	
	3	-	-	-	9	11	2	-	-	-	-	-	-	-	
	4	-	-	-	16	5	7	-	-	-	-	-	-	-	
	Avg	-	-	-	15,3	8,3	2,8	0,5	-	-	-	-	-	-	

Source : Primary Data, 2019

put in a mosquito cage, and then mated. One to two days after copulation, female mosquitoes and male mosquitoes were separated. An easy way to distinguish *Ae. aegypti* female and male mosquitoes was by sticking a clean hand (without smearing anything, and does not smell anything) to the side of the cage. If a mosquito approached the hand, it was most likely that it was female, because it was trying to find

blood to ripen its eggs. Another way is to look at the special characteristics of the mosquito's body. Generally female is bigger than male. You can also see the mosquito palpus, which is the female is shorter than male. The tip of the female mosquito's abdomen also looks darker colored and pointed edges.

Female *Ae. aegypti* was taken using an aspirator, and then separated into other cages.

Table 3. Development of *Ae. aegypti* Mosquito Larva in various pH condition

Water pH	Data taking	Number of Larva becoming Mosquito	Percentage (%)
3	1	0	0
	2	0	0
	3	0	0
	4	0	0
	Average	0	0
4	1	11	44
	2	9	36
	3	9	36
	4	10	40
	Average	9,75	39
5	1	18	72
	2	15	60
	3	16	64
	4	17	68
	Average	16,5	66
6	1	21	84
	2	18	72
	3	17	68
	4	20	80
	Average	19	76
7	1	21	84
	2	19	76
	3	22	88
	4	19	76
	Average	20,3	81
8	1	25	100
	2	25	100
	3	22	88
	4	21	84
	Average	23,3	93
9	1	24	96
	2	24	96
	3	23	92
	4	25	100
	Average	24	96
10	1	20	80
	2	16	64
	3	18	72
	4	20	80
	Average	18,5	74

511	1	0	0
	2	1	4
	3	1	4
	4	0	0
	Average	0,5	2
12	1	0	0
	2	0	0
	3	0	0
	4	0	0
	Average	0	0

Source: Primary Data, 2019

The female mosquito separated from the male was then fed with guinea pig blood. Guinea pig that would be put into mosquito cage was slightly scraped off their feathers first, so that mosquito can easily suck blood. After 2 hours, the guinea pig was removed from the mosquito cage, and the Female *Ae. aegypti* had now become gravid 1, by looking at a mosquito's stomach which becomes bigger and forms like a blood sac. It was then taken to be put into a cage that had been prepared for oviposition.

In research, the tendency of *Ae. Aegypti* mosquito eggs placement was carried out for 14 days, this is because after 14 days all mosquitos have died. The number of *Ae. aegypti* female mosquito samples in this experiment were 25 animals. During the experiment, the room temperature was measured and managed in accordance with the optimal temperature for mosquitoes.

From table 1, it can be seen that the temperature in the laboratory was very optimal for mosquito breeding, where *Ae. aegypti* can develop optimally at a temperature of 25-27°C, but *Ae. aegypti* can live normally at temperatures of 20-30°C. Thus, the temperature factor did not affect the tendency of mosquito breeding places.

To see the tendency of laying *Ae. aegypti* mosquito eggs on various types of container materials / water containers, then an experiment was conducted by placing 5 kinds of containers in a mosquito cage containing 25 gravid female mosquitoes. The container was filled with well water. The tendency of laying eggs was seen from the number of larva contained in the container. The larva found were then

exterminated and thrown away, so as not to mix with ones that hatch later, so as not to confuse the calculation. Based on table 2. it can be seen that the tendency of *Ae. aegypti* mosquito for laying eggs is in a plastic container. It can be seen that the average total number of larva found in plastic containers was 108, followed by cans of 67. For containers made of soil, on average only one larva was found.

To see the development of mosquito in water with various pH conditions, an experiment was carried out using instar 2 larva. The instar 2 larva were obtained from *Ae. aegypti* mosquito female gravid 1 which had been separated from male mosquito. Then the hatched eggs were observed until they became instar 2 larva. These instar 2 larva were put into containers containing water with a different pH, and each container was put 25 instar 2 larva mosquito. The reason of Instar 2 chosen for this experiment was because Instar 1 was still too weak, so there was a higher risk of dying. instar 3 and 4 were not chosen because in this research we wanted to know the maximum breeding of mosquito. So by choosing instar 2, the observation phase could be longer. The results of this larva breeding can be seen in table 3. below.

From table 3, it could be seen that largest percentage of instar 2 larva that successfully hatch into mosquito were larva placed in water with a pH of 9, which was as much as 96%, followed by larva placed in water with a pH of 8, which is as much as 93%, and larva placed in water with a pH of 7, which is as much as 81%. Instar 2 larva were placed in water with a pH of 3 and a pH of 12 no larva have hatched into

mosquito.

To see the development of mosquitoes in water with various salinity conditions, an experiment was carried out using instar 2 larva. The larva were obtained from gravid 1 *ae. aegypti* female mosquito which had been separated from male mosquito. Then the hatched eggs were observed until they become instar 2 larva. These larva were put into containers containing water with different salinity, and each container was put 25 larva. The reason Instar 2 was chosen for this experiment was because Instar 1 is still too weak, so there is a higher risk of dying. As for instar 3 and 4 were not chosen because in this study we wanted to know the maximum breeding of mosquito, so by choosing instar 2, the observation phase could be longer. The results of this breeding larva can be seen in table 4 following.

Based on table 4. it can be seen that the largest percentage of instar 2 larva that successfully hatched into mosquito were larva placed in water with a salinity of 0, which was as much as 90%, followed by larva placed in water with a pH of 8, which was as much as 93%, and larva placed in water with salinity 4, which was as much as 81%. Instar 2 larva placed in water with salinity 7 and salinity 8, none successfully hatched into mosquito, although in salinity 7 there were 3 larva that successfully become pupae, but died before becoming mosquitoes.

From the results it can be seen that the most preferred container by the *Ae.* for oviposition was one made of plastic and can. In this research, the tendency of *Ae. aegypti* mosquito for laying eggs was in a plastic container. It can be seen that the average total number of larva found in plastic containers was 108, followed by cans with 67. For container made of soil, on average only one larva was found. In theory, *Ae. aegypti* mosquito will lay the eggs slightly above the surface of the clear water. In addition, it will avoid water where the wall is made of soil (Budiyanto, 2012; Schneider, 2011). But in certain circumstances, as in this research, *Ae. aegypti* mosquito wants to lay their eggs in a container made of soil or a plastic container that has soil bed, although the percentage is very small when compared to the number of eggs laid in a plastic container or can (Baserra et al., 2010).

Mature *Ae. aegypti* who have just come out of the pupa shell will remain for a while in the pupa shell to dry their wings. After the wings are felt dry, then *Ae. aegypti* immediately flies. To ripen the eggs, mature female *Ae. aegypti* must suck blood as food, while the male only eats fruit and flower fluids. After copulation, female *Aedes aegypti* will suck blood and three days later it will lay about 125 eggs and an average of 100 eggs, then it will suck blood again.

The results of this research are in accordance with the research of Hendri et al., (2010) which states that of the 39 positive larva of containers, the highest number is found in plastic-based buckets. The attractiveness of female mosquito to lay their eggs is affected by the color of the container, temperature, humidity, light, and environmental conditions. Other factors that also affect the tendency of female mosquitoes to lay their eggs include the ability to absorb water, texture, and color of the material. In Buenos Aires, Argentina, it was found that containers of black and black plastic materials contained a lot of *Ae. aegypti* larva (82.1%). Based on the results of these studies it can be seen that mosquito in each particular region have tendency in different water reservoirs, in the type, basic material, and color used. These factors are estimated to affect the percentage of larval acquisition in various containers in each region (CDC, 2016).

In this research, *Ae. aegypti* mosquito larva was placed in regulated water, so that it had a pH between 3 - 12. The pH range was chosen because we wanted to know whether the larva can adapt to pH conditions that are rather extreme and capable of breeding. In previous studies, no results had yet been obtained that *Ae. mosquito* larva was able to breed at pH less than 4 or more than 10. According to the research of Sayono et al., (2011), eggs from oviposition of gravid I female *Ae. aegypti* can be found in water with a pH range of 4-10 in the attached ovitrap, and have different results in each treatment. Research conducted by Yulidar (2015) stated that the hatchability of *Ae. Aegypti* eggs at the laboratory scale reached 80.09%. Eggs produced by *Ae. Aegypti* mosquito were then collected to be hatched. In this research, gravid female mosquito can lay their eggs in

Table 4. Development of *Ae. aegypti* Mosquito Larva in Various Salinity Condition

Water (gr/l)	Salinity	Data taking	Number of Larva becoming Mosquito	Percentage (%)
0		1	23	92
		2	23	92
		3	23	92
		4	21	84
		Average	22,5	90
4		1	21	84
		2	19	76
		3	21	84
		4	20	80
		Average	20,3	81
5		1	18	72
		2	18	72
		3	16	64
		4	19	76
		Average	17,8	71
6		1	12	48
		2	8	32
		3	7	28
		4	8	32
		Average	8,8	35
7		1	0	0
		2	0	0
		3	0	0
		4	0	0
		Average	0	0
8		1	0	0
		2	0	0
		3	0	0
		4	0	0
		Average	0	0

Source: Primary Data, 2019

acidic pH 3 conditions. But after being hatched, only a small number that managed to hatch into larva. The hatched larva were only able to survive less than 24 hours, then died (mosquito larva cannot develop). This might be caused by environmental conditions that were too acidic. Dead larva were white. Water pH is one of the factors that affect larval development. In

research conducted by Clark (2004), it stated that mosquito larva will die at pH <3 or >12. Based on the result of research conducted by Sayono et al., (2011), instar III larva cannot develop into pupa at pH 4. But the results of the research are different from this one, because in this research, the development of larva into adult mosquito can occur in water with pH 4.

¹ Based on the results of the research it can be seen that the most optimal water pH for breeding instar 2 larva into *Ae. aegypti* mosquitoes is water with a pH of 9, followed by water with a pH of 8 and 7. One important factors that can affect the percentage of hatching of *Ae. Aegypti* mosquito egg is temperature and pH. In addition, the process of developing an egg to the next stage (becoming an adult stage) is also affected by temperature and pH. If the temperature and pH factors are at optimal conditions, the process of mosquito breeding will also be higher. Vice versa, if the temperature is too hot or cold, it will interfere with mosquito growth, even stop or kill the mosquito (Agustin et al., 2017; Wahyuningsih, 2009).

Based on the results of Jacob's research (2014), *Ae. aegypti* cannot develop in alkaline water, and can develop optimally in neutral water. However, this research shows that the biggest percentage is *Ae. aegypti* can develop into adult mosquito at pH 9. That is, based on this research, the most optimal mosquito larva develop in more alkaline water (pH 9) than in neutral water (pH 7). The ability of *Ae. aegypti* larva to develop in water conditions with an alkaline pH due to it's adaptability in various environmental conditions (CDC, 2016).

The chemical condition of water that is acidic or alkaline does not affect gravid female mosquito to lay their eggs (Agustina, 2013). That is, if there is a pool of water that is considered safe by the mosquito to lay its eggs, then that's where the gravid female mosquito will lay her eggs. This is consistent with the results of research Tilak et al., (2005) which states that the pH of water does not affect the oviposition response of female mosquito. However, the pH of water will affect the ability of larva to develop into adult mosquito. The results of this study are in line with research of Sayono (2011) which states that the *Ae. aegypti* mosquito can grow and develop well in water drains

that are still. This is because the pH of the sewage water itself is not all neutral, because the water comes from various sources of waste water.

In this research, the largest percentage of instar 2 larva that successfully hatched into mosquitoes were larva placed in water with a pH of 9, which was as much as 96%, followed by one placed in water with a pH of 8, which was as much as 93%, and larva placed in water with a pH of 7, which is as much as 81%. Instar 2 larva placed in water with a pH of 3 and a pH of 12 could not hatch into mosquito.

In the condition of water salinity 0 gr / l up to 6 gr / l, *Ae. aegypti* larva is able to develop into an adult mosquito. When the salinity of water is 6 g / l, the mosquito larva developed some appear to be larger than usual size, where the larval instar IV size is 5 mm. This can occur because the liquid absorbed the body of the larva, so the size of the larva appear larger than normal size (CDC, 2016). In Clark's research (2004) stated that *Ae. aegypti* develops in water salinity with a concentration of 3.5 gr/l.

Mortality of *Ae. aegypti* larva in various water salinity in the laboratory in the first 24 hours occurred in the treatment with salinity of 18 gr / l - 22 gr / l. High concentration of NaCl resulted in an imbalance between the body fluids of larva and brood media. This difference in osmotic pressure results in the death of *Ae. aegypti* larva. Osmosis plays an important role in the body of a living creature, because the fluid in the body of the larva is pulled out so that the body of the dead larva shrinks and is damaged called crenation (CDC, 2011). This is in accordance with research by Badrah & Nurul (2011) which stated that the content of Cl in water can interfere with the process of development and hatching of mosquito eggs. *Ae. Aegypti* larva can survive on salt levels at concentrations of 10.0 mg - 59.5 mg. When viewed from the length of the development of the larva to

adult mosquito, at a concentration of 6 gr / l some mosquito developed abnormally. The metamorphosis cycle of the mosquito exceeds the normally from egg to adult + 10 days (Ahmad et al., 2009; MOH RI, 2005). This has an impact on the size of larva and pupa in normal conditions. Larva and pupa that do not develop, will not survive and died.

In addition to temperature and pH, the presence of organic matter also affects the development process of mosquito, ranging from larva (pre-mature stage) to mature mosquito (Olayemi et al., 2010; Curto & Schweigmann., 2006). The growth of mosquito larva is divided into several instars, which then become pupa. The process is affected by the presence of detritus or organic matter as food. The media used in this research was clean water from well, which is used as a breeding ground for *Ae. aegypti* mosquito.. Usually well water has a sufficient amount of organic matter. The content of organic matter also affects the penetration of light and oxygen content in a medium. If the content of organic matter is too much, it can interfere with the penetration of light into the media so that it can interfere the development of pre-mature mosquito. So, too much organic matter can also inhibit the breeding of *Ae. aegypti* larva (Agustin et al., 2017).

The longevity of *Ae.aegypti* also depends on the temperatures, humidity of the air, water supply, food and predators (Wong et al., 2011). *Aedes aegypti* can live at 20°C with a humidity of 70%, but can only last for about 100 days. *Ae.aegypti* will die if it is at 6°C for 24 hours (Yulidar & Wilya, 2015)

Based on the results of the research it can be seen that the most optimal water salinity for instar 2 larva breeding into *Ae. aegypti* mosquito is water with a 0-6gr / l salinity.

Conclusion

The results of this research showed that

the most preferred container of the *Ae. Aegypti* mosquito for oviposition is a container made of plastic and can. The most optimal water pH for breeding instar 2 larva into *Ae. aegypti* mosquito is 9, followed by pH of 8 and 7. The optimal water salinity for breeding instar 2 larva into *Ae. Aegypti* mosquito is water with a 0-6gr / l salinity.

Reference

- Ahmad, I., Astari, S., Rahayu, R., & Hariani, N. (2009). Status Kerentanan *Aedes aegypti* (Diptera: Culicidae) pada Tahun 2006-2007 terhadap Malathion di Bandung, Jakarta, Surabaya, Palembang, dan Palu. *Biosfera*, 26 (2): 82-89.
- Agustina, E., 2013. Pengaruh Media Air Terpolusi Tanah terhadap Perkembangbiakan Nyamuk *Aedes aegypti*. *J Biotechnology*, 1: pp. 103-107.
- Agustin, Indira., Tarwotjo, U., & Rahadian, R., 2017. Perilaku Bertelur dan Siklus Hidup *Aedes aegypti* Pada Berbagai Media Air. *Jurnal Biologi*, 6(4): 71-81.
- Badrah, Sitti., Nurul Hidayah., 2011. Hubungan Antara Tempat Perindukan Nyamuk *Aedes Aegpty* dengan Kasus Demam Berdarah Dengue Di Kelurahan Penajam Kecamatan Penajam Kabupaten Penajam Paser Utara. *J. Trop. Pharm. Chem*, 1(2): 153-160.
- Baserra, R., Amador, M., & Clark, G.C., 2010. Competition and Resstance to Satrvation in Larva of Container Inhibiting *Aedes* Mosquitoes. *Ecologycal Entomology*, 5: 117-127.
- Budiyanto, A., 2012, Karakteristik Kontainer Terhadap Keberadaan Jentik *Aedes aegypti* di Sekolah Dasar. *Jurnal Pembangunan Manusia*, 6(1).
- Curto, S.I., & Schweigmann, N.J., 2006. Spatial Distribution Pattern of Oviposition in the Mosquito *Aedes aegypti* in Relation to Urbanization in Buenos Aires: Southern Fringe Bionomics of an Introduced Vector. *Medical and Veterinary Entomology*, 20(2): 209-218.
- CDC. 2016. *Entomology Dengue*. Diakses pada tanggal 26 April 2016. <http://www.cdc.gov/dengue/entomologyEcology/index.html>.
- Clark, Thomas M., Benjamin J. Flis, Susanna K. Remold. 2004. pH Tolerances and Regulatory Abilities of Freshwater and Euryhaline Aedine Mosquitoes Larva. *The Journal of Experimental Biology*. 207:2297-2304.
- Depkes RI. 2005. *Pencegahan Dan Penanggulangan Penyakit Demam Dengue Dan Berdarah*

- Dengue*. Ditjen PPM & PLP. Jakarta.
- Hendri, J., Roy, N.R.E.S., & Prasetyowati, H., 2010. Tempat Perkembangbiakan Nyamuk *Aedes* spp. Di Pasar Wisata Pangandaran. *Aspirator*, 2(1): 23-31.
- Jacob, Aprianto, D. Pijoh, Victor Wahongan., 2014. Ketaahanan Hidup dan Pertumbuhan Nyamuk *Aedes* spp pada berbagai Jenis Air Perindukan. *Jurnal e-Biomedik (eBM)*, 2(3).
- Olayemi, I. K., Omalu, I. C. J., Famotele, O. I., Shegna, S. P., & Idris, B., 2010. Distribution Of Mosquito Larva In Relation To Physico-Chemical Characteristics Of Breeding Habitats In Minna, North Central Nigeria. *Reviews in Infection*, 1(1): 49-53.
- Sayono, S. Qoniatun, Mifbakhuddin. 2011. Pertumbuhan Larva *Aedes aegypti* Pada Air Tercemar. *Jurnal Kesehatan Masyarakat Indonesia*. 7 (1).
- Schneider, J.R., Dave, D.C., Akio, M., Jeafne, R., David W.S., 2011. Heritability and Adaptive Phenotypic Plasticity of Adult Body Size in the Mosquito *Aedes aegypti* with Implications for Dengue Vector Competence. *J. Inf, Gen, Evol*, 11(1): 11-16.
- Tilak, DR Rina, Maj Vivek Gupta, Maj Vani Suryam, Mrs JD Yadav, Brig KK Dutta Gupta., 2005. A Laboratory Investigation into Oviposition Responses of *Aedes aegypti* to Some Common Household Substances and Water from Conspecific Larva. *MJAFI*. 61(3): 227-229.
- Wahyuningsih, E., 2009. Kefektifan Penggunaan Dua jenis Ovitrap untuk Pengambilan Contoh Telur *Aedes* sp. di Lapangan. *Jurnal Entomologi Indonesia Universitas Diponegoro*, 2.
- Wong, J., Stoddard, S.T., Aste, H., Morrison, C.A., & Scott, T.W., 2011. Oviposition Site Selection by the Dengue Vector *Aedes aegypti* and It's Implication for Dengue Control. *PLoS Neglected Tropical Disease*, 5: 12.
- Yotopranoto, S., Rosmanida S., & Sulaiman., 1998. Dinamika Populasi Vektor pada Lokasi dengan Kasus Demam Berdarah Dengue yang Tinggi di Kotamadya Surabaya. *Majalah Kedokteran Tropis Indonesia*, 9(1)
- Yulidar & Wilya, Veny., 2015. Siklus Hidup *Aedes aegypti* Pada Skala Laboratorium. *Sel*, 2(1): 22-28.

Determination of Oviposition, pH, and Salinity of Aedes aegypti's Breeding Places in Semarang Regency.pdf

ORIGINALITY REPORT

3%

SIMILARITY INDEX

%

INTERNET SOURCES

2%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | Submitted to Universitas Negeri Semarang
Student Paper | 1% |
| 2 | Submitted to Surabaya University
Student Paper | 1% |
| 3 | Malik Saepudin, Soeharyo Hadisaputro, Ari Suwondo, Suhartono Suhartono. "The Effect of Rekattidiri Ovitrap towards Aedes aegypti Larval Density", International Journal of Public Health Science (IJPHS), 2017
Publication | <1% |
| 4 | Submitted to Indiana State University
Student Paper | <1% |
| 5 | Y. C. Tsang. "Hunger motivation in gastrectomized rats.", Journal of Comparative Psychology, 1938
Publication | <1% |
| 6 | Submitted to Sriwijaya University
Student Paper | <1% |

7

Chou, Lee Yiung, Gary A. Dykes, Robyn F. Wilson, and Charles M. Clarke. "Nepenthes ampullaria (Nepenthaceae) Pitchers Are Unattractive to Gravid *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae)", *Environmental Entomology*, 2015.

Publication

<1%

8

Ely Hayati Nasution, Roswita Silalahi, Vivi Adryani Nasution. "FOREIGNIZATION AND DOMESTICATION IN THE MINISTERIAL WEBSITE LOCALIZATION", *JOALL (Journal of Applied Linguistics & Literature)*, 2019

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Determination of Oviposition, pH, and Salinity of Aedes aegypti's Breeding Places in Semarang Regency.pdf

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
