

Larvicidal and pupicidal activities from *Citrus hystrix* against *Aedes aegypti* mosquito

by Niken Subekti

Submission date: 25-Jul-2021 01:38PM (UTC+0700)

Submission ID: 1623659162

File name: EEC-58.pdf (92.36K)

Word count: 3939

Character count: 20398

Larvicidal and pupicidal activities from *Citrus hystrix* against *Aedes aegypti* mosquitoes

Niken Subekti^{1*}, Asni Puraedah¹, Dyah Rini Indriyanti¹ and Agoes Soegianto^{2*}

¹Biology Department, Faculty of Mathematics and Natural Sciences, Semarang State University, Indonesia

²Biology Department, Faculty of Sciences and Technology, Universitas Airlangga, Surabaya, Indonesia

(Received 4 April, 2020; Accepted 25 June, 2020)

ABSTRACT

Aedes aegypti is a dengue vector that is known to be very threatening to human's life. Dengue virus can cause dengue haemorrhagic fever (DHF) disease that are known to kill human beings. Control is done during this using abate. However, abate is synthetic insecticide that is not environmentally friendly. Therefore, the control of mosquitoes by using natural materials such as essential oil of *Citrus hystrix* one of the wise choice to mosquito control. This study aims to analyse the content of compounds in the essential oil of *Citrus hystrix* and analyse the mortality of larvae and pupa due to the provision of essential oil of kaffir lime based on nano emulsion technology. Essential oil is produced through the process of extracting *Citrus hystrix* using steam distillation, to determine the chemical compound of kaffir lime peel oil GC-MS was used, to make nano emulsion sonicator was used. Nano emulsion is then tested on larvae and pupa of *Aedes aegypti*. The identified major compounds were *beta-pinene* and *limonene*. The particle size identified from nano emulsion was ± 129.45 nm and ± 109.15 nm. The highest larvicidal activity was $21.6 \pm 1.67\%$ ($LC_{50} = 546.592$ ppm) and pupicidal activity $36 \pm 1.67\%$ ($LC_{50} = 3045.323$ ppm). Nano emulsion of essential oil of *Citrus hystrix* showed larvicidal and pupicidal activity on larva and pupa of *Aedes aegypti* and it is more environmentally friendly than abate. The results of this study can be recommended for control systems on *Aedes aegypti* mosquitoes.

Key words: *Aedes aegypti*, Essential oil, *Citrus hystrix*, Larvicidal, Pupicidal

21

Introduction

Aedes aegypti mosquito is the main vector of dengue virus. Dengue virus is very dangerous for humans. The virus can cause Dengue Haemorrhagic Fever (DHF), filariasis, yellow fever, malaria and various types of encephalitis (Waritkoo *et al.*, 2011). By 2016, the number of dengue cases in Indonesia is reported to reach 160.000 with the death of 10.000 (Ministry of Health of the Republic of Indonesia, 2020).

Efforts have been carried out by the Ministry of Health of the Republic of Indonesia in 2011

which includes: prevention, discovery, relief and reporting, epidemiological investigation and observation of dengue fever and counselling. However, due to lack of awareness and ignorance of people about dengue fever, dengue virus cannot be stopped. Therefore, *Aedes aegypti* mosquito control is required.

Generally, the mosquito control system is done using chemical pesticides. Chemical pesticides are pesticides that are not environmentally friendly and can harm other organisms (Nugroho, 2011). Therefore, it needs controlling technology from natural

materials that are environmentally friendly and do not harm other organisms and are effective in controlling mosquitoes.

Potential natural ingredients for mosquito control are kaffir lime peel (*Citrus hystrix*). Peel of Kaffir lime is underutilized. Characteristic from leave oil dominant are essential (-)-(S)-citronellal (80%), and citronellol (10%), nerolid limonene. *Citrus hystrix* is another citrus enantiomer dominant (4) (R)-citronellal. Some of the chemical compounds in the essential oil of kaffir lime peel are the limonene and β -pinene compounds (Kasuan *et al.*, 2011). The limonene and α -pinene compounds are known to have larvicidal activity (Akono *et al.*, 2015).

Essential oil have volatile and unstable properties (Noveriza *et al.*, 2017). Therefore, the essential oil of kaffir lime peel needs to be formulated in a more stable form, such as in the form of nano emulsion. In addition, the reduction of particle size in the form of nano emulsion of essential oil can also be effective as a larvicidal agent (Anjali *et al.* 2011). Nano emulsion has average droplet size is between 100 to 500 nm (Shah *et al.*, 2010). This latest technology becomes one of the solutions for economic problems and also environmental friendliness (Elkhal *et al.*, 2015). Small size particles are expected to improve the absorption of active ingredients quickly and deeply. The nanoemulsion formulation which is widely used is ultrasonic emulsification method because of its ease of use and economics (Sugumar *et al.*, 2014). The use of nanotechnology as a pesticide will reduce the toxic effects from synthetic pesticides.

Recently, control of vector-borne diseases was performed using nano emulsion from essential oil and was a good alternative against *Aedes aegypti* (Duarte *et al.*, 2015). This study was conducted to analyse the larvicidal and pupacidal activity of nano emulsion of *Citrus hystrix* oil against larvae and pupa mosquitoes. Instar III larvae and *Aedes aegypti* mosquito pupa were exposed for 24 hours at various concentrations. The small particle size of nano emulsion will make it the ideal larvicide and pupicide against *Aedes aegypti*.

Methodology

Collection and Processing of Plants (Othman *et al.*, 2016)

The material used in this study is the peel of *Citrus*

hystrix obtained from Citrus Plantation Semarang, Indonesia. Kaffir lime was chosen based on diameter, and color of the lime peel. Diameter of *Citrus hystrix* kaffir used is 5.0-7.5 cm.

Mosquito Rearing (Kumar *et al.*, 2010)

Mosquitoes were reared by inserting albino mice into mosquito coils. Female mosquitoes will suck the blood of albino rats for egg maturation. The breeding of *Aedes aegypti* mosquitoes was conducted at Biology Laboratory of Universitas Negeri Semarang at $28 \pm 1^\circ\text{C}$ and humidity $80 \pm 5\%$ RH. While for larvicide and pupicide, the eggs produced from rearing mosquitoes will be hatched and allowed to grow into instar larvae III and pupa.

Extraction of Essential Oils (Khandagle *et al.*, 2011)

The steam distillation process began by inserting 4 kg of dried lime peel of *Cytrus hystrix* into the extractor container. Then, the process was followed by heating the extractor by using steam generated from water heating. Furthermore, the essential oil was carried through the steam generated from the heating process then it went to condensation process. The essential oil produced was 54 ml. The essential oil was then stored in a small bottle at 4°C .

Gas Chromatography – Mass Spectroscopy (GC-MS) (Loh *et al.*, 2011)

Content analysis of the compound in the essential oil of kaffir lime peel used GC-MS. The GC-MS process method using Shimadzu QP 5050 which is a mass spectrometer system equipped with a capillary column of BPX 5 (film trapping $30\text{ m} \times 250\text{ m}, 0.25$). Helium is used as gas carrier. MS operating conditions are: ionization induced by electron (EI) at 70 eV, ion source 200°C . This compound is identified by performing a library search using Shimadzu NIST/ EPA/ NIH + mass spectral databases.

Nanoemulsion Preparation (Sugumar *et al.*, 2014)

Crude emulsion was prepared in proportion 16.66% oil: 16.66% tween 20: 66.66% water in (v/v) was made by stirring it using magnetic stirrer at 250 rpm for 10 minutes. The next process was ultrasonic emulsification by using 20 kHz sonicator (Ultrasonics, USA). The sonicator with a 13 mm probe diameter was immersed in a coarse emulsion. The temperature difference between the initial crude emulsions for the final nanoemulsion should be less than 10°C . The heat was generated during the high-en-

¹⁶ ergy ultrasonic emulsification process. This heat is reduced by the storage of emulsion samples in a beaker glass containing relatively larger ice.

Analysis of Nanoemulsion Size (Anjali et al., 2011)

¹¹ The size and morphology of nano emulsion particles of essential oil of kaffir lime peel (*Citrus hystrix*) can be determined by using a transmission electron-microscopy (TEM) device. For the use of this TEM tool, a drop of nano emulsion was placed on the copper grid and allowed to dry in a vacuum. This electron micrograph transmission used Tecnai G-10 (Philips) instrument, 80 kVTEM with W-search and a piece of ultrahigh-resolution pillar with point-to-point resolution 1.9Å.

Larvicidal Test (WHO, 2005)

Nano emulsions of *Cytrushystrix* were prepared at concentrations of 50, 100, 150, 200 and 250 ppm. Nano emulsions of kaffir lime peel with various concentrations respectively were tested on 25 instar larvae III of *Aedes aegypti*. The controls in this study were 250 mL of water. The rate of larvicidal mortality was assessed after 24 hours. Larvae are considered dead if the larvae cannot move and cannot transfer the siphon to the water surface.

Pupicidal Test (Krishnappa et al., 2012)

Pupicide test by using experimental protocol was performed according to. Ten pupas of *Aedes aegypti* were taken into 500 mL nano emulsion solution of *Citrus hystrix* kaffir lime peel oil and abate with various concentration. The control was 500 mL of water. Bioassay at test concentrations of 62.5, 125, 250, 500, 1000 ppm (WHO, 2005). The study room is preferably at temperature (28 ± 2) °C with general photoperiod (12: 12h/ L:D). Pupa was considered dead when the pupa ¹⁴ cannot move when it is touched for many times with a soft brush. Mortality of each pupa was recorded after 24 hours of exposure to the selected concentration of chemical compositions following the Abbott formula.

Statistical analysis

⁷ The data were calculated by using the software statistical package of social science (SPSS) version 23.0 for windows. The significant ³² level was set at $P < 0.05$. Probit analysis SPSS 23.0 was applied to determine lethal concentrations causing 50% (LC_{50}) mortality of larvae and pupae 24h after treatment application.

25

Results and Discussion

Chemical Compounds of Essential Oil of *Citrus hystrix*

³⁹ Chemical compounds in essential oil of kaffir lime peel can be determined by Chromatography Gas-Mass Spectroscopy (GC-MS). There are 18 compounds identified. The highest peak of GC-MS chromatogram is peak 4 (β -pinene) and peak 9 (limonene). The β -pinene compound was seen at retention time of 8.142 minutes and by molecular weight 136. The presence of β -pinene compound in the essential oil of *Citrus hystrix* was 38.60%. The limonene compound was seen at retention time of 9.750 minutes and with molecular weight of 136. The presence of limonene compound in the *Citrus hystrix* oil was 15.46%. Limonene compounds are the second highest compound after β -pinene in the essential oil of *Citrus hystrix*. Both these compounds give a distinctive aroma. The limonene compound gives a strong orange scent ³ while β -pinene gives the woody-green scent in the essential oil of the kaffir lime peel (Kasuan et al., 2013). The identified compounds can be ²⁴ seen in Table 1.

Table 1 shows that the major compounds in the essential oil of *Citrus hystrix* are β -pinene (38.6%) and limonene (15.46%). This is confirmed by research

35

Table 1. Chemical compound of essential oil of *Cytrushystrix*

Peak	Compound	Retention Time	% of Essential Total
³³ 8	Bicyclo 3.1.0 hex-2-ene	6.617	0.7
2	Alpha-pinene	6.817	3.05
3	Champhene	7.233	0.22
4	Beta-pinene	8.142	38.6
5	Beta-myrcene	8.525	1.26
6	Alpha-phellandrene	8.933	0.41
7	Alpha-terpinene	9.325	3.18
8	Benzene	9.595	0.88
9	Limonene	9.75	15.46
10	Gamma-terpinene	10.692	5.12
11	Cis- Linalol oxide	11.142	2.53
12	(+)-2-Carene	11.658	2.4
13	6-Octenal, 3, 7-dimethyl	13.767	11.85
14	3-Cyclohexen	14.567	11.38
15	Copaene	20.4	0.84
16	Caryophyllene	21.625	0.63
17	Napthalene	24.342	1.33
18	9-Octadecenoid acid	36.975	0.15

from Kasuan *et al.* (2013) which shown that **11** limonene compounds (17.2%), β -pinene (13.5%) are the main compounds in essential oil of kaffir lime *Cytrus hystrix*. However, the presence of β -pinene compounds from the results of this study is greater. β -pinene and limonene compounds are compounds that have larvicidal activity (Akono *et al.*, 2015).

Morphology and Size of Nanoemulsion

Transmissions Electron Microscopy (TEM) analysis aims to find out particle size and morphology of nanoemulsion of *Cytrus hystrix* oil. The TEM test results revealed that by looking the nanoemulsion of *Cytrus hystrix* oil by microscope at 500 nm magnification, showed 2 nanoemulsion particles. Both particles of nanoemulsion of essential oil of *Cytrus hystrix* skim has a size of ± 129.45 nm and ± 109.15 nm. According to Shah *et al.* (2010), the average nanoemulsion size ranges from 20-200 nm. The visible nanoemulsion morphology is spheric in shape. Similar results were obtained from Anjali *et al.* (2011) which stated that the nanoemulsion results of Neem oil nanoemulsion are spheric in shape.

Larvicidal Activity of Nanoemulsion of *Cytrus hystrix* Oil

Nanoemulsion of essential oil of *Cytrus hystrix* was then tested on *Aedes aegypti* larvae. The larvicidal activity of nanoemulsion of kaffir lime peel oil and abate can be seen from larvae mortality calculated after 24 hours of treatment. The result of observation of the effect of nanoemulsion of essential oil of *Cytrus hystrix* and abate to mortality of larvae *Aedes aegypti* is presented in Table 2.

Based on Table 2, it can be seen that the highest percentage of mortality in nanoemulsion of essential

Table 2. Larvicidal activity of nanoemulsion of *Cytrus hystrix* oil and abate against larvae of *Aedes aegypti*

Concentration (ppm)	Average Mortality of larvae (%)*	
	Nanoemulsion	Abate
0	0 \pm 0.00 ^a	0 \pm 0.00 ^a
50	0 \pm 0.00 ^a	100 \pm 0.00 ^f
100	8 \pm 1.58 ^b	100 \pm 0.00 ^f
150	15.2 \pm 1.09 ^c	100 \pm 0.00 ^f
200	16.8 \pm 1.48 ^d	100 \pm 0.00 ^f
250	21.6 \pm 1.67 ^e	100 \pm 0.00 ^f

*Values are means \pm standard deviations from five replication

oil of *Cytrus hystrix* is 21.6 \pm 1.67% and abate has mortality 100 \pm 0.00%. No mortality was observed in the control groups. ANOVA test indicated statistical differences in mortality between tested concentrations after 24 hours (F value = 17.733; P = 0.000). According to Balasubramani *et al.* (2017), mortality of larvae of nanoemulsion *Vitex negundo* reach 80.66 \pm 0.66% for 24 hours of application. However, concentration of *Vitex negundo* nanoemulsion used is higher at 400 ppm. This condition resulted in higher mortality. Therefore, if the nanoemulsion concentration is increased then the mortality of the *Aedes aegypti* larvae will also increase. In addition, the results of study by Duarte *et al.* (2015) also confirm that the length of time of exposure may affect larval mortality. The mortality results from the essential oil of nanoemulsion of *Rosmarinus officinalis* reached mortality > 75%. The results are quite high because of the length of application from *Rosmarinus officinalis* which is up to 48 hours. Based on this fact, if this experiment is conducted for 48 hours, the mortality of the larvae will also increase.

The toxic potential of essential oils of *Cytrus hystrix* and their compounds against *Aedes aegypti* larvae may vary significantly according to extrinsic and intrinsic factors. Plant species, age of the plant, chemotypes, plant parts, and geographic conditions (such as season, humidity percentage, temperature, sunlight, rainfall, and altitude) in which the plant was collected, methods used and larval source, in general, induce different larval responses (Subekti *et al.*, 2018).

Additionally, the different ecological niches of mosquito larvae can influence their susceptibility to toxic compounds. It was hypothesized that the field collected larvae are more resistant to chemicals than those reared in laboratory because the former are better adapted to adjust to environmental variations and have a higher genetic variability.

Pupicidal Activity Nanoemulsion Essential Oil of *Cytrus hystrix*

Nanoemulsion of essential oil of *Cytrus hystrix* was then tested on *Aedes aegypti* pupae. The pupicidal activity of nanoemulsion essential oil of *Cytrus hystrix* and abate can be seen from larvae mortality calculated after 24 hours of treatment. The result of observation of the effect of nanoemulsion of essential oil of *Cytrus hystrix* and abate to mortality of pupae *Aedes aegypti* is presented in Table 3.

The result of the pupicide test in Table 3 shows

Table 3. Pupical activity of nanoemulsion essential oil of *Cytrus hystrix* and abate against pupa of *Aedes aegypti*

Concentration (ppm)	Pupae Mortality \pm SD		% Mortality \pm SD	
	Nanoemulsion	Abate	Nanoemulsion	Abate
Control	0 \pm 0.55	0 \pm 00	4 \pm 0.55	0 \pm 00
62.5	14.58 \pm 1.30	2 \pm 0.44	18 \pm 1.30	2 \pm 0.44
125	16.67 \pm 1.87	8 \pm 0.84	20 \pm 1.87	8 \pm 0.84
250	16.67 \pm 2.34	18 \pm 1.48	20 \pm 2.34	18 \pm 1.48
500	25 \pm 1.09	42 \pm 2.68	28 \pm 1.09	42 \pm 2.68
1000	33.34 \pm 1.67	50 \pm 1.87	36 \pm 1.67	50 \pm 1.87

that the highest mortality in the essential oil of *Citrus hystrix* was 36 \pm 1.67% and abate 50 \pm 1.67%. The mortality observed in the control groups was 4 \pm 0.55%. ANOVA test indicated statistical differences in mortality between tested concentrations after 24 h (F value = 2.296; P = 0.077). According to study of Priyadarshi *et al.* (2012) which was aimed at investigating the potential of silver nanoparticles synthesized from *Euphorbia hirta* to *Anopheles stephensi* showed the highest mortality 55.8% at 250 ppm.

Another study on mosquito control was reported by Lame *et al.* (2015) stating that the chloroform fraction could kill 36% larvae at 2500 ppm concentrations. The concentration used in the study was quite high, but the mortality of larvae was only 36%. When compared with the results of this study, nanoemulsion of essential oil of kaffir lime peel has a higher mortality.

Mortality in larvae and pupa of mosquito caused by nanoemulsion of essential oil of *Citrus hystrix* is caused by the active compound content in essential oil of *Cytrus hystrix*. According to Akono *et al.* (2015), larval mortality is caused by the presence of β -pinene and limonene compounds in *Citrus limon*, *Citrus sinensis* and *Citrus reticulata*. In this experiment, β -pinene and limonene compounds are the main compounds with the highest content in the essential oil of kaffir lime peel. Deaths of larvae and pupa of *Aedes aegypti* are most likely due to β -pinene and limonene compounds in the essential oil of *Citrus hystrix*.

The use of essential oil as an insecticide material is a good alternative to reduce the impact of synthetic insecticide used in the environment. The results showed that nanoemulsion of essential oil of kaffir lime peel has the ability to kill larvae and pupa of mosquito *Aedes aegypti* L. The larvicidal activity of nanoemulsion of essential oil of *Citrus hystrix* and abate is higher than the activity of its pupicidal activity. The mortality of larvae and pupa

because of the presence of abate is higher than that of the presence of nanoemulsion of essential oil of *Citrus hystrix*. However, nanoemulsion of essential oil of *Citrus hystrix* is more environmentally friendly than that of abate.

Conclusion

The essential oil of *Citrus hystrix* could be promoted as an efficient repellent larvicidal and pupicidal agent. The major chemical compounds were *beta-pinene* and *limonene*, both as active components that can kill the larvae and pupae. The particle size from nanoemulsion was \pm 129.45 nm and \pm 109.15 nm. The larvicidal activity was 21.6 \pm 1.67% (LC₅₀ = 546.592 ppm) and pupicidal activity 36 \pm 1.67% (LC₅₀ = 3045.323 ppm). Nanoemulsion essential oil of *Citrus hystrix* is more environmentally friendly than abate.

Acknowledgement

The authors acknowledge the financial of Directory Research and Service Society and Ministry Education and Cultural of Indonesia No Project 547/B3.1/KM/2017.

References

- Akono, P.N., Dongmo, P.M.J., Tonga, C., Kouotou, S., Kekeunou, S., Magne, G.T., Lehman, L.G. and Menut, C. 2015. Larvicidal activity of essential oils from pericarps of ripe *Citrus* fruits cultivated in Cameroon on pyrethroids sensitive and resistant strains of *Anopheles gambiae* Giles, 1902. *Journal of Entomology and Zoology Studies*. 3: 334-339
- Anjali, C.H., Sharma, Y., Mukherjee, A. and Chandrasekaran, N. 2011. Neem oil (*Azadirachta indica*) nanoemulsion - a potent larvicidal agent against *Culex quinquefasciatus*. *Pest Manag Sci*. 68 : 158-163.
- Balasubramani, S., Rajendhiran, T., Moola, A.N. and

- Diana, P.K.B. 2017. Development of nanoemulsion from *Vitex negundo* L. essential oil and their efficacy of antioxidant, antimicrobial and larvicidal activities (*Aedes aegypti* L.). *Environ Sci Pollut Res.* 1-9.
- Duarte, J.L., Amado, J.R.R., Oliveira, A.E.M.F.M., Cruz, R.A.S., Ferreira, A.M., Souto, R.N.P., Falcao, D.Q., Carvalho, J.C.T. and Fernandes, C.P. 2015. Evaluation of larvicidal activity of a nanoemulsion of *Rosmarinus officinalis* essential oil. *Brazilian Journal of Pharmacognosy.* 25 : 189-192.
- El-khal, F., Lalami, A.E.O. and Guemmouh, R. 2015. Larvacidal activity of essential oils of *Citrus sinensis* and *Citrus aurantium* (Rutaceae) cultivated in Morocco against the malaria vector *Anopheles labranchiae* (Diptera: Culicidae). *Asian Pac J Trop Dis* 5: 458-462.
- Kementrian Kesehatan Republik Indonesia (KKRI). 2020. Profil Kesehatan Indonesia 2016. Jakarta: Kementerian Kesehatan RI
- Khandagle, A.J., Tare, V.S., Raut, K.D. and Morey, R.A. 2011. Bioactivity of essential oils of *Zingiber officinalis* and *Achyranthes aspera* against mosquitoes. *Parasitol Res.* 109: 339-343.
- Krishnappa, K., Dhanasekaran, S. and Elumalai, K. 2012. Larvicidal, ovicidal and pupicidal activities of *Gliricidia sepium* (Jacq.) (Leguminosae) against the malarial vector, *Anopheles stephensi* Liston (Culicidae: Diptera). *Asian Pacific Journal of Tropical Medicine.* 598-604.
- Kumar, S., Warikoo, R. and Wahab, N. 2010. Larvicidal potential of ethanolic extracts of dried fruits of three species of peppercorns against different instars of an indian strain of dengue fever mosquito, *Aedes aegypti* L. (Diptera: Culicidae) *Parasitol Res.* 107 : 901-907.
- Lame, Y., Nukene, E.N., Pierre, D.Y.S., Elijah, A.E. and Esimone, C.O. 2015. Laboratory evaluation of the fractions efficacy of *Annona senegalensis* (Annonaceae) leaf extract on immature stage development of malaria filarial mosquito vektors. *J Arthropod-Borne Dis.* 9 : 226-237.
- Loh, F.S., Awang, R.M., Omar, D. and Rahman, M. 2011. Insecticidal properties of *Citrus hystrix* DC leaves essential oil against *Spodopteralitura fabricius*. *Journal of Medicinal Plants Research.* 5 : 3739-3744.
- Noveriza, R., Mariana, M. and Yuliani, S. 2017. Keefektifan formula nanoemulsiminyakseraiwangiterhadap Potyvirus penyebabpenyakitmosaik pada tanamannilam. *Bul. Litro.* 28 : 47-56.
- Nugroho, A.D. 2011. Kematian larva *Aedes aegypti* setelah pemberian abate dibandingkan dengan pemberian serbuk serai. *Jurnal Kesehatan Masyarakat.* 7 : 91-96.
- Othman, S.N.A.M., Hassan, M.A., Nahar, L., Basar, N., Jamil, S., Sarker, S.D. 2016. Essential oils from the Malaysian *Citrus* (Rutaceae) medical plants. *Medicines.* 3: 1-11.
- Shah, P., Bhalodia, D. and Shelat, P. 2010. Nanoemulsion: a pharmaceutical review. *Systematic Reviews in Pharmacy.* 1 : 24-32.
- Subekti, N., Patien, M., Kawulur, E.I.J.J., Sirait, S.H.K. and Mohammed, S. 2018. Types of Plasmodium and the Effect of Environmental factor Against Malaria in Manokwari, West Papua. *Jurnal Pendidikan IPA Indonesia.* 7 (3): 322-332 URL: <https://journal.unnes.ac.id/nju/index.php/jpii/article/view/14236/8255>
- Sugumar, S., Clarke, S.K., Nirmala, M.J., Tyagi, B.K., Mukherjee, A. and Chandrasekaran, N. 2014. Nanoemulsion of eucalyptus oil and its larvicidal activity against *Culex quinquefasciatus*. *Bull Entomol Res.* 104 : 393-402.
- Warikoo, R., Wahab, N. and Kumar, S. 2011. Oviposition altering and ovicidal potentials of five essential oils against female adult of the dengue vektor. *Parasitol Res.* 109 : 1125-1131.
- World Health Organization (WHO). 2005. Guidelines for laboratory and field testing of mosquito larvicides. World health organization communicable disease control, prevention and eradication Who pesticide evaluation scheme. WHO, Geneva, Switzerland.
-

Larvicidal and pupicidal activities from *Citrus hystrix* against *Aedes aegypti* mosquito

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	link.springer.com Internet Source	3%
2	journals.plos.org Internet Source	1%
3	"Proceedings of the Second International Conference on the Future of ASEAN (ICoFA) 2017 – Volume 2", Springer Science and Business Media LLC, 2018 Publication	1%
4	Tran Thi Kim Ngan, Tran Thien Hien, Xuan Tien Le, Trieu Tuan Anh et al. "Physico-Chemical Profile of Essential oil of Kaffir Lime (<i>Citrus hystrix</i> DC) Grown in An Giang Province, Vietnam", Asian Journal of Chemistry, 2019 Publication	1%
5	Submitted to Universiti Teknologi MARA Student Paper	1%
6	www.readbag.com Internet Source	1%

7 Mathivanan Thangarasu, Krishnappa Kaliyamoorthy, Elumalai Kuppusamy. "Pesticidal and Mosquitocidal Activities of *Burm. F.* (Rutaceae) against (*Fab.*) (Noctuidae: Lepidoptera) and *L., Liston, and Say.* (Diptera: Culicidae) ", *Advances in Zoology*, 2014

Publication

1 %

8 Ahn, Jeong-Hyeon, Akash Deep, and Ki-Hyun Kim. "The storage stability of biogenic volatile organic compounds (BVOCs) in polyester aluminum bags", *Atmospheric Environment*, 2016.

Publication

1 %

9 www.biotech-asia.org

Internet Source

1 %

10 Linda Julianti Wijayadi, Taty Rusliati Rusli. "Encapsulated Lime Peel Essential Oil () Into Chitosan Nanoparticle: New Entity to Enhanced Effectivity Against *Propionilbacterium Acne* in Vitro ", *IOP Conference Series: Materials Science and Engineering*, 2020

Publication

<1 %

11 T T Hien, T T K Ngan, H X Phong, T Q Toan. "Application of green technology in the process of extracting essential oil from Vietnam's Kaffir lime () leaves ", *IOP*

<1 %

Conference Series: Materials Science and Engineering, 2020

Publication

12

CH Anjali, Yamini Sharma, Amitava Mukherjee, Natarajan Chandrasekaran. "Neem oil (Azadirachta indica) nanoemulsion-a potent larvicidal agent against Culex quinquefasciatus", Pest Management Science, 2012

Publication

<1 %

13

Cleidjane Gomes Faustino, Fernando Antônio de Medeiros, Allan Kardec Ribeiro Galardo, Alex Bruno Lobato Rodrigues et al. "Larvicide Activity on Aedes aegypti of Essential Oil Nanoemulsion from the Protium heptaphyllum Resin", Molecules, 2020

Publication

<1 %

14

Kaliyamoorthy Krishnappa, Shanmugam Dhanasekaran, Kuppusamy Elumalai. "Larvicidal, ovicidal and pupicidal activities of Gliricidia sepium (Jacq.) (Leguminosae) against the malarial vector, Anopheles stephensi Liston (Culicidae: Diptera)", Asian Pacific Journal of Tropical Medicine, 2012

Publication

<1 %

15

Sylvia Jiero, Ayodhia Pitaloka Pasaribu. "Haematological profile of children with

<1 %

malaria in Sorong, West Papua, Indonesia",
Malaria Journal, 2021

Publication

16

Vijayalakshmi Ghosh, Amitava Mukherjee,
Natarajan Chandrasekaran. "Ultrasonic
emulsification of food-grade nanoemulsion
formulation and evaluation of its bactericidal
activity", Ultrasonics Sonochemistry, 2013

Publication

<1 %

17

Saranya Sugumar, Vijayalakshmi Ghosh, M.
Joyce Nirmala, Amitava Mukherjee, Natarajan
Chandrasekaran. "Ultrasonic emulsification of
eucalyptus oil nanoemulsion: Antibacterial
activity against Staphylococcus aureus and
wound healing activity in Wistar rats",
Ultrasonics Sonochemistry, 2014

Publication

<1 %

18

Tássio Rômulo Silva Araújo Luz, Ludmilla
Santos Silva de Mesquita, Flavia Maria
Mendonça do Amaral, Denise Fernandes
Coutinho et al. "Essential oils and their
chemical constituents against Aedes aegypti
L. (Diptera: Culicidae) larvae", Acta Tropica,
2020

Publication

<1 %

19

Mohamed M. Soumanou, Euloge S. Adjou.
"Sweet Fennel (Ocimum gratissimum) Oils",
Elsevier BV, 2016

Publication

<1 %

20 T. K. Lim. "Edible Medicinal And Non-Medicinal Plants", Springer Science and Business Media LLC, 2012
Publication <1 %

21 jurnal.umsu.ac.id
Internet Source <1 %

22 serisc.org
Internet Source <1 %

23 www.journalcra.com
Internet Source <1 %

24 A Senthilkumar, V Venkatesalu. "Larvicidal potential of Acorus calamus L. essential oil against filarial vector mosquito Culex quinquefasciatus (Diptera: Culicidae)", Asian Pacific Journal of Tropical Disease, 2012
Publication <1 %

25 S. Mohan, R. Ezhumalai, S. H. Jain, G. Ravikumar. "Chemical constituents of the essential oil from Dysoxylum malabaricum Bedd. wood of Western Ghats, India", Journal of the Indian Academy of Wood Science, 2011
Publication <1 %

26 academic.oup.com
Internet Source <1 %

27 coek.info
Internet Source <1 %

28

mafiadoc.com

Internet Source

<1 %

29

www.refsynjournals.com

Internet Source

<1 %

30

Mahmoud Osanloo, Amir Amani, Hassan Sereshti, Mohammad Reza Abai, Fariba Esmaeili, Mohammad Mehdi Sedaghat. "Preparation and optimization nanoemulsion of Tarragon (*Artemisia dracunculus*) essential oil as effective herbal larvicide against *Anopheles stephensi*", *Industrial Crops and Products*, 2017

Publication

<1 %

31

journalarticle.ukm.my

Internet Source

<1 %

32

journals.sagepub.com

Internet Source

<1 %

33

www.freepatentsonline.com

Internet Source

<1 %

34

www.mdpi.com

Internet Source

<1 %

35

Ricardo Marcelo dos Anjos Ferreira, Naima Pontes D'haveloose, Rodrigo Alves Soares Cruz, Raquel Silva Araújo et al. "Nano-emulsification Enhances the Larvicidal Potential of the Essential Oil of Siparuna

<1 %

guianensis (Laurales: Siparunaceae) Against Aedes (Stegomyia) aegypti (Diptera: Culicidae)", Journal of Medical Entomology, 2019

Publication

36

Shivani Sharma, Neha Loach, Shweta Gupta, Lalit Mohan. "Phyto-nanoemulsion: An Emerging Nano-Insecticidal Formulation", Environmental Nanotechnology, Monitoring & Management, 2020

Publication

<1 %

37

Abdulrhman A. Almadiy, Gomah E. Nenaah, Dalia M. Shaver. "Facile synthesis of silver nanoparticles using harmala alkaloids and their insecticidal and growth inhibitory activities against the khapra beetle", Journal of Pest Science, 2017

Publication

<1 %

38

Hamidah Jaafar Sidek, Fatin Fathihah Abdullah. "Chapter 52 Potential Antibacterial Activity of Essential Oil of Citrus hystrix and Chromolaena odorata Leaves", Springer Science and Business Media LLC, 2018

Publication

<1 %

39

Siti Md Othman, Muhammad Hassan, Lutfun Nahar, Norazah Basar, Shajarahtunnur Jamil, Satyajit Sarker. "Essential Oils from the

<1 %

Malaysian Citrus (Rutaceae) Medicinal Plants", Medicines, 2016

Publication

40

Y W Wulandari, C Anwar, S Supriyadi. " Effects of drying time on essential oil production of kaffir lime (DC) leaves at ambient temperature ", IOP Conference Series: Materials Science and Engineering, 2019

Publication

<1 %

Exclude quotes On

Exclude matches < 4 words

Exclude bibliography On