

# Lipid Profiles and Postprandial Glucose Status of Hypertensive Individuals with Stroke in Indonesia

*by* Oktia Woro Kasmini Handayani

---

**Submission date:** 04-Jan-2021 10:36AM (UTC+0700)

**Submission ID:** 1482798650

**File name:** Asian\_Journal\_of\_Epidemiology.pdf (539.49K)

**Word count:** 4518

**Character count:** 23932



# Asian Journal of Epidemiology

ISSN 1992-1462

**science**  
alert  
<http://www.scialert.net>

**ANSI***net*  
an open access publisher  
<http://ansinet.com>



## Research Article

# Lipid Profiles and Postprandial Glucose Status of Hypertensive Individuals with Stroke in Indonesia

<sup>1,2</sup>Mahalul Azam, <sup>1</sup>Fatkhiatun Nida, <sup>1</sup>Oktia Woro Kasmini Handayani and <sup>1</sup>Arulita Ika Fibriana

<sup>1</sup>Department of Public Health, Sports Science Faculty, Universitas Negeri Semarang, Semarang, Indonesia

<sup>2</sup>Department of Doctoral Program in Medicine and Health Sciences, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

## Abstract

**Background and Objective:** Stroke is the second leading cause of death worldwide. Stroke caused 23.48% of all deaths in Indonesia, a country with the highest prevalence of stroke in the world. The main risk factor for stroke is hypertension. In Indonesia, hypertension is the most prevalent non-communicable disease (25.8%). The objective of this study was to determine lipid profile and postprandial (PPG) status among hypertensive people with stroke in Indonesia. **Materials and Methods:** In this study cross-sectional study design was used and obtained data from secondary data of Riskesdas 2013. The number of respondents was 8,511. Data was analyzed using Cochran's and Mantel-Haenzel test. **Results:** Prevalence of stroke cases in people with hypertension was 16%. Most of stroke respondents have high T-cholesterol, LDL-C, triglyceride and PPG. Factors related to stroke are: age, LDL-C level (PR = 1.21; 95% CI = 1.026-1.431) and PPG levels ((PR = 1.28; 95% CI = 1.102-1.1492). After adjusting for gender and age, PPG was associated with stroke prevalence in hypertensive people in Indonesia. **Conclusion:** Prevalence of high LDL-C and PPG levels in hypertensive people with stroke is high. Those related to stroke incident among hypertensive people. It needs an educational effort to the stroke group to control LDL-C levels for prevention of recurrent stroke and other impacts.

**Key words:** Stroke, T-Cholesterol, LDL-C, HDL-C, triglyceride, postprandial glucose

**Citation:** Mahalul Azam, Fatkhiatun Nida, Oktia Woro Kasmini Handayani and Arulita Ika Fibriana, 2017. Lipid profiles and postprandial glucose status of hypertensive individuals with stroke in Indonesia. Asian J. Epidemiol., 10: 116-122.

**Corresponding Author:** Mahalul Azam, Department of Public Health, Sports Science Faculty, Universitas Negeri Semarang, Semarang, Indonesia  
Tel: +62 812-2853-982

**Copyright:** © 2017 Mahalul Azam *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

**Competing Interest:** The authors have declared that no competing interest exists.

**Data Availability:** All relevant data are within the paper and its supporting information files.

## INTRODUCTION

Stroke is a medical condition of concern both in Indonesia and globally, it is the second leading cause of death worldwide, causing 11.3% of all deaths<sup>1</sup>. In 2011, it was reported that 30.7 million individuals globally had previously experienced a stroke, 12.6 million individuals had moderate to severe disability as a result of stroke<sup>2</sup>. The burden of stroke globally is estimated to increase from 38-61 million disability-adjusted life years (DALYs) from 1990-2020. Additionally, stroke has been the single largest cause of disability in the United Kingdom (UK)<sup>3</sup>. In the United States (US), stroke and coronary heart disease (CHD) related health service costs were the most expensive, reaching more than 320.1 billion US dollars (USD) in 2013<sup>1</sup>.

Riset Kesehatan Dasar (Riskesdas), the Indonesian Basic Health Survey conducted by Ministry of Health Indonesia, reported that the incidence of stroke was 8.3/1000 population in 2007. This incidence increased to 12.1/1000 population in 2013<sup>4,5</sup>. Additionally, the incidence of mortality due to stroke also reached 0.16/100,000 population in 2013 in Indonesia. Stroke was the leading cause of death in Indonesia based on World Health Organization (WHO) data in 2015, with the number of deaths reaching 328,525 or 23.48% of all deaths in Indonesia. Moreover, Indonesia has the highest burden of strokes in the world<sup>6</sup>, thus increasing the risk of a large number of individuals living with poor health and quality of life<sup>7</sup>. According to Jaminan Kesehatan Nasional (JKN, National Health Insurance) data, stroke is the most costly medical condition to treat in Indonesian hospitals with total costs of treatment and care reaching 115.35 million USD<sup>8</sup>.

Hypertension is a major risk factor of stroke. Hypertension increased the risk of stroke 3.89 (99% confidence interval [CI] = 3.33-4.54) times when compared to the risk in non-hypertensive individuals<sup>9</sup>. Hypertension is also the most prevalent non-communicable disease in Indonesia with a prevalence of 25.8%<sup>5</sup>.

The relationship between lipid profiles and the postprandial glucose (PPG) status with stroke are well known. The risk of stroke increases 3.75 (95% CI 1.83-7.7) fold for higher total cholesterol<sup>10</sup>, 2.29 (95% CI 1.1-4.8) fold for higher low-density lipoprotein cholesterol (LDL-C)<sup>10</sup>, 0.52 (95% CI 0.28-0.96) fold for lower high-density lipoprotein cholesterol (HDL-C)<sup>10</sup>, 0.97 (95% CI 0.956-0.978) fold for higher triglyceride<sup>11</sup> and 2.71 (95% CI 1.2-91) fold for higher PPG<sup>12</sup>. Lipid profile<sup>13</sup> and PPG<sup>14</sup> status also play a pivotal role in prevention of recurrent stroke. However, there is lack of

information about the lipid profiles and PPG status in hypertensive individuals, who experienced stroke. Riskesdas 2013<sup>15</sup> was the latest survey conducted by Ministry of Health, Indonesia. This survey collected comprehensive health data, nevertheless lipid profiles and PPG levels in hypertensive individuals with stroke in Indonesia have not yet been reported on. The aim of this study was to determine the prevalence of stroke in hypertensive individuals in Indonesia and to report on their lipid profiles and PPG status based on the data from the Riskesdas, 2013.

## MATERIALS AND METHODS

**Design and sample research:** This cross-sectional study used secondary survey data from Riskesdas 2013<sup>15</sup>. This survey was the latest Riskesdas done and provide current condition describe health status and its determinants. This survey also provide comprehensive secondary health data and describe current condition of lipid profiles and PPG status that associated with hypertensive people with stroke in Indonesia. Subjects included were hypertensive respondents of Riskesdas 2013, subjects with incomplete data were excluded. Detailed information about the methods employed for the completion of Riskesdas 2013<sup>15</sup>, in addition to the data collected using this survey has been described in previous reports<sup>5,15</sup>.

**Data collection:** Riskesdas 2013<sup>15</sup> was a community-based survey covering all regions of Indonesia. Riskesdas collected baseline data and health indicators that represented a national, provincial and district/city description. Riskesdas is intended to be implemented every 3 years and the latest survey was conducted in 2013. This was a cross-sectional study with respondents from 33 provinces and 497 districts/cities in Indonesia, with a total of 1,027,763 respondents. This study was approved by the Ethics Committee, the National Institute of Health Research and Development (NIHRD), Ministry of Health, Indonesia in 2013. Respondents with hypertension were defined by having: Blood pressure  $\geq 140$  mm Hg (systolic pressure) and/or  $\geq 90$  mm Hg (diastolic pressure),<sup>16</sup> who were diagnosed with hypertension by a health professional, or prescribed antihypertensive medication. Stroke was determined by asking respondents whether or not they had been diagnosed as having a stroke by a health professional (doctor or nurse) in the past or were suddenly showing symptoms of a stroke. Symptoms, as assessed in Riskesdas 2013<sup>15</sup> included: experiencing paralysis on one side of the body, tingling or numbness on one side of the body, mouth droops without eye

muscle paralysis, slurred speech and difficulty in speaking and/or understanding conversation<sup>5</sup>. Factors that were analyzed were limited to age and biomedical parameters such as total cholesterol, LDL-C, HDL-C, triglyceride and PPG levels. Total cholesterol, LDL-C, HDL-C and triglyceride levels were measured using an auto analyzer TRX 7010<sup>®</sup> and PPG was measured using an Accu-Chek<sup>®</sup> Performa<sup>®</sup>.

Age was categorized as young (<45 years), middle aged (45-65 years) and elderly (>65 years). Total cholesterol, LDL-C, HDL-C and triglyceride levels were determined based on the dyslipidemia guidelines according to the Per kumpulan Endokrinologi Indonesia (PERKENI, The Indonesia Society of Endocrinology)<sup>17</sup>. PPG levels were determined according to the Ministry of Health, Indonesia monitoring guidelines on the factors for non-communicable diseases<sup>18</sup>. High risk total cholesterol, LDL-C, HDL-C and triglyceride levels were defined as a concentration  $\geq 190$  mg dL<sup>-1</sup>, a concentration  $\geq 110$  mg dL<sup>-1</sup>, a concentration  $< 40$  mg dL<sup>-1</sup> for men and  $< 45$  mg dL<sup>-1</sup> for women and a concentration  $\geq 150$  mg dL<sup>-1</sup>. Abnormal PPG levels were defined as a concentration  $\geq 180$  mg dL<sup>-1</sup>.

**Statistical analysis:** Characteristics of respondents were presented as frequencies and proportions. Here explored lipid profiles and PPG levels that were related to the prevalence of stroke adjusted for age and gender in hypertension patients using Cochran's and Mantel-Haenszel analysis. p-values  $< 0.05$  were considered statistically significant. All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS) software (version 13.0 for Windows, IBM SPSS Inc., Chicago, IL).

## RESULTS

In total, 8,511 respondents were included in the study of which 767 (9.0%) had stroke. The majority of the respondents were female (65.5%), aged 45-55 years (51.1%), had high risk total cholesterol (57.3%) and LDL-C (75.5%) and normal HDL-C (70%), triglyceride levels (71.8%) and PPG levels (77.9%). The clinical and demographic characteristics of respondents are presented in Table 1.

Out of seven variables assessed, the bivariate analysis found three related variables: Age, LDL-C level and PPG levels (Table 2). The prevalence of stroke was more common in women (9.2%) than in men (8.6%), however, this difference was not statistically significant ( $p = 0.347$ ).

Overall, 10.5, 10 and 6.7% of the prevalence of stroke occurred among those aged >65, 45-65 and <45 years, respectively. Current study found that there was a significant

Table 1: Characteristics of the subjects included in the analysis

Characteristics	Frequency (n)	Proportion (%)
<b>Gender</b>		
Male	2,933	34.5
Female	5,578	65.5
<b>Age group</b>		
<45 years	1,386	16.3
45-65 years	4,345	51.1
>65 years	2,780	32.7
<b>T-Cholesterol levels</b>		
$\geq 190$ mg dL	4,873	57.3
$< 190$ mg dL <sup>-1</sup>	3,638	42.7
<b>LDL-C levels</b>		
$\geq 100$ mg dL <sup>-1</sup>	6,426	75.5
$< 100$ mg dL <sup>-1</sup>	2,085	24.5
<b>HDL-C levels</b>		
At risk (M $< 40$ mg dL <sup>-1</sup> ; F $< 45$ mg dL <sup>-1</sup> )	2,556	30.0
No risk (M $\geq 40$ mg dL <sup>-1</sup> ; F $\geq 45$ mg dL <sup>-1</sup> )	5,955	70.0
<b>Triglyceride levels</b>		
$\geq 150$ mg dL <sup>-1</sup>	2,397	28.2
$< 150$ mg dL <sup>-1</sup>	6,114	71.8
<b>PPG levels</b>		
$\geq 180$ mg dL	1,885	22.1
$< 180$ mg dL <sup>-1</sup>	6,626	77.9
<b>Stroke</b>		
Yes	767	9
No	7744	91

LDL-C: Low density lipoprotein cholesterol, HDL-C: High density lipoprotein cholesterol

relationship between age and stroke prevalence in hypertensive individuals in Indonesia in all age groups when compared to reference group (<45 years). Respondents aged >65 and 45-65 years were at 1.57 and 1.48 times the risk of stroke when compared to respondents aged <45 years, respectively.

The prevalence of stroke was found to be higher in the high risk total cholesterol group (9.5%) compared to the no risk group (8.4%). However, there was no significant association between total cholesterol levels and the prevalence of stroke in hypertensive individuals in Indonesia ( $p = 0.087$ ,  $\alpha = 0.05$ ).

A significantly higher prevalence of stroke prevalence was found in the high risk LDL-C group (9.4%) compared to the no risk group (7.2%). ( $p = 0.025$ ,  $\alpha = 0.05$ ). Respondents in the high risk LDL-C group were 1.21 times more likely to have a stroke than respondents, who were in the no risk LDL-C group. The prevalence of stroke was higher in the no risk HDL-C group (9.2%) than in the high risk HDL-C group (8.6%), however this difference was not significant ( $p = 0.465$ ,  $\alpha = 0.05$ ). The prevalence of stroke was higher in the high risk triglyceride group (9.8%) than the no risk triglyceride group (8.7%), however, this difference was not significant ( $p = 0.120$ ,  $\alpha = 0.05$ ). Finally, the prevalence of stroke was significantly higher in the abnormal PPG group (10.9%) when compared to

Table 2: Relationship between parameters and stroke among respondents with hypertension

Parameter	Stroke (n = 747)		Stroke (n = 7744)		p-value	PR	95% CI
	No.	%	No.	%			
<b>Gender</b>							
Male	252	8.6	2,681	91.4	0.347	0.93	0.806-1.075
Female	515	9.2	5,063	90.8			
<b>Age group</b>							
>65 years	146	10.5	1,240	89.5	0.0001	1.57	1.274-1.925
45-65 years	434	10.0	3,911	10.0	0.0001	1.48	1.259-1.751
<45 years	187	6.7	2,593	93.3			
<b>T-Cholesterol levels</b>							
≥190 (mg dL <sup>-1</sup> )	462	9.5	4,411	90.5	0.087	1.13	0.985-1.298
<190 (mg dL <sup>-1</sup> )	305	8.4	3,333	91.6			
<b>LDL-C levels</b>							
≥100 (mg dL <sup>-1</sup> )	605	9.4	5,821	90.6	0.025	1.21	1.026-1.431
<100 (mg dL <sup>-1</sup> )	162	7.8	1,923	92.2			
<b>HDL-C levels</b>							
At Risk (M <40 mg dL <sup>-1</sup> ; F <45 mg dL <sup>-1</sup> )	221	8.6	2,335	91.4	0.465	0.94	0.812-1.095
No Risk (M ≥40 mg dL <sup>-1</sup> ; F ≥45 mg dL <sup>-1</sup> )	546	9.2	5,409	90.8			
<b>Triglyceride levels</b>							
≥150 (mg dL <sup>-1</sup> )	235	9.8	2,162	90.2	0.12	1.13	0.974-1.304
<150 (mg dL <sup>-1</sup> )	532	8.7	5,582	91.3			
<b>PPG levels</b>							
≥180 (mg dL <sup>-1</sup> )	205	10.9	1,680	89.1	0.002	1.28	1.102-1.1492
<180 (mg dL <sup>-1</sup> )	562	8.5	6,064	91.5			

Table 3: Comparison of prevalence ratio

Parameters	Model 1* PR (95% CI)	Model 2** PR (95% CI)	Model 3*** PR (95% CI)	Model 4**** PR (95% CI)
T-Cholesterol	1.13 (0.985-1.298)	1.14 (0.976-1.325)	1.10 (0.941-1.277)	1.09 (0.937-1.277)
LDL-C	1.30* (1.054-1.597)	1.23* (1.024-1.473)	1.20 (0.996-1.433)	1.20 (0.997-1.439)
HDL-C	0.94 (0.812-1.095)	0.94 (0.798-1.108)	0.98 (0.830-1.153)	0.99 (0.840-1.168)
Triglyceride	1.13 (0.974-1.304)	1.15 (0.976-1.349)	1.12 (0.954-1.319)	1.16 (0.981-1.359)
PPG	1.28* (1.02-1.1492)	1.31* (1.107-1.552)	1.25* (1.057-1.483)	1.24* (1.043-1.464)

\*Significant p value at <0.05, \* Model 1: Crude prevalence ratio, \*\*Model 2: Prevalence ratio after adjusted by gender, \*\*\* Model 3: Prevalence ratio after adjusted by age, \*\*\*\* Model 4: Prevalence ratio after adjusted by gender and age

the normal group (8.5%) (p = 0.002, α = 0.05). Respondents with abnormal PPG levels had 1.28 times the risk of stroke than respondents, who had normal PPG levels.

The comparisons of the prevalence ratios in each model is shown in Table 3. Before adjusting for other variables, LDL-C and PPG were significantly related to stroke prevalence among individuals with hypertension in Indonesia (Model 1). Similarly, after adjustment for gender, LDL-C and PPG were significantly related to stroke incident among hypertensive individuals in Indonesia (Model 2). However, the prevalence ratio (PR) for LDL-C increased from 1.21-1.23, while from PPG it increased from 1.28-1.31. After adjustment for age in Model 3, only PPG was significantly related to stroke prevalence among individuals with hypertension in Indonesia leading to a reduction in the PR from 1.28 (Model 1) -1.25 (Model 3).

Similarly, after adjustment for age and gender, only PPG was significantly related to the prevalence of stroke among individuals with hypertension in Indonesia leading to a reduction in the PR from 1.28 (Model 1)-1.24 (Model 4).

## DISCUSSION

In this study found that 9% of individuals in Indonesia with hypertension included in this study experienced stroke. This prevalence rate is lower than that previously reported from a study conducted in Nigeria<sup>19</sup>. Here determined that age, LDL-C levels and PPG status were significantly associated with the occurrence of stroke among hypertensive individuals in Indonesia. However, there was no association between the prevalence of stroke and gender, total cholesterol, HDL-C and

triglyceride levels. Previous report<sup>19</sup> on lipid profiles and PPG status among hypertensive individuals with stroke are limited.

<sup>7</sup> Nigerian study<sup>19</sup> showed that modifiable factors related to stroke occurrence included higher maximum and minimum blood pressure in preceding 3 months, higher fasting plasma glucose levels, large amounts of alcohol consumption, less physical activity and increased carotid intima medial thickness (CIMT).

<sup>8</sup> In this study found a significant relationship between age and the prevalence of stroke in hypertensive individuals in Indonesia, the risk of stroke increases with increasing age. The majority of the cases of stroke were identified among those aged >65 years (10.5%), while among those <45 years it was much lower (6.7%). Generally, stroke occurs due to aging. Naturally, the blood vessels will become more rigid as a result of plaque buildup<sup>20</sup>. In young adults, increased risk of infarction and vascular risk factors in developing countries are associated with behaviors such as increased smoking and urbanization<sup>21</sup>. Although the relationship between age and stroke incident has been well documented, this study adds to the existing body of evidence on the association between age and stroke, particularly among hypertensive individuals in Indonesia.

<sup>25</sup> In this study found a significant relationship between LDL-C levels and the prevalence of stroke. Respondents, who had LDL-C levels >100 mg dL<sup>-1</sup> had 1.30 times the risk of experiencing stroke than respondents with LDL-C levels 100 mg dL<sup>-1</sup>. However, after adjustment for age and gender, LDL-C levels were not significantly related to the occurrence of stroke.

<sup>24</sup> The association between LDL-C levels and coronary heart disease has previously been proven. However, the relationship between LDL-C levels and the prevalence of stroke is controversial. The findings of this study were in line with those of a study performed in Kerala, India<sup>10</sup> which reported that respondents with a cholesterol level >200 mg dL<sup>-1</sup> were 2.29 times more likely to have a stroke than those with normal cholesterol levels (p = 0.009, 99% CI:1.27-4.15). Likewise, a study conducted in the Neurology Department of Sina Hospital, Iran<sup>11</sup> also revealed similar findings, LDL-C levels were a predictor of hemorrhagic stroke. Increased cholesterol and LDL-C levels were associated with a higher risk of ischemic stroke. Investigators concluded that LDL-C was a risk factor in both ischemic and hemorrhagic stroke<sup>11</sup>. The LDL-C levels were also associated with mortality after long-term stroke but was not a prognostic factor<sup>22</sup>. The relationship between LDL-C levels and stroke was different in an Indonesian cohort study<sup>12</sup> in the common population and in a Nigerian-African study<sup>19</sup> in a hypertensive population. These differences may be due to

differences in the number of respondents, control of other confounding factors, or limitations in the data.

<sup>8</sup> In this study found a significant relationship between PPG levels and the prevalence of stroke in hypertensive individuals in Indonesia. Respondents with abnormal PPG levels had 1.28 times the risk of experiencing stroke than respondents with normal PPG levels. However, after adjusted for age and gender, PPG was a significant variable related to stroke incident, decreasing the PR from 1.28 (Model 1)-1.24 (Model 4).

Research on the relationship between PPG levels and the occurrence of stroke is limited both in the general and hypertension populations. Most previous studies<sup>12,19</sup> have included fasting blood Glucose (FBG) levels as a research variable. A cohort study in Indonesia<sup>12</sup> showed a significant association between PPG levels and stroke incidence in the general population, however, after controlling for other potentially confounding variables, PPG was not the dominant variable causing stroke<sup>12</sup>. However the study<sup>6</sup> in Nigeria (2013)<sup>19</sup> used FBG levels in order to determine the relationship between FBG and stroke incidence in patients with hypertension. The investigators found a significant relationship between FBG and the incidence of stroke in patients with hypertension. However, when other potentially confounding variables were controlled for, FBG levels were not the dominant variable causing stroke in hypertensive patients.<sup>19</sup> The San Luigi diabetes study showed that PPG was a strong cardiovascular predictor of FBG in individuals with type 2 diabetes mellitus, particularly in women.<sup>23</sup> High PPG levels or postprandial hyperglycemia are characterized by hyperglycemic spikes that cause endothelial dysfunction, inflammatory reactions and oxidative stress, which can lead to the development of atherosclerosis and the occurrence of cardiovascular events. Research has shown that postprandial hyperglycemia may predispose individuals to the development of atherosclerosis and cardiovascular events<sup>24</sup>.

This study showed a significant association between PPG levels and stroke incidence in hypertensive patients in Indonesia. Findings of this study might be different from those observed in the Nigerian study<sup>19</sup> because this study included more respondents and collected more robust data. Patients with hypertension, who had abnormal PPG levels had 1.28 times the risk of stroke when compared to those with normal PPG levels. After controlling for age, this risk decreased from 1.28-1.25 times the risk. Nonetheless, this study provides new insights as to stroke among hypertensive patients, that there is a significant relationship between PPG levels and the incidence of stroke in hypertensive individuals, especially in Indonesia.

This study was subject to several limitations. As this was a cross-sectional study, exposures and effects were measured in the same time period, thus, it is unknown exactly which events occurred first (i.e., whether respondents had hypertension, stroke, or high-risk LDL-C levels first). Additionally, we were unable to explain the LDL-C status before the data collection. However, from Table 2 it can be observed that 9.4% of stroke patients still had high risk LDL-C levels ( $\geq 100$  mg dL<sup>-1</sup>). These findings suggest that LDL-C levels were poorly controlled in the stroke group. Future studies are needed to explain the role of PPG levels on the increased risk of stroke.

## CONCLUSION

The majority of hypertensive individuals with stroke have high LDL-C and PPG levels. There was a significant relationship between age and LDL-C and PPG levels and the prevalence of stroke in hypertensive individuals in Indonesia. After adjustment for gender and age, PPG levels were significantly associated with prevalence of stroke among hypertensive patients. Education should be provided to those, who have experienced a stroke in the past in order to provide them with mechanisms to control LDL-C and PPG levels for the prevention of recurrent stroke.

## SIGNIFICANCE STATEMENTS

This study investigated the LDL-C and PPG levels related to hypertensive individuals with stroke. This study help the researchers to uncover the critical area risk factors of stroke, especially lipid profiles and PPG levels in individuals with hypertension that many researchers have not been able to explore in past. Current study found that LDL-C and PPG were the main risk factors of stroke in hypertensive individuals, thus interventions should target these high risk populations.

## ACKNOWLEDGMENTS

We thank the Research Institute of Universitas Negeri Semarang for supporting the research and as the National Institute of Health Research and Development (NIHRD) and the Ministry of Health in Indonesia for permission to access Riskesdas 2013 data.

## REFERENCES

- Go, A.S., D. Mozaffarian, V.L. Roger, E.J. Benjamin and J.D. Berry *et al*, 2014. Heart disease and stroke statistics-2014 update: A report from the American Heart Association. *Circulation*, 129: e28-e292.
- WHO., 2011. Global Atlas on Cardiovascular Disease Prevention and Control: Policies, Strategies and Interventions. WHO., Geneva, Switzerland, ISBN: 9789241564373, Pages: 164.
- WHO., 2004. Global Burden of Stroke. In: The Atlas of Heart Disease and Stroke, WHO. (Eds.), World Health Organization, Geneva, Switzerland, ISBN-13: 9789241562768, pp: 50-51.
- Badan Penelitian dan Pengembangan Kesehatan, 2007. Riset kesehatan dasar riskesdas 2007. Departemen Kesehatan, Jakarta, Republik Indonesia.
- Badan Penelitian dan Pengembangan Kesehatan, 2013. Riset kesehatan dasar riskesdas 2013. December 1, 2013, Kementerian Kesehatan, Jakarta, Republik Indonesia. <http://www.depkes.go.id/resources/download/general/Hasil%20Riskesdas%202013.pdf>
- WHO., 2015. Indonesia: WHO statistical profile. Volume 34, World Health Organization, Geneva, Switzerland.
- Endarti, A.T. and A. Handito, 2016. The relationship between non-communicable diseases history and health-related quality of life. *J. Kesehatan Masyarakat*, 12: 120-130.
- Moeloek, N.F., 2016. Kebijakan pembangunan kesehatan. Rapat Kerja Kesehatan Nasional, Jakarta, April 5, 2016. [http://www.depkes.go.id/resources/download/informasi/rakerkesnas\\_gel2\\_2016/Paparan%20Menkes.pdf](http://www.depkes.go.id/resources/download/informasi/rakerkesnas_gel2_2016/Paparan%20Menkes.pdf)
- Tu, J.V., 2010. Reducing the global burden of stroke: INTERSTROKE. *Lancet*, 376: 74-85.
- Grace, M., K.J. Jacob, A.V. Kumar and V.K. Shameer, 2016. Role of dyslipidemia in stroke and comparison of lipid profile in ischemic and hemorrhagic stroke -a case control study. *Int. J. Adv. Med.*, 3: 694-698.
- Togha, M., M.R. Gheini, B. Ahmadi, P. Khashaiar and S. Razeghi, 2011. Lipid profile in cerebrovascular accidents. *Iran. J. Neurol.*, 10: 1-4.
- Riyadina, W. and E. Rahajeng, 2013. Determinan penyakit stroke. *Kesmas: Nat. Public Health J.*, 7: 324-330.
- Chen, W., Y. Pan, J. Jing, X. Zhao and L. Liu *et al*, 2017. Recurrent stroke in minor ischemic stroke or transient ischemic attack with metabolic syndrome and/or diabetes mellitus. *J. Am. Heart Assoc.* 10.1161/JAHA.116.005446.
- Alte, M., S.M. Lai, G. Friday, V. Singh, V.M. Kumar and E. Sobel, 1997. Stroke recurrence in diabetics. Does control of blood glucose reduce risk? *Stroke*, 28: 1153-1157.
- Badan Penelitian dan Pengembangan Kesehatan, 2013. Riskesdas biomedis riset kesehatan dasar 2013. Kementerian Kesehatan, Jakarta, Republik Indonesia.
- U.S. Department of Health and Human Health and Human Services, 2003. Prevention, detection, evaluation and treatment of high blood pressure: Report. National Institutes of Health, National Heart, Lung and Blood Institute, USA.
- Arsana, P.M., R. Rosandi, A. Manaf, A. Budhiarta and K.W. Sucipta *et al*, 2015. Panduan Pengelolaan Dislipidemia di Indonesia. PERKENI., Indonesia, pp: 51.



18. Kemenkes, R.I., 2015. Buku monitoring faktor risiko penyakit tidak menular. Kementerian Kesehatan, Jakarta, Republik Indonesia.
19. Owolabi, M.O. and A.M. Agunloye, 2013. Risk factors for stroke among patients with hypertension: A case-control study. *J. Neurol. Sci.*, 325: 51-56.
20. Noviyanti, R.D., 2014. Faktor risiko penyebab meningkatnya kejadian stroke pada usia remaja dan usia produktif. *Profesi*, 10: 52-56.
21. Brainin, M., Y. Teuschl and L. Kalra, 2007. Acute treatment and long-term management of stroke in developing countries. *Lancet Neurol.*, 6: 553-561.
22. Xing, Y., Z. An, N. Yu and W. Zhao, X. Ning and J. Wang, 2016. Low density lipoprotein cholesterol and the outcome of acute ischemic stroke: Results of a large hospital-based study. *Eur. Neurol.*, 76: 195-201.
23. Cavalot, F., A. Petrelli, M. Traversa, K. Bonomo and E. Fiora *et al.*, 2005. Postprandial blood glucose is a stronger predictor of cardiovascular events than fasting blood glucose in type 2 diabetes mellitus, particularly in women: Lessons from the San Luigi Gonzaga diabetes study. *J. Clin. Endocrinol. Metab.*, 91: 813-819.
24. Node, K. and T. Inoue, 2009. Postprandial hyperglycemia as an etiological factor in vascular failure. *Cardiovasc. Diabetol.*, Vol. 8. 10.1186/1475-2840-8-23.

# Lipid Profiles and Postprandial Glucose Status of Hypertensive Individuals with Stroke in Indonesia

## ORIGINALITY REPORT

18%	%	16%	7%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

## PRIMARY SOURCES

- 1** "Minutes of The 43rd General Assembly of The European Association for The Study of Diabetes", Diabetologia, 2008  
Publication 2%
- 2** Mahalul Azam, Rina Sulistiana, Arulita Ika Fibriana, Soesmeyka Savitri, Syed Mohamed Al Junid. "Prevalence of Mental Health Disorders among Elderly Diabetics and Associated Risk Factors in Indonesia", Cold Spring Harbor Laboratory, 2020  
Publication 2%
- 3** Submitted to Universitas Diponegoro  
Student Paper 2%
- 4** Submitted to Universitas Jember  
Student Paper 1%
- 5** Mahalul Azam, Fetty Nur Hidayati, Arulita Ika Fibriana, Udin Bahrudin, Syed Mohamed Aljunid. "Prevalence of Isolated Systolic Hypertension among People with Diabetes in Indonesia", Cold Spring Harbor Laboratory, 2020 1%

6

Noer Bahry Noor, Ridwan Amiruddin, Muhammad Awal, Sukri Palutturi, Anwar Mallongi. "Proxy Model of Comorbidities with Stroke Incident in South Sulawesi", Pakistan Journal of Nutrition, 2017

Publication

---

7

M.O. Owolabi, A.M. Agunloye. "Risk factors for stroke among patients with hypertension: A case–control study", Journal of the Neurological Sciences, 2013

Publication

---

8

Ruisheng Duan, Wenjun Xue, Kunpeng Wang, Nan Yin, Hongyu Hao, Hongshan Chu, Lijun Wang, Peng Meng, Le Diao. "Estimation of the LDL Subclasses in Ischemic Stroke as a Risk Factor in a Chinese population", Research Square, 2020

Publication

---

9

Koichi Node. "Postprandial hyperglycemia as an etiological factor in vascular failure", Cardiovascular Diabetology, 2009

Publication

---

10

Yuan Zhang, Manuel Cifuentes, Xiang Gao, Gina Amaral, Katherine L. Tucker. "Age- and gender-specific associations between insomnia and falls in Boston Puerto Rican adults", Quality of Life Research, 2016

Publication

---

1%

1%

1%

1%

1%

11

Submitted to Universitas Negeri Semarang

Student Paper

&lt;1 %

12

Submitted to Methodist College

Student Paper

&lt;1 %

13

Sohan Singh Hayreh. "Ischemic Optic Neuropathies", Springer Science and Business Media LLC, 2011

Publication

&lt;1 %

14

Wahyu Septiono, Mirte A.G. Kuipers, Nawi Ng, Anton E. Kunst. "Changes in adolescent smoking with implementation of local smoke-free policies in Indonesia: Quasi-experimental repeat cross-sectional analysis of national surveys of 2007 and 2013", Drug and Alcohol Dependence, 2020

Publication

&lt;1 %

15

Krishnamoorthy Srilatha, Zachariah Bobby, D.K. Subrahmanyam, Uppoor NithinKumar. "Insulin resistance and elevated C-reactive protein among first-degree relatives of ischemic stroke patients", Diabetes &amp; Metabolic Syndrome: Clinical Research &amp; Reviews, 2017

Publication

&lt;1 %

16

Miwako Nagasu. "Prevalence and risk factors for low back pain among professional cooks working in school lunch services", BMC Public Health, 2007

Publication

&lt;1 %

- 
- 17 R. Jago. "Prevalence of Abnormal Lipid and Blood Pressure Values Among an Ethnically Diverse Population of Eighth-Grade Adolescents and Screening Implications", PEDIATRICS, 2006  
Publication <1%
- 
- 18 Submitted to Higher Education Commission Pakistan  
Student Paper <1%
- 
- 19 Submitted to Universitas Sebelas Maret  
Student Paper <1%
- 
- 20 Bartoll, X., L. Palencia, D. Malmusi, M. Suhrcke, and C. Borrell. "The evolution of mental health in Spain during the economic crisis", The European Journal of Public Health, 2013.  
Publication <1%
- 
- 21 M. G Duerden. "Guidelines from the British Hypertension Society: BHS is set to bankrupt NHS", BMJ, 9/4/2004  
Publication <1%
- 
- 22 Takuya Oikawa, Yasuhiko Sakata, Kotaro Nochioka, Masanobu Miura et al. "Prognostic Impact of Statin Intensity in Heart Failure Patients With Ischemic Heart Disease: A Report From the CHART-2 (Chronic Heart Failure Registry and Analysis in the Tohoku District 2) Study", Journal of the American

23

P. Armario. "Antihypertensive treatment and stroke prevention: are angiotensin receptor blockers superior to other antihypertensive agents?", *Therapeutic Advances in Cardiovascular Disease*, 06/01/2009

Publication

---

<1%

24

Musunuru, K.. "A synergistic relationship of elevated low-density lipoprotein cholesterol levels and systolic blood pressure with coronary artery calcification", *Atherosclerosis*, 200810

Publication

---

<1%

25

S. Haffner. "Metabolic syndrome - a new risk factor of coronary heart disease?", *Diabetes Obesity and Metabolism*, 11/2003

Publication

---

<1%

26

Mohd Razif Shahril, Syed Amirfaiz, Pei Lin Lua, Ali Nurnazahiah et al. "Prevalence of Metabolic Syndrome among Breast Cancer Survivors in East Coast of Peninsular Malaysia", *Research Square*, 2020

Publication

---

<1%

27

S E Humphries. "Genetic causes of familial hypercholesterolaemia in patients in the UK: relation to plasma lipid levels and coronary heart disease risk", *Journal of Medical Genetics*, 7/6/2006

<1%

28

I. Holme. "Lipoprotein components and risk of congestive heart failure in 84 740 men and women in the Apolipoprotein MOrtality RISk study (AMORIS)", European Journal of Heart Failure, 10/03/2009

Publication

---

<1 %

29

Huihui Liu, Ruojun Wang, Jijun Shi, Yanlin Zhang, Zhichao Huang, Shoujiang You, Guodong Xiao, Dapeng Wang, Yongjun Cao. "Baseline Neutrophil Counts and Neutrophil Ratio May Predict a Poor Clinical Outcome in Minor Stroke Patients with intravenous Thrombolysis", Journal of Stroke and Cerebrovascular Diseases, 2019

Publication

---

<1 %

30

Kristian Ade Chandra, Dessie Wanda. "Traditional Method of Initial Diarrhea Treatment in Children", Comprehensive Child and Adolescent Nursing, 2017

Publication

---

<1 %

31

Resty Ryadinency, Suharyo Hadisaputro, Banundari Rachmawati. "Effect of zinc supplementation on triglyceride and malondialdehyde levels: study on diabetic Wistar rats induced with streptozotocin", Medical Journal of Indonesia, 2018

Publication

---

<1 %

32

Titus Tambaip, Marni Br Karo, Mochammad

Hatta, Ressay Dwiyanthi et al.

"Immunomodulatory Effect of Orally Red Fruit (Pandanus conoideus) Extract on the Expression of CC Chemokine Receptor 5 mRNA in HIV Patients with Antiretroviral Therapy", Research Journal of Immunology, 2018

Publication

<1%

33

Cecilia M Shikuma, Yang Yang, Marshall J Glesby, William A Meyer et al. "Metabolic Effects of Protease Inhibitor-Sparing Antiretroviral Regimens Given as Initial Treatment of HIV-1 Infection (AIDS Clinical Trials Group Study A5095)", JAIDS Journal of Acquired Immune Deficiency Syndromes, 2007

Publication

<1%

34

Seung Hee Kim, Ki Young Son. "Association between lipoprotein cholesterol and future cardiovascular disease and mortality in older adults: a Korean nationwide longitudinal study", Research Square, 2020

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On