

Baseline and post

by Mahalul Azam

Submission date: 17-Feb-2021 05:25AM (UTC+0700)

Submission ID: 1511045384

File name: 560-2314-1-PB-dikompresi.pdf (398.51K)

Word count: 5900

Character count: 30717

RESEARCH ARTICLE

3
Baseline and Post-exercise High-Sensitivity C-Reactive Protein Levels in Endurance Cyclists: The Indonesian North Coast and Tour de Borobudur 2017 Study

Mahalul Azam^{1,2,*}, Susanti Lestari¹, Sri Ratna Rahayu¹, Arulita Ika Fibrina¹, Budhi Setianto³,
Nyoman Suci Widyastiti⁴, Suhartono⁵, Hardhono Susanto⁶, Martha Irene Kartasurya⁷,
Udin Bahrudin⁸, Thijs Eijssvogels⁹

¹Department of Public Health, Faculty of Sport Sciences, Universitas Negeri Semarang, Jl. Sekaran, Gunungpati, Semarang, Indonesia

²Doctoral Program in Medicine and Health Sciences, Faculty of Medicine, Universitas Diponegoro, Jl. Imam Bardjo SH No. 5, Semarang, Indonesia

³Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Indonesia, Jl. Letjen S. Panman Kav 87, Slipi, Jakarta, Indonesia

⁴Department of Clinical Pathology, Faculty of Medicine, Universitas Diponegoro, Jl. Doktor Sutomo No. 16, Semarang, Indonesia

⁵Department of Environmental Health, Faculty of Public Health, Universitas Diponegoro, Jl. Prof. Soedarto SH, Tembalang, Semarang, Indonesia

⁶Department of Anatomy, Faculty of Medicine, Universitas Diponegoro, Jl. Prof. Soedarto SH, Tembalang, Semarang, Indonesia

⁷Department of Public Health Nutrition, Faculty of Public Health, Universitas Diponegoro, Jl. Prof. Soedarto SH, Tembalang, Semarang, Indonesia

⁸Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Diponegoro, Jl. Doktor Sutomo No. 16, Semarang, Indonesia

⁹Department of Physiology, Radboud University Medical Center, Geert Grooteplein Zuid 30, Netherlands

*Corresponding author. E-mail: mahalul.azam@mail.unnes.ac.id

Received date: Sep 24, 2018; Revised date: Oct 25, 2018; Accepted date: Nov 21, 2018

Abstract

BACKGROUND: Inflammation plays an important role in the atherosclerotic process. High-sensitivity C-reactive-protein (hs-CRP) is commonly used as inflammatory biomarker. It is well known that regular physical activity lowers hs-CRP levels, while prolonged exercise induces hs-CRP elevations. However, the relationship of training and exercise characteristics with hs-CRP levels remains not well elucidated. We evaluated baseline and post-exercise hs-CRP levels and its association with training and exercise characteristics.

METHODS: Eighty-eight male endurance cyclists were involved. Demographic data, health condition and training characteristics were collected. Baseline and post-exercise blood-samples were collected to determine hs-CRP concentrations. A hs-CRP cut-off point of 3 mg/L was used. Blood-cell count and biochemical parameters were measured at baseline. Heart rate (HR) was measured during exercise.

RESULTS: Cyclists performed 7.3 hours (interquartile-range (IQR) = 5.4-7.5) of endurance exercise at intensity

of 81.8 % (IQR = 74.9-85.8). Cyclists with baseline hs-CRP \geq 3 mg/L reported higher body mass, body mass index (BMI), waist-circumference and total-cholesterol. An increase in hs-CRP was following endurance exercise. Cyclists with any elevation of hs-CRP reported a higher BMI, HR during exercise and exercise intensity. Binary logistic regression analysis showed BMI (OR = 1.24, 95% CI = 1.04-1.48) and cycling distance (OR = 0.22, 95% CI = 0.06-0.76) were associated with post-exercise hs-CRP elevations.

CONCLUSION: Body mass, BMI, waist-circumference, total- and HDL-cholesterol are associated with baseline hs-CRP, whereas BMI and cycling distance were associated with hs-CRP elevations. These findings suggest that anthropometry parameters and lipid levels attributed to baseline hs-CRP, while anthropometry parameters and cycling intensity attributed to post-exercise hs-CRP elevations.

KEYWORDS: C-reactive-protein, exercise, endurance-cycling, inflammation, acute-phase-response

Indones Biomed J. 2019; 11(1): 91-9

Introduction

46
Inflammation plays a pivotal role in the mechanism of atherosclerotic and is a risk factor for the development of coronary artery diseases (CAD) risk factor.(1) High-sensitivity C reactive protein (hs-CRP) is a common clinical biomarker to assess inflammation and is established as an independent predictor for CAD.(2) A systematic review reported that regular physical activity lowers resting hs-CRP levels, however as an acute-phase response hs-CRP is elevated after exercise.(3,4) CRP and tumor necrosis factor-alpha (TNF- α) were significantly increased after cross-country skiing tour (5), while no changes in interleukin (IL)-6 were observed. Another study reported that CRP levels were influenced by exercise capacity as assessed by a treadmill exercise test.(6)

Alongside with exercise-induced changes in hs-CRP, several cardiac injury markers such as creatine kinase-muscle/brain (CK-MB), cardiac troponin, and N terminal pro B-type natriuretic peptide (NT-proBNP) are known to increase as well.(7) There is still controversy whether these biomarker increases represent a deleterious effect or not.(7,8) Previous studies concluded that exercise-induced inflammatory responses are related to muscle injury and muscle damage.(9-11) An evidence also reported that highly post-exercise hs-CRP and cardiac troponin elevations were related to sub-clinical CAD event.(12,13)

10
A previous study reported that baseline hs-CRP was associated with body mass index (BMI), daily life activity level and high-sensitive cardiac troponin T (hs-cTnT) level in subjects suspected of the CAD.(14) Hs-CRP elevated following an exercise test (ET), but was not related to the CAD, exercise time, metabolic equivalents (METs), ST-segment depression, angina during ET, increase in heart rate, and increase in systolic blood pressure. Exercise-induced hs-CRP elevations are well-published (4-7), however, the relationship of training and exercise characteristics with hs-CRP elevation remains unclear.

Cycling is a popular sports activities and has many health benefits. However, endurance cycling is characterized by prolonged exercise at a high intensity, which is likely to cause exercise-induced inflammation. The present study explored the relationship of training and exercise characteristics with baseline and post-exercise hs-CRP levels among endurance cyclists. We hypothesized that anthropometric parameters *i.e.*, body mass, BMI, and waist circumference, lipid levels *i.e.*, total- and HDL- cholesterol, and training characteristics are associated with baseline

hs-CRP levels, while antropometric parameters and exercise intensity are associated with post-exercise hs-CRP levels among endurance cyclists.

Methods

Study Population

Ninety-two endurance cyclists of two Indonesian long distance cycling tours, *i.e.*, 2017 Indonesia North Coast tour (NC) (<http://ina-northcoast-cyclingtour.com>) and 2017 Tour de Borobudur (TdB) (<https://event.sambabikers.com>), were recruited for this prospective observational study. A detailed description of participants recruitment is presented in Figure 1.

Ethical Approval

The study protocol conformed to the concepts of the Declaration of Helsinki and approved by the local Ethics Committee in the Faculty of Medicine, Diponegoro University - Dr. Kariadi General Hospital, Semarang, Indonesia, and informed consent was obtained from all participants. The study was also registered at Protocol and Registration and Results System under Clinicaltrial.gov identifier (NCT number) NCT03310450.

Demographic Data and Blood Sampling

One day before the cycling tour, all cyclists completed a questionnaire on their personal characteristics, training characteristics, and health conditions, including a medical history. After completing the questionnaire, a 10 mL venous blood sample was drawn from the antecubital vein. All measurements *i.e.*, completing a post-exercise questionnaire, body weighing, and blood sampling were repeated in the same manner. Post-exercise blood sampling was taken immediately post-exercise (less than 5 minutes).

Anthropometric Measurements

We used common anthropometric measurements including waist and hip circumference (cm), body mass (kg), BMI (kg/m²), and waist to hip ratio. Anthropometric measurements for waist and hip circumference, body mass, and height respectively measured using circumference measuring tape SECA 201®, weighing scale SECA 813®, and height scale SECA Bodymeter 206® (SECA Corp, Birmingham, UK).

27 Heart Rate (HR)

HR was monitored during exercise using a chest band HR monitor (Polar RC3 GPS®, Polar Corporation, Kempele,




Figure 1. Participants recruitment flow diagram.

Finland). We determined mean HR as the average HR during cycling and peak HR as the highest HR during cycling. Maximal predicted heart rate ($HR_{max} = 208 - 0.7 * \text{age}$), and exercise intensity ($\text{Exercise intensity} = 100 * \text{mean HR} / HR_{max}$) were calculated.(15).

Laboratory Tests

The blood samples (10 mL) taken before and immediately after exercise were analyzed in the laboratory.^{3,2} The whole blood sample for routine blood was analyzed on an automated blood cell counter (Sysmex Xn-350, Sysmex Corporation, Kobe, Japan). The other fractions of whole blood were collected in serum-gel vacutainer tubes and allowed to clot for 45 minutes. After centrifugation, the serum was aliquoted, frozen, and stored at -80°C for future analysis. All analyses were conducted using the same calibration and setting to minimize variation. The hs-CRP was measured using a high-sensitivity CRP assay (CR120C, Calbiotech, El Cajon, USA). The assay detection limit was 0.02 mg/L. A hs-CRP value of 3 mg/L was used as the cut-off point. (14,16) Post-exercise hs-CRP elevation was determined by a positive delta hs-CRP ($\Delta\text{hs-CRP} = \text{post-exercise hs-CRP}$

level – baseline hs-CRP level). Biochemical parameters (*i.e.*, hemoglobin, total cholesterol and HDL cholesterol) were analyzed using automated analyzer Cobas C-501 (Roche Diagnostics, Indianapolis, USA).

Statistical Analysis

All numeric data with a Gaussian distribution were reported as a mean±standard deviation (SD) and statistical significance was considered at a $p\text{-value} < 0.05$. Statistical analyses were performed using the Statistical Package for the Social Sciences (IBM SPSS Statistics for Windows, Version 23.0, IBM Corp., New York, USA). The normality of the data distribution was examined by the Kolmogorov–Smirnov test. When the data demonstrated a non-Gaussian distribution,^{3,1} natural logarithmic transformation was applied. If the data remained not normally distributed after transformation, non-parametric statistics were used and data were presented as median with interquartile range.

A paired Student's t-test was used to test the significance of the differences between baseline and post-exercise hs-CRP levels. Bivariate logistic regression analysis with enter method was used to determine parameters that

were related to a baseline hs-CRP levels ≥ 3 mg/L and a post-exercise hs-CRP elevation. Based on our hypothesis and potential confounding factors, we included age, BMI, medical history (*i.e.*, history of hypertension, dyslipidemia, diabetes mellitus, family history of CAD), systolic and diastolic blood pressure, blood cell count, total and HDL cholesterol, exercise duration, mean HR, peak HR and exercise intensity in our model as potential determinants to predict post-exercise hs-CRP elevation. Thereafter, all predictors with a p -value < 0.10 were retained in the final regression model.

Results

Subject Characteristics

Cyclists completed the NC240K tour (n=31), TdB140K (n=207) and TdB100K (n=973) in 2017. NC240K participants cycled for 240 km with a total elevation of 826 m, in a sunny weather with an average temperature of 33°C (28°-41°C). TdB140K participants cycled for 140 km, with a total elevation of 2754 m, in a drizzle and mostly rainy weather, with an average temperature of 26°C (21°-37°C). The TdB100K group cycled 100 km, with a total elevation of 1117 m, and had sunny weather with an average temperature of 30°C (28°-34°C).

We recruited 92 male cyclists for this study, but n=4 athletes dropped out since they did not complete the post-exercise examinations and were excluded from further analyses. The remaining 88 cyclists covered 240 km (n=28), 140 km (n=30) and 100 km (n=30). Table 1 provides an overview of subject characteristics. CAD family history, HDL cholesterol levels, neutrophil count and resting HR, showed significant differences between the different cycling distance groups ($p < 0.05$). NC240K cyclists had a higher HDL cholesterol and lower neutrophil count and resting HR compared to the TdB140K and TdB100K cyclists. All other subject characteristics were comparable across groups.

Table 2 provides an overview of the training and exercise characteristics of the subjects. Training characteristics between groups were comparable. Due to the difference of cycling distance, we found differences in exercise duration, recovery HR and exercise intensity across groups ($p < 0.05$). Peak HR and mean HR did not differ across groups ($p > 0.05$).

Baseline hs-CRP

Baseline hs-CRP in NC240K cyclists had the lower levels (2.6 \pm 2.24 mg/L) than other groups (TdB140K = 3.3 \pm 3.31 mg/L; TdB100K = 3.0 \pm 3.23 mg/L) (Table 1). Cyclists with baseline hs-CRP levels ≥ 3 mg/L (n=24, mean age 46.5 \pm 11.10 years) had a higher body mass, BMI, waist

Table 1. Cyclist's characteristics based on touring groups.

Variables	NC240K (n=28)	TdB140K (n=30)	TdB100K (n=30)	Total (n=88)	P
Age (years old; mean \pm SD)	43.3 \pm 9.33	43.5 \pm 10.18	49 \pm 13.70	45.3 \pm 11.47	0.090*
Height (cm; mean \pm SD)	167.2 \pm 5.02	167.1 \pm 4.35	167.3 \pm 4.5	167.2 \pm 4.58	0.969*
Body mass (kg; mean \pm SD)	67.9 \pm 7.51	68.5 \pm 9.10	66.8 \pm 11.21	67.7 \pm 9.36	0.779*
BMI (kg/m ² ; mean \pm SD)	24.2 \pm 2.26	24.5 \pm 3.11	23.7 \pm 3.60	24.2 \pm 3.03	0.653*
Medical history					
Hypertension (n)	4	3	6	13	0.549**
Dyslipidemia (n)	7	12	9	28	0.456**
Diabetes mellitus (n)	1	1	0	2	0.588**
Family history of CAD (n)	3	0	0	3	0.036**
Lipid levels					
Total cholesterol levels (mg/dL; mean \pm SD)	216.6 \pm 55.04	208.1 \pm 27.13	203.1 \pm 31.95	209.1 \pm 39.45	0.427*
HDL cholesterol levels (mg/dL; mean \pm SD)	68.98 \pm 19.09	52.1 \pm 13.9	53.6 \pm 12.45	58.0 \pm 16.97	<0.001*
Neutrophil (10 ⁶ /L; mean \pm SD)	50.8 \pm 9.03	51.8 \pm 7.94	55.7 \pm 7.12	52.8 \pm 8.23	0.050*
Lymphocyte (10 ⁶ /L; mean \pm SD)	37.3 \pm 8.93	36.7 \pm 7.7	33.8 \pm 6.8	35.9 \pm 7.9	0.182*
Monocyte (10 ⁶ /L; mean \pm SD)	7.8 \pm 1.65	7.8 \pm 1.40	7.1 \pm 1.61	7.6 \pm 1.57	0.190*
Resting HR (bpm; mean \pm SD)	57.6 \pm 8.7	63.2 \pm 9.3	66.7 \pm 8.05	62.6 \pm 9.3	0.001*
Baseline hs-CRP (mg/L; mean \pm SD)	2.6 \pm 2.24	3.3 \pm 3.11	3.0 \pm 3.23	2.9 \pm 2.89	0.645*

Table 2. Training and exercise characteristics based on touring groups.

Variables	NC240K (n=28)	TdB140K (n=30)	TdB100K (n=30)	Total (n=88)	<i>p</i>
Cycling distance in previous month (km; median (range))	897 (205-1,360)	495 (180-1350)	582 (170-1,350)	602 (170-1,360)	0.205***
Long-distance cycling participation					
Yes (n)	26 (92.9%)	21 (70%)	24 (80%)	71 (80.7%)	0.088**
Never (n)	2 (7.1%)	9 (30%)	6 (20%)	17 (19.3%)	
Exercise characteristics					
Exercise duration (hours; median (range))	7.5 (7.5-8.0)	7.5 (4.0-9.8)	5.0 (3.5-6.5)	7.3 (3.5-9.8)	<0.001***
Recovery HR (bpm; median (range))	80.5 (57.0-133.0)	91.0 (59.0-168.0)	94.0 (73.0-113.0)	90.0 (57.0-168.0)	<0.001***
Mean HR (bpm; median (range))	140.0 (104.0-165.0)	133.0 (94.0-160.0)	141.0 (96.0-180.0)	140.0 (94.0-180.0)	0.151***
Peak HR (bpm; median (range))	168.0 (128.0-196.0)	180.8 (122.0-229.0)	172.5 (116.0-233.0)	173.5 (116.0-233.0)	0.075***
Exercise intensity (%; median (range))	84.3 (75.14-90.80)	80.8 (53.73-92.16)	74.5 (45.58-90.98)	81.8 (45.58-92.16)	<0.001***
Post-exercise hs-CRP (mg/L; mean±SD)	3.5±3.33	4.5±2.86	3.3±3.75	3.8±3.34	0.335 [†]

circumference, total cholesterol level and ratio of total cholesterol to HDL cholesterol (Table 3) compared to cyclists with baseline hs-CRP levels < 3 mg/L (n=64; mean age 44.9±11.66 years). Alongside with baseline hs-CRP levels among cycling distance groups, Table 3 also showed that there was no difference in the proportion of the cycling distance groups categorized by a cut-off point of 3 mg/L.

Post-exercise hs-CRP

A significant increase in hs-CRP levels *i.e.*, 0.9 mg/L was found ($p=0.022$) from baseline (2.9±2.89 mg/L) to post-exercise (3.8±3.34 mg/L) in our study population (Figure 2). However, the exercise-induced increase in hs-CRP appeared to be different among subgroups. A significant increase in hs-CRP levels *i.e.*, 0.9 mg/L was found in the NC240K group (2.6±2.24 mg/L to 3.5±3.33 mg/L, $p=0.020$) and 1.1 mg/L in the TdB140K group (3.3±3.31 mg/L to 4.5±2.86 mg/L, $p=0.003$), but was not found in the TdB100K participants (3.0±3.23 mg/L to 3.3±3.75 mg/L, $p=0.960$).

Cyclists with post-exercise hs-CRP elevations (n=58, mean age 43.7±11.20 years) had a higher BMI, higher HR during exercise, and higher exercise intensity (Table 3) compared to cyclists without post-exercise hs-CRP elevations (n=30; mean age 48.5±11.47 years). NC240K and TdB140K participants had a higher prevalence of post-exercise hs-CRP elevations (*i.e.*, 67.8 % and 80 % respectively) compared to TdB 100K participants (50 %).

Binary logistic regression analysis showed that BMI (OR: 1.241, 95% CI: 1.043-1.476); $p=0.015$) and cycling distance (OR: 0.217, 95% CI: 0.062-0.758); $p=0.017$) were the only parameters with a significant association

with post-exercise hs-CRP elevation in our multivariable model ($r=0.216$, $p=0.007$). The other variables included in the analysis did not reach statistical significance and were therefore excluded from regression analysis as determinants of post-exercise hs-CRP elevation (Table 4).

Discussion

The present study revealed several interesting findings. First, we found that BMI, waist circumference, total- and HDL-cholesterol were significantly associated with baseline hs-CRP levels. Second, we found a significant increase in hs-CRP levels after prolonged exercise and its related to BMI and cycling distance. These findings indicate that anthropometry measurement and lipid profile were related to resting hs-CRP concentration, whereas, besides anthropometry, cycling intensity was related to elevated hs-CRP as an acute phase response.

The predictors for baseline hs-CRP levels that were identified in the present study (*i.e.*, BMI, waist circumference, total- and HDL-cholesterol) align with previous studies in this field of research for the subjects of general large population (17), female elderly (18) and diabetic patients (19). A recent study reported that IL-6 and hs-CRP concentrations were related to anthropometric parameters, such as BMI, waist to hip ratio, body adiposity index and neck circumference.(18) Another study reports that hs-CRP, white blood count and fibrinogen were associated with waist circumference, body mass, BMI, body adiposity index, waist/hip ratio and waist/height ratio.

Table 3. Parameters based on baseline hs-CRP and post-exercise hs-CRP elevation.

Variables	Baseline hs-CRP		P	Post-exercise hs-CRP elevation		P
	≥3 mg/L (n=24)	<3 mg/L (n=64)		Yes (n=58)	No (n=30)	
Age (years old; mean±SD)	46.5±11.10	44.9±11.66	0.571*	43.7±11.20	48.5±11.47	0.064*
Body mass (kg; mean±SD)	70.5±6.78	66.7±10.00	0.043*	66.6±8.98	69.9±9.83	0.115*
BMI (kg/m ² ; mean±SD)	25.3±2.75	23.7±3.06	0.030*	25.1±3.56	23.7±2.62	0.029*
Waist circumference (cm; mean±SD)	88.3±6.06	84.1±8.93	0.015*	84.4±8.20	86.8±8.73	0.210*
Hip circumference (cm; mean±SD)	90.8±4.77	88.6±7.45	0.104*	88.5±6.80	90.4±6.95	0.233*
Waist-hip ratio	0.97±0.062	0.94±0.053	0.084*	0.95±0.059	0.96±0.051	0.714*
Systolic blood pressure (mmHg; mean±SD)	120±13.5	122±13.8	0.669*	120±12.8	124±15.8	0.178*
Diastolic blood pressure (mmHg; mean±SD)	82±6.6	80±7.7	0.250*	79±7.3	82±7.6	0.179*
Neutrophil (10 ⁶ /L; mean±SD)	53.7±8.05	52.5±8.34	0.543*	53.5±8.58	51.6±7.50	0.312*
Lymphocyte (10 ⁶ /L; mean±SD)	35.5±8.24	36.1±7.83	0.740*	35.2±7.98	37.4±7.68	0.224*
Monocyte (10 ⁶ /L; mean±SD)	7.2±1.14	7.6±1.68	0.203*	7.5±1.66	7.6±1.37	0.714*
Resting HR (bpm; mean±SD)	64.8±8.46	61.8±9.64	0.116*	62.6±9.38	62.7±9.54	0.989*
Medical history						
Hypertension (n)	6	7	0.097**	6	7	0.097**
Dyslipidemia (n)	10	18	0.169**	16	12	0.172**
Diabetes mellitus (n)	1	1	0.473**	0	2	0.114**
Family-history of CAD (n)	2	1	0.179**	2	1	0.179**
Smoking (n)	1	5	0.476**	5	1	0.328**
Lipid levels						
Total cholesterol levels (mg/dL; mean±SD)	222±35.18	203±40.01	0.038*	207±41.3	213±35.8	0.505*
HDL cholesterol levels (mg/dL; mean±SD)	55.3±17.60	58.9±16.75	0.361*	58.4±16.81	57.1±17.52	0.743*
TC to HDL ratio	4.4±3.62	3.66±1.04	0.010*	3.7±1.11	4.00±1.17	0.367*
Average cycling distance in previous month (km; median (range))	675 (214-1,220)	585 (170-1,360)	0.715***	675 (214-1,220)	585 (170-1,360)	0.715***
Long-distance cycling experience						
Yes (n)	20 (83.3%)	51 (79.7%)	0.479**	12 (20.7%)	5 (16.7%)	0.441**
Never (n)	4 (16.7)	13 (20.3%)		46 (79.3%)	25 (83.3%)	
Exercise characteristics						
Cycling distance NC 240K (n)	7 (29.2%)	21 (32.8%)		19 (32.7%)	9 (30%)	
Cycling distance TdB 140K (n)	9 (37.5%)	21 (32.8%)		24 (41.4%)	6 (20%)	
Cycling distance TdB 100K (n)	8 (33.3%)	22 (34.4%)		15 (25.9%)	15 (50%)	
Exercise duration (hours; median (range))				7.5 (3.5-9.8)	6.3 (3.5-8.7)	0.161***
Mean HR (bpm; median (range))				140 (94-180)	134 (96-158)	0.008***
Peak HR (bpm; median (range))				175 (128-229)	168 (116-233)	0.444***
Exercise intensity (%; median (range))				82.8 (53.73-92.16)	80.7 (45.6-90.8)	<0.001***

(17) These findings suggest that anthropometry parameters are related to baseline hs-CRP and other inflammation biomarkers. We also found that BMI was associated to post-exercise hs-CRP elevation as well. Inflammation is more prominent in individuals with CVD risk factors include high BMI. It is well known that obesity associated with inflammation through the mechanism of platelet activation (20) and lipoinflammation.(21)

A higher total cholesterol and lower HDL cholesterol was found in cyclists with baseline hs-CRP ≥ 3mg/L. It is well known that long-term exercise training increases HDL-cholesterol levels but decreases BMI, IL-6 and hs-CRP levels.(22-24) So less trained individuals are expected to demonstrate higher resting concentrations of hs-CRP. (22,25) Indeed, a Chinese study reported that hs-CRP and HDL cholesterol levels were inversely associated with




Figure 2. Baseline and post-exercise hs-CRP levels.

the dose of exercise session (23) and the dose of training intensity (22,25). These studies concluded that the least trained individuals have higher hs-CRP and lower HDL cholesterol compared to the most trained individuals.(24) Another study in Austria reports that heavy workers tend to have better oxidation status, determined by total oxidant capacity and MDA levels compared to office workers. (26) The study in the United States of America and Brazil also concluded that higher physical activity levels tend to have lower hs-CRP levels.(27,28) The previous studies concluded that the more intensity of exercise session the better status of hs-CRP levels and lipid profile. However, in this study based on touring groups, there were no significant differences in baseline hs-CRP levels, but the lipid levels remain significantly different. This condition can be explained that training habit characteristics between the groups were almost comparable, impacts on hs-CRP levels status accordingly (Table 3). The condition why lipid levels remain different between groups, leaving unclear explanation and further study must be conducted.

The current study also confirmed that exercise-induced hs-CRPelevations are common following prolonged cycling. Almost all groups have significantly different post-exercise and baseline hs-CRP levels, except TdB100K group (Figure 2). This finding may directly relate to exercise duration and intensity as TdB100K participants cycled for 5.0 (3.5-6.5) hours at 74.5% (45.58-90.98) of HRmax, whereas the other groups cycled longer and at a higher intensity (Table 2).

Exercise-induced hs-CRP elevation has been widely reported and attributed to the acute heavy physical activity. (29,30) In response to heavy physical activity, inflammation can provoke the production of monocyte tissue factor, stimulate hyperreactivity of platelet, promote biosynthesis of fibrinogen, and enhance formation of the microparticle and erythrocyte aggregability, thus agitate prothrombotic state.(29) Present study demonstrated a significant increase of 0.9 mg/L of post-exercise hs-CRP level over the baseline.

A study in Turkey reported that after exercise stress test there was a significant increase of 0.5 mg/L of post-exercise stress test hs-CRP level over the baseline in subjects with obstructive coronary lesions, while there was no increase in subjects without obstructive coronary lesions.(31)

The study in Turtu concluded exercise induced a significant increment in inflammatory markers *i.e.*, white blood count, hs-CRP, IL-6, IL-10 and MCP-1 concentrations. The long-distance sculling intensity variables such as the average rating of perceived exertion, HR and blood lactate were correlated with changes in IL-8, IL-1 α and IL-1 β levels. Maximal aerobic performance variables (maximum rate of oxygen consumption (VO₂ max) and maximal aerobic power) were related to changes in IL-2, IL-4, IL-8 and IL-1 β levels.(30) It was also clearly stated in a systematic review that acute moderate and heavy physical exercise influences in the elevation of pro-inflammatory *i.e.*, IL-6, TNF- α , IL-8 and hs-CRP and lowering the anti-inflammatory markers *i.e.*, IL-1, IL-10 and adiponectin, while regular moderate physical exercise will make favorable effects in lowering pro-inflammatory markers and increasing anti-inflammatory markers.(29)

The current study also showed that exercise intensity significantly influenced the magnitude of post-exercise hs-CRP elevations. This finding is in accordance with an American study reporting that hs-CRP elevations were directly linked to exercise intensity.(32) We also found

Table 4. Regression logistic model for post-exercise hs-CRP elevation.

Variables	β	OR adjuste d	95% CI	p
BMI	0.216	1.241	1.043-1.476	0.015
Age	0.034	1.034	0.989-1.082	0.137
Cycling distance	-1.527	0.217	0.062-0.758	0.017
Constant	-6.806			0.007

that cycling distance was related to post-exercise hs-CRP elevations. The training status or training volume of these subgroups may be responsible for these findings as a previous study found these characteristics to be predictive for changes in hs-CRP levels following exercise.(32) We also must take notice that training volume influences to the exercise intensity, as seen in this study, NC240K group had relative higher training volume and adapted longer duration and distance with lower exercise intensity than other groups. Similar results were found in both of these studies, however, the present study just evaluated hs-CRP while the previous study measured almost whole pro- and anti-inflammatory markers such as interferon- γ , IL-1b, IL-2, IL-8, IL-10 and IL-12p70, and granulocyte and monocyte phagocytosis (GR-PHAG and MO-PHAG).

Conclusion

In endurance cyclists, this study confirmed that body mass, BMI, waist circumference and lipid levels (total- and HDL-cholesterol) were associated with baseline hs-CRP levels, while BMI and cycling distance were associated with post-exercise hs-CRP elevations. Further research is needed to elucidate exercise-induced changes in pro- and anti-inflammatory biomarkers and to unravel the underlying mechanisms that are responsible for these responses.

Acknowledgements

Research was funded by the Directorate of Research and Public Service, Ministry of Research, Technology and Higher Education of the Republic of Indonesia (DIPA 042.06.1.401516). The authors would like to thank all the investigators and organizers of 2017 Indonesia North Coast and Tour de Borobudur cycling tours, and all the study participants.

References

1. Yousuf O, Mohanty BD, Martin SS, Joshi PH, Blaha MJ, Nasir K, *et al.* High-sensitivity C-reactive protein and cardiovascular disease: A resolute belief or an elusive link? *J Am Coll Cardiol.* 2013; 62: 397-408.
2. Shrivastava AK, Singh HV, Raizada A, Singh SK. C-reactive protein, inflammation and coronary heart disease. *Egypt Hear J.* 2015; 67: 89-97.
3. Kasapis C, Thompson PD. The effects of physical activity on serum C-reactive protein and inflammatory markers: a systematic review. *J Am Coll Cardiol.* 2005; 45: 1563-9.
4. Fallon KE. The acute phase response and exercise: The ultramarathon as prototype exercise. *Clin J Sport Med.* 2001; 11: 38-43.
5. Andersson J, Jansson JH, Hellsten G, Nilsson TK, Hallmans G, Boman K. Effects of heavy endurance physical exercise on inflammatory markers in non-athletes. *Atherosclerosis.* 2010; 209: 601-5.
6. Nunes RAB, Araújo F, Correia GF, Da Silva GT, Mansur AJ. High-sensitivity C-reactive protein levels and treadmill exercise test responses in men and women without overt heart disease. *Exp Clin Cardiol.* 2013; 18: 124-8.
7. Eijssvogels TMH, Fernandez AB, Thompson PD. Are there deleterious cardiac effects of acute and chronic endurance exercise? *Physiol Rev.* 2016; 96: 99-125.
8. Stanojevic D, Jakovljevic V, Barudic N, Zivkovic V, Srejsovic I, Parezanovic Ilic K, *et al.* Overtraining does not induce oxidative stress and inflammation in blood and heart of rats. *Physiol Res.* 2016; 65: 81-90.
9. Tidball JG. Inflammatory processes in muscle injury and repair. *Am J Physiol Regul Integr Comp Physiol.* 2005; 288: 345-53.
10. Peake JM, Suzuki K, Wilson G, Hordem M, Nosaka K, MacKinnon L, *et al.* Exercise-induced muscle damage, plasma cytokines, and markers of neutrophil activation. *Med Sci Sports Exerc.* 2005; 37: 737-45.
11. Wallberg L, Mikael Mattsson C, Enqvist JK, Ekblom B. Plasma IL-6 concentration during ultra-endurance exercise. *Eur J Appl Physiol.* 2011; 111: 1081-8.
12. Skadberg Ø, Kleiven Ø, Bjørkavoll-Bergseth M, Melberg T, Bergseth R, Selvåg J, *et al.* Highly increased Troponin I levels following high-intensity endurance cycling may detect subclinical coronary artery disease in presumably healthy leisure sport cyclists: The North Sea Race Endurance Exercise Study (NEEDED) 2013. *Eur J Prev Cardiol.* 2017; 24: 885-94.
13. Bessa AL, Oliveira VN, Agostini GG, Oliveira RJS, Oliveira ACS, White GE, *et al.* Exercise intensity and recovery: biomarkers of injury, inflammation, and oxidative stress. *J Strength Cond Res.* 2016; 30: 311-9.
14. Mouridsen MR, Nielsen OW, Carlsen CM, Mattsson N, Ruwald MH, Binici Z, *et al.* High-sensitivity C-reactive protein and exercise-induced changes in subjects suspected of coronary artery disease. *J Inflamm Res.* 2014; 7: 45-55.
15. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. *J Am Coll Cardiol.* 2001; 37: 153-6.
16. Ridker PM. High-sensitivity c-reactive protein, potential adjunct for global risk assessment in the primary prevention of cardiovascular disease. *Circulation.* 2001; 103: 1813-8.
17. Arbel Y, Birati EY, Shapira I, Finn T, Berliner S, Rogowski O. Comparison of different anthropometric measurements and inflammatory biomarkers. *Int J Inflamm.* 2012; 2012: 1-5. doi: 10.1155/2012/124693.
18. Zarzecczny R, Nawrat-Szoltysik A, Polak A, Manasar A, Maliszewski J, Kiełtyka A, *et al.* The relationship between selected body fatness indices and the level of blood interleukin-6 in female nursing home residents aged 80+ years without inflammation: A pilot study. *Exp Gerontol.* 2018; 108: 240-6.
19. Sigdel M, Kumar A, Gyawali P, Shrestha R, Tuladhar ET, Jha B. Association of high sensitivity C-reactive protein with the components of metabolic syndrome in diabetic and non-diabetic individuals. *J Clin Diagnostic Res.* 2014; 8: 11-3.
20. Samad F, Ruf W. Inflammation, obesity, and thrombosis. *Blood.* 2013; 122: 3415-22.
21. Izaola O, de Luis D, Sajoux I, Domingo JC, Vidal M. Inflammation and obesity (Lippoinflammation). *Nutr Hosp.* 2015; 31: 2352-8.

22. Sponder M, Kopecky C, Campean IA, Emich M, Fritzer-Szekeres M, Litschauer B, *et al.* Sports and HDL-quality reflected by serum amyloid A and surfactant protein B. *Int J Med Sci.* 2017; 14: 1040-8.
23. Tsao TH, Yang C Bin, Hsu CH. Effects of different exercise intensities with isoenergetic expenditures on C-reactive protein and blood lipid levels. *Res Q Exerc Sport.* 2012; 83: 293-9.
24. Rehnuma B, Hassan Z, Ibrahim M, Ali L. Serum levels of high sensitivity C-Reactive protein and its association with lipidemic status in Bangladeshi healthy adults. *J Pathol Nepal.* 2014; 4: 644-8.
25. Sponder M, Campean IA, Emich M, Fritzer-Szekeres M, Litschauer B, Bergler-Klein J, *et al.* Long-term endurance training increases serum cathepsin S and decreases IL-6 and hsCRP levels. *J Sports Sci.* 2017; 35: 2129-34.
26. Zelzer S, Tatzber F, Herrmann M. Work intensity, low-grade inflammation, and oxidative status: a comparison between office and Slaughterhouse Workers. *Oxid Med Cell Longev.* 2018; 2018: 2737563. doi: 10.1155/2018/2737563.
27. Femandes RA, Ritti-Dias RM, Balagopal PB, Conceição RDO, Santos RD, Cucato GG, *et al.* Self-initiated physical activity is associated with high sensitivity C-reactive protein: A longitudinal study in 5,030 adults. *Atherosclerosis.* 2018; 273: 131-5.
28. McFarlin BK, Flynn MG, Campbell WW, Craig BA, Robinson JP, Stewart LK, *et al.* Physical activity status, but not age, influences inflammatory biomarkers and toll-like receptor 4. *J Gerontol A Biol Sci Med Sci.* 2006; 61A: 388-93.
29. Chen YW, Apostolakis S, Lip GYH. Exercise-induced changes in inflammatory processes: Implications for thrombogenesis in cardiovascular disease. *Ann Med.* 2014; 46: 439-55.
30. Jurimae J, Tillmann V, Purge P, Jurimae T. Acute inflammatory response to prolonged sculling in competitive male rowers. *J Sport Med Phys Fit.* 2016; 56: 1368-75.
31. Basar N, Saatci Yasar A, Erden G, Cagli K, Sen N, Metin F, *et al.* The relation between exercise-induced myocardial ischemia and high-sensitive C-reactive protein and neopterin levels. *Anadolu Kardiyol Derg.* 2010; 10: 502-7.
32. Nieman DC, Konrad M, Henson DA, Kennerly K, Shanelly RA, Wallner-Liebmann SJ. Variance in the acute inflammatory response to prolonged cycling is linked to exercise intensity. *J Interf Cytokine Res.* 2012; 32: 12-7.

Baseline and post

ORIGINALITY REPORT

15%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to iGroup Student Paper	1%
2	Eijsvogels, Thijs M.H., James L. Januzzi, Beth A. Taylor, Stephanie K. Isaacs, Pierre D'Hemecourt, Amanda Zaleski, Sophia Dyer, Chris Troyanos, Rory B. Weiner, Paul D. Thompson, and Aaron L. Baggish. "Impact of Statin Use on Exercise-Induced Cardiac Troponin Elevations", The American Journal of Cardiology, 2014. Publication	1%
3	scholar.ui.ac.id Internet Source	1%
4	www.ncmls.nl Internet Source	1%
5	www.hindawi.com Internet Source	1%
6	icgh2016.ui.ac.id Internet Source	1%

7	wp1191596.server-he.de Internet Source	1%
8	anatoljcardiol.com Internet Source	<1%
9	transforming-science.com Internet Source	<1%
10	watermark.silverchair.com Internet Source	<1%
11	Submitted to Universitas Sebelas Maret Student Paper	<1%
12	Nilay Yuksel, Burak Tanriverdi, Betül Ipteç, Ozcan Erel. "Thiol-disulfide homeostasis as an oxidative stress marker in patients with Graves' ophthalmopathy", Orbit, 2018 Publication	<1%
13	A Guntur Hermawan. "Development of Immunopathobiogenesis on SIRS-Sepsis", The Indonesian Biomedical Journal, 2009 Publication	<1%
14	Jaak Jürimäe, Sille Vaiksaar, Priit Purge. "Circulating Inflammatory Cytokine Responses to Endurance Exercise in Female Rowers", International Journal of Sports Medicine, 2018 Publication	<1%
15	Isti Pudjihastuti, Noer Handayani, Siswo	

Sumardiono. "The Effect of Emulsifier and Hydrocolloid on Baking Expansion and Texture of Bread from Modified Cassava", MATEC Web of Conferences, 2018

Publication

<1%

16

Gach, O.. "Long-Term Prognostic Significance of High-Sensitivity C-Reactive Protein Before and After Coronary Angioplasty in Patients With Stable Angina Pectoris", The American Journal of Cardiology, 20070101

Publication

<1%

17

Ryszard Zarzeczny, Agnieszka Nawrat-Szołtysik, Anna Polak, Ahmed Manasar et al. "The relationship between selected body fatness indices and the level of blood interleukin-6 in female nursing home residents aged 80+years without inflammation: A pilot study", Experimental Gerontology, 2018

Publication

<1%

18

ard.bmj.com
Internet Source

<1%

19

Choirun Nissa, Martha Irene Kartasurya, Banundari Rahmawati. "Effects of Chlorophyll in Papaya Leaves on Superoxide Dismutation and Blood Glucose Level of Diabetic Rats", Makara Journal of Health Research, 2015

Publication

<1%

20 Lene Rørholm Pedersen, Rasmus Huan Olsen, Christian Anholm, Arne Astrup et al. "Effects of 1 year of exercise training versus combined exercise training and weight loss on body composition, low-grade inflammation and lipids in overweight patients with coronary artery disease: a randomized trial", Cardiovascular Diabetology, 2019
Publication <1%

21 Juho Tynkkynen, Jussi A. Hernesniemi, Tiina Laatikainen, Aki S. Havulinna et al. "High-sensitivity cardiac troponin I and NT-proBNP as predictors of incident dementia and Alzheimer's disease: the FINRISK Study", Journal of Neurology, 2016
Publication <1%

22 otorhinolaryggologos.blogspot.com
Internet Source <1%

23 ejournal2.undip.ac.id
Internet Source <1%

24 Y. FENG. "Effect of Agmatine on the Time Course of Brain Inflammatory Cytokines After Injury in Rat Pups", Annals of the New York Academy of Sciences, 12/01/2003
Publication <1%

25 Isti Pudjihastuti, Noer Handayani, Siswo <1%

Sumardiono. "Effect of pH on Physicochemical Properties of Cassava Starch Modification Using Ozone", MATEC Web of Conferences, 2018

Publication

26

www.japsonline.com

Internet Source

<1%

27

Kevin Zwetsloot, Casey John, Marcus Lawrence, Rebecca Battista, R. Andrew Shanely. "High-intensity interval training induces a modest systemic inflammatory response in active, young men", Journal of Inflammation Research, 2014

Publication

<1%

28

journal.unnes.ac.id

Internet Source

<1%

29

www.spandidos-publications.com

Internet Source

<1%

30

"Abstracts", Catheterization and Cardiovascular Interventions, 2007

Publication

<1%

31

bmcpublikealth.biomedcentral.com

Internet Source

<1%

32

Katsumi Abo, Takaya Mio, Kimiaki Sumino. "Comparative analysis of plasma and erythrocyte 7-ketocholesterol as a marker for oxidative stress in patients with diabetes

<1%

33

Nathalie Van Aerde, Philippe Meersseman, Yves Debaveye, Alexander Wilmer et al. "Five-year impact of ICU-acquired neuromuscular complications: a prospective, observational study", Intensive Care Medicine, 2020

Publication

<1%

34

Te Hung Tsao, Chang Bin Yang, Chin Hsing Hsu. "Effects of Different Exercise Intensities With Isoenergetic Expenditures on C-Reactive Protein and Blood Lipid Levels", Research Quarterly for Exercise and Sport, 2012

Publication

<1%

35

savoirs.usherbrooke.ca

Internet Source

<1%

36

edoc.pub

Internet Source

<1%

37

jmscr.igmpublication.org

Internet Source

<1%

38

www.docstoc.com

Internet Source

<1%

39

Leonid Roytblat, Maxim Rachinsky, Allan Fisher, Lev Greemberg, Yoram Shapira, Amos Douvdevani, Simon Gelman. "Raised Interleukin-6 Levels in Obese Patients", Obesity

<1%

Research, 2000

Publication

40

synapse.koreamed.org

Internet Source

<1%

41

Vincent L. Aengevaeren, Arend Mosterd, Thijs L. Braber, Niek H.J. Prakken et al. "Relationship Between Lifelong Exercise Volume and Coronary Atherosclerosis in Athletes", *Circulation*, 2017

Publication

<1%

42

D. Aronson. "Association between fasting glucose and C-reactive protein in middle-aged subjects", *Diabetic Medicine*, 1/2004

Publication

<1%

43

Fadzlina, AA, Fatimah Harun, MY Nurul Haniza, Nabilla Al Sadat, Liam Murray, Marie M Cantwell, Tin Su, Hazreen Majid, and Muhammad Jalaludin. "Metabolic syndrome among 13 year old adolescents: prevalence and risk factors", *BMC Public Health*, 2014.

Publication

<1%

44

Raul A. Martins, Ana P. Neves, Manuel J. Coelho-Silva, Manuel T. Veríssimo, Ana Maria Teixeira. "The effect of aerobic versus strength-based training on high-sensitivity C-reactive protein in older adults", *European Journal of Applied Physiology*, 2010

<1%

45

Stefanos Zafeiropoulos, Ioannis Farmakis, Anastasios Kartas, Alexandra Arvanitaki et al. "Reinforcing Adherence to Lipid-lowering Therapy after an Acute Coronary Syndrome: A Pragmatic Randomized Controlled Trial", Research Square, 2020

Publication

<1%

46

G McGwin. "The relation between C reactive protein and age related macular degeneration in the Cardiovascular Health Study", British Journal of Ophthalmology, 2005

Publication

<1%

47

Mohammad Nasir Pour Eshmanan Talemi, Seyed Mojtaba Paydar Ardakani, Behnam Roozbeh. "Tribulus Terrestris may decrease muscle damage markers following a high-intensity resistance exercise: a pilot study", International Journal for Vitamin and Nutrition Research, 2020

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Baseline and post

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
