

Path Analysis of Blood Glucose Determinant on Diabetes Mellitus Patients through Intervening Variables of Medication Adherence

by Sri Ratna Rahayu

Submission date: 12-May-2023 09:15AM (UTC+0700)

Submission ID: 2090927704

File name: tients_through_Intervening_Variables_of_Medication_Adherence.pdf (225.01K)

Word count: 4239

Character count: 22073



Path Analysis of Blood Glucose Determinant on Diabetes Mellitus Patients through Intervening Variables of Medication Adherence

Oktia Woro Kasmini Handayani^{1✉}, Pamela Lindasari¹, Sri Ratna Rahayu¹, Ari Yuniastuti², Efa Nugroho³

¹ Postgraduate Program in Public Health, Semarang State University

² Department of Biology, Faculty of Mathematics and Natural Sciences, Semarang State University

³ Department of Public Health, Faculty of Sport Sciences, Semarang State University

Article Info

Article History:
Submitted May 2019
Accepted June 2019
Published July 2019

Keywords:
Family Support, Adherence to taking drugs, Blood Sugar, Knowledge

Abstract

The prevalence of Diabetes Mellitus (DM) patients in Indonesia in 2013 has increased compared to 2007. The non-adherence of patients in undergoing treatment therapy is one of the drug therapy problems (DTP) that needs special attention. The purpose of the study was to obtain the results of an analysis of direct and indirect relationships related to knowledge, attitudes, family support, and adherence to taking medication for blood glucose levels in patients with DM of type II. Research with correlation analytic research design, with a population of DM patients of type II with outpatient care in the Sleman region, DIY, were 530 people. The sample was determined by the proportionate stratified random sampling technique of 119 people. The independent variables include knowledge, attitude, family support, while the dependent variable is blood glucose levels in patients with DM of type II and the intervening variable is adherence with taking medication. This study used path analysis techniques. The results of this study draw conclusions, knowledge, attitudes, family support and medication adherence to DM patients with directly influence blood glucose levels. Apart from knowledge, attitudes and family support through adherence to taking drugs indirectly affect blood glucose levels.

INTRODUCTION

The World Health Organization (WHO) predicts that there will be an increase in the number of diabetics from 135 million in 1995 to 300 million in 2025. The highest increase is predicted to occur in the Continent of Asia (Zahra et al., 2016). Diabetes mellitus (DM) is rapidly becoming one of the most common non-infectious diseases globally. Population growth, population aging, and urbanization with associated lifestyle changes are likely to cause 55% increase worldwide, with diabetes by 2035. China has become the country with the largest number of

diabetics in the world (Sugang et al, 2015). Based on WHO data in 2013, the prevalence of DM patients in Indonesia in 2013 (2.1%) experienced an increase compared to 2007 (1.1%). The most common type of DM and the increasing prevalence is type II diabetes mellitus with the most cases being 90% of all cases of DM in the world.

DM is a disease associated with lifestyle, so the success or failure of the management of DM is very dependent on the patient himself in changing his behavior. Rosyida et al. (2015), stated that patient non-adherence to undergoing treatment thera-

py is one of the Drug Therapy Problems (DTP) that needs special attention. In addition, the research of Meidikayanti & Chatarina (2017) stated that family support is believed to have an influence on the quality of life of people with DM. The patient's knowledge about DM is a tool that can help patients with diabetes management (Perdana, 2013). The results of the study of Amalia et al. (2016), stated that the level of knowledge of DM patients in the less category, so most experienced symptoms of DM but they did not know that it was signs of DM. They also do not know what things can increase sugar levels. The results of the study state that almost half of people with DM of type II have a level of knowledge that is less to bad and less satisfying (Okatiranti & Nur'aeni, 2016; Anju et al., 2015; Kaniz et al., 2017). In addition, research of Herath et al. (2017) found that attitudes toward diabetes were mostly poor (90%) and the level of education did not significantly influence attitudes.

Previous studies conducted in the study area showed that 11% of knowledge about DM disease was lacking, and lack of support from the family, because family members were busy with their work, and 70% were not obedient to taking medication for forgetting or being lazy. The purpose of this study was to obtain the results of an analysis of direct and indirect relationships related to knowledge, attitudes, family support, and adherence to taking medication for blood glucose levels in DM patients of type II.

METHODS

This study used a quantitative approach with correlation analytic research design, using a cross sectional study approach. The population in this study were all DM patients of type II with outpatient care in the Sleman region, DIY, was an area with a high DM case of 530 people. The sample was determined using the proportionate stratified random sampling technique from 4 work areas of the community health center. The sample size was determined by the Slovin formula (Nursalam, 2013), so that a sample of 119 people was determined. Independent variables include knowledge, attitudes, family support, while the dependent variable is blood sugar levels in patients with DM of type II and the intervening variable is adherence to taking medication. The instrument used for primary data is a questionnaire that contains knowledge, attitudes, family support and adherence of respondents in taking medication, while secondary data was obtained from medical record documents to obtain respondents' blood sugar levels. This study used path analysis techniques.

RESULTS AND DISCUSSION

The results showed that respondents' characteristics as illustrated in Table 1. as many as 56-65 years old (45.4%), senior high school graduation (53.8%), good knowledge (84%), good attitude (82.4%), Good family support (69.7%), medication compliance (65%) and controlled blood sugar (57.1%).

Based on the results of the normality and linearity test, it was concluded that the research data were normally distributed and there was a linear relationship between the independent variables consisting of knowledge, attitudes, family support, and adherence to taking drugs with the dependent variable namely blood sugar levels, so path analysis which is measured by multiple linear regression analysis. Then after testing the model validation with total determinant coefficients and theory trimming, the diagram is as follows:

Interpretation of Figure 1 is as follows:

Direct influence X_1 to $Y = P_4$, the influence of knowledge variables (X_1) the variable of blood glucose levels (Y) are 0,193.

Direct influence X_2 to $Y = P_5$, the influence of attitude variable (X_2) on the variable of blood glucose level (Y) is 0.203.

Direct influence X_3 to $Y = P_6$, the effect of family support variable (X_3) on the variable of blood glucose level (Y) is 0.217.

Direct influence X_4 to $Y = P_7$, the effect variable of the adherence to taking medication (X_4) on the variable of blood glucose level (Y) is 0,223.

Indirect influence X_1 to Y through $X_4 = P_1 \times P_7$. The effect of the knowledge variable (X_1) on the variable of blood glucose level (Y) through medication adherence (X_4) is $0,290 \times 0,223 = 0,065$.

Indirect influence X_2 to Y through $X_4 = P_2 \times P_7$. The effect of attitude variable (X_2) on the variable of blood glucose level (Y) through medication adherence (X_4) is $0,274 \times 0,223 = 0,0611$.

Indirect influence X_3 to Y through $X_4 = P_3 \times P_7$. Effect of family support variables (X_3) the variable of blood glucose level (Y) through medication compliance (X_4) is $0,275 \times 0,223 = 0,0613$.

Total Influence X_1 to Y through X_4 , $Y = P_4 + (P_1 \times P_7) = 0,193 + 0,065 = 0,258$

Total Influence X_2 to Y through X_4 , $Y = P_5 + (P_2 \times P_7) = 0,203 + 0,061 = 0,264$

Total Influence X_3 to Y through X_4 , $Y = P_6 + (P_3 \times P_7) = 0,217 + 0,061 = 0,278$

Direct Effects and Indirect Effects of Knowledge on Blood Glucose Levels

Based on the results of path analysis, it was shown that knowledge proved to have a direct effect on blood glucose levels with a path coefficient

Table 1. Overview of the Distribution of Respondents Characteristics

Respondents Characteristics	Total	Percentage
Age		
36 - 45 years old	8	6.7%
46 - 55 years old	28	23.5%
56 - 65 years old	54	45.4%
> 65 years old	29	24.4%
Education		
Graduate of Master Degree	2	1.7%
Graduate of Bachelor Degree	17	14.3%
Graduate of Diploma 3	9	7.6%
Graduate of Senior High Graduate	64	53.8%
Graduate of Junior High School	16	13.4%
Graduate of Elementary School	11	9.2%
Knowledge		
Good	100	84%
Enough	19	16%
Less	0	0.0%
Attitude		
Good	98	82.4%
Enough	17	14.3%
Less	4	3.4%
Family support		
Good	83	69.7%
Enough	34	28.6%
Less	2	1.7%
Adherence to taking drugs		
Obey	91	76.5%
Disobey	28	23.5%
Blood Sugar Levels		
Controlled	68	57.1%
Uncontrolled	51	42.9%

of 0.193 and knowledge was proven to influence blood glucose levels through medication adherence to a path coefficient of 0.065 with a total effect of 0.253. Good knowledge related to health problems will affect the way of view which then influences a person's behavior on the health problems they face, the better level of knowledge of a patient has about DM and its control, the more controlled his blood glucose levels are.

In this study the patient's knowledge is good and directly influences the blood sugar level of 19.3%, besides that knowledge through medication adherence will have a greater influence on blood glucose levels which is equal to 25.8%, where adherence to taking medication from patients at this research is 76.5% is good. Knowledge can directly affect the patient's blood sugar levels, because of other factors,

such as diet, activity or sports factors, and stress factors, so the value of the influence obtained is smaller (19.3%). Knowledge can also influence adherence to taking medication which is a variable intervening, then it will affect blood sugar levels, so the total effect value is greater (25.8%), although there are several other factors that can influence such as family support or other support in the environment.

Knowledge is the basis for health behavior, especially the lack of knowledge about the rules of treatment, the benefits and complications caused by not adhering to the treatment recommendations. This is in line with studies that state that good knowledge related to knowledge about diabetes mellitus, both signs and symptoms and its handling will affect the control of blood glucose levels (Kassahun & Alemayehu, 2017; Qurratuaeni, 2010). The results




Figure 1. Path Analysis Diagram

Other studies say that ignorance of the treatment of diabetes will cause a lack of control of blood sugar levels and can trigger distress conditions that have an impact on rising blood sugar levels (Fisher et al., 2010).

New behavior, especially in adults, begins in the cognitive domain in the sense that the subject knows first about the stimulus in the form of material objects outside, giving rise to an inner response in the form of attitude. Finally, the stimulus that is the object that has been known and fully realized will cause a further response, namely in the form of an action against a stimulus or object. Knowledge is the first step for someone to determine their attitudes and behavior. So the level of knowledge will greatly influence the acceptance of a program (Notoatmodjo, 2007). Because of the importance of knowledge that must be known by DM patients, various ways are done to be able to transfer health education and patient behavior guidance that can reach large areas, including coastal areas (Gunn et al., 2011). According to Alsous et al. (2019), the need for more educational interventions to overcome negative attitudes and promote healthy lifestyle practices and routine health checks, especially in the subgroup of patients with certain characteristics, such as low education is very important.

Direct Effects and Indirect Effects of Attitudes on Blood Glucose Levels

Based on the results of path analysis, it was shown that attitudes proved to have a direct effect on blood glucose levels with a path coefficient of

0.203, besides that attitudes also influence blood glucose levels through medication compliance with a path coefficient of 0.061, with a total effect of 0.264. Attitude is a reaction or response that is still closed from someone to the stimulus. Attitudes will not necessarily be realized in the form of an action, supporting factors are needed or conditions are possible, such as facilities. A DM patient who has intended to go on a diet as recommended, sometimes gets out of the lane because the situation at home or office is not supportive.

The results of this study are in line with the results of previous studies, such as the results of a study by Al-Maskari et al. (2013), Roux et al. (2018), Ansari et al. (2019), who said that poor levels of knowledge and attitudes related to patient control of DM, which reflected in poor glycemic control and can contribute to morbidity and mortality. Although many research results say knowledge, and attitudes influence patients in controlling their blood sugar, but to improve the ability of DM patients to manage their own blood sugar, they need to think about demographics, education and age and economic conditions (Kheir et al., 2011).

Direct and Indirect Effects of Family Support on Blood Glucose Levels

Based on the path analysis results showed that family support proved to have a direct effect on blood glucose levels with a path coefficient of 0.217, besides that family support was proven to influence blood glucose levels through medication adherence to a path coefficient of 0.061 and a total influence

of 0.278. Family is the social environment that is the closest to DM patients so that it is expected to help, control and shape the behavior of DM patients, including in this case self-management behavior. Family support plays a crucial role in compliance with self-management and indirectly affects the metabolic control. Family support is very important to motivate patients in an effort to create a stress-free environment due to the treatment they undergo. Family social support is as a protector in stress trigger factors and creates a comfortable environment so as to maintain blood sugar control. So that Diabetic Self-management Education (DMSE), which is a process of providing health education to DM patients, regarding the application of self-care strategies independently is to optimize metabolic control, prevent complications and improve the quality of life of patients.

In line with the research of Rosland & Piette (2011), stated that family support can affect the health of patients with chronic diseases, including DM. Communication patterns and good family coping mechanisms can increase the motivation of clients to always maintain their health. Family support is defined as part of social support, is a form of interaction between individuals who provide physical and psychological comfort through the fulfillment of the need for affection and security, which can be empathetic, encouragement, facilitative and participative (Hensarling, 2009). The results of the study concluded that the relationship of family support with medication adherence showed a moderate positive ($r=0.527$), which means that the higher the value of family support, the higher the compliance value of DM patients in undergoing treatment, including the relationship between family social support and patient compliance in implementing diet program (Nurleli, 2016; Anggina et al., 2010). Research of Awodele & Osuolale (2015), explained that there is a significant relationship between patient age, gender, and adherence to treatment which needs to be taken into consideration especially related to health education and counseling that can produce levels of adherence and improvement of clinical parameters

Good and correct treatment therapy if supported by patient compliance will be very beneficial for the patient themselves, both in terms of health and costs that must be incurred. Adherence of patients in taking drugs is important, especially for patients who have to take drugs for a long time or for life like DM sufferers. Therefore, the efficiency and effectiveness of adherence to drug use are important factors to consider (Rochana, 2009). A new nonpharmacological and pharmacological approach needs to be considered that has a clinically signifi-

cant and sustained long-term impact on patient adherence. The goal of treatment for DM of type 2 is to prevent or delay complications and maintain quality of life, so it is very necessary to use a patient and environment-centered approach to increase patient involvement in self-care activities (Davies et al., 2018).

The availability of drugs in patients that includes daily, weekly, or even monthly doses must be available in the near future, in order to increase adherence to, resulting in increased glycemic control, reduced complications of diabetes, and health care with minimal costs can be reduced. Aymen et al. (2017), explained that the longer duration of illness, old age, female sex, and self-perception and patient's belief about the severity of DM were factors associated with adherence to treatment. Increasing interaction and having partner relationships with patients is key to improving patient adherence to drugs.

The direct effect of medication adherence is an influence with the largest total value (22.3%), because for people with DM, drugs are a very influential external factor for controlling blood sugar levels, besides there are other factors that influence, such as diet done. The indirect effect on blood sugar levels with the greatest value is the effect of family support on blood sugar levels through adherence to medication (27.8%). Support from the environment of people with DM, especially the family environment is a factor that needs attention because the characteristics of the largest respondents in the age group are 56-65 years (45.4%), including the age group that begins to depend on the family environment.

CONCLUSIONS

Based on the research conducted, the conclusions are as follows: 1) Knowledge, attitudes, family support and medication adherence in with DM patients directly influence blood glucose levels. 2) Knowledge through adherence to taking drugs indirectly affects the blood glucose level by 25.8%. 3) There is an influence of attitude through adherence to taking medication indirectly to blood glucose levels of 26.4%. 4) Family support through adherence to taking medication has an effect on blood glucose levels of 27.8%.

REFERENCES

- Al-Maskari, F., El-Sadig, M., Al-Kaabi, J.M., Afandi, B., Nagelkerke, N., Yeatts, K.B. 2013. Knowledge, Attitude and Practices of Diabetic Patients in the United Arab Emirates. *PLoS ONE*, 8(1): e52857.
- Alsous, M., Jalil, M.A., Odeh, M., Al Kurdi, R., & Alnan, M. 2019. Public knowledge, attitudes and practices toward diabetes mellitus: A

- cross-sectional study from Jordan. *PLoS ONE*, 14(3): e0214479.
- Amalia, W.C., Sutikno, E., & Nugraheni, R. 2016. Hubungan antara Tingkat Pengetahuan tentang Diabetes Mellitus dan Gaya Hidup dengan Tipe Diabetes Mellitus di Puskesmas Wonodadi Kabupaten Blitar. *Jurnal Preventia*, 1(1): 14-20.
- Anggina, L.L., Hamzah, A., & Pandhit. 2010. Hubungan antara Dukungan Sosial Keluarga dengan Kepatuhan Pasien Diabetes Mellitus dalam Melaksanakan Program Diet di Poli Penyakit Dalam RSUD Cibabat Cimahi. *Jurnal Penelitian Kesehatan Suara Forikes*, 2(1): 1-9.
- Anju, G., Bhatta, D.N., & Aryal, U.R. 2015. Diabetes related health knowledge, attitude and practice among diabetic patients in Nepal. *BMC Endocrine Disorders*, 15(25): 1-8.
- Ansari, T, Sami, W., Alsubaie, N.R., Althaqib, A.A., Alenezi, A.A., Almutairy, A.N., Alshamrani, M.A., Alanazi, M.A., Almutairi, M.A. 2019. Assessment of knowledge, attitude, and practice of dietary pattern in patients with type 2 diabetes mellitus: a locality-based perspective study. *International Journal of Medicine in Developing Countries*, 3(7): 581–585.
- Awodele, O., & Osuolale, J.A. 2015. Medication Adherence in Type 2 Diabetes Patients: Study of Patients in Alimosho General Hospital, Igan-do, Lagos, Nigeria. *Journal African Health Sciences*, 15(2): 513-522.
- Aymen, E., Radwan, M., Al-Sharif, H., & Mustafa, A.A. 2017. Medications Adherence and Associated Factors among Patients with Type 2 Diabetes Mellitus in the Gaza Strip, Palestine. *Frontiers in Endocrinology*, 8(100): 1-9.
- Davies, M., D'Alessio, D., Fradkin, J., Kernan, W., Mathieu, C., et. al. (2018). Management of hyperglycaemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia*, 61(12).
- Fisher, L., Glasgow, R. E., & Strycker, L.A. 2010. The Relationship between Diabetes Distress and Clinical Depression with Glycemic Control Among Patients with Type 2 Diabetes. *BMC Public Health*, 33(5).
- Hensarling. 2009. *Development and Psychometric Testing of Hensarling's Diabetes Family Support Scale*. Doctoral Thesis. Denton: Graduate School of the Texas Woman's University.
- Herath, H.M.M., Weerasinghe, N.P., Dias., H., & Weeraratna, T.P. 2017. Knowledge, Attitude and Practice Related to Diabetes Mellitus among the General Public in Galle District in Southern Sri Lanka: A Pilot Study. *BMC Public Health*, 17(535): 1-7.
- Kaniz, F., Hossain, S., Natasha, K., Chowdhury, H.A., Akter, J., Khan, T., & Ali, L. 2017. Knowledge Attitude and Practice Regarding Diabetes Mellitus among Nondiabetic and Diabetic Study Participants in Bangladesh. *BMC Public Health*, 17(364): 1-10.
- Kassahun, C., & Alemayehu, G. 2017. Knowledge, Attitude, Practices and Their Associated Factors towards Diabetes Mellitus among Non Diabetes Community Members of Bale Zone Administrative Towns, South East Ethiopia: A Cross-sectional Study. *Journal Pone*, 12(2): 1-18.
- Kheir, N., Greer, W., Yousif, A., Al Geed, H., & Al Okkah, R. 2011. Knowledge, attitude and practices of Qatari patients with type 2 diabetes mellitus. *International Journal of Pharmacy Practice*, 19(3).
- Meidikayanti & Chatarina. 2017. Hubungan Dukungan Keluarga dengan Kualitas Hidup Diabetes Melitus Tipe 2 di Puskesmas Pademawu. *Jurnal Berkala Epidemiologi*, 5(2): 240-252.
- Notoatmodjo, S. 2007. *Promosi Kesehatan dan Ilmu Perilaku*. Jakarta: Rineka Cipta
- Nurleli. 2016. Dukungan Keluarga dengan Kepatuhan Pasien Diabetes Melitus dalam Menjalani Pengobatan di BULD RSUD ZA Banda Aceh. *Idea Nursing Journal*, 7(2).
- Nursalam. 2013. *Konsep Penerapan Metode Penelitian Ilmu Keperawatan*. Jakarta: Salemba Medika.
- Okatiranti & Nur'aeni. 2016. Gambaran Pengetahuan, Sikap dan Kepercayaan terhadap Diet Penderita DM di RSUD Kota Bandung. *Jurnal Ilmu Keperawatan*, 4(1).
- Perdana, Burhanudin, & Devi. 2013. Hubungan Tingkat Pengetahuan Tentang Penyakit DM dengan Pengendalian Kadar Glukosa Darah pada Pasien DM Tipe II di RSUD PKU Muhammadiyah Surakarta. *Jurnal Biomedika*, 5(2): 17-21.
- Quinn, C.C., Shardell, M.D., Terrin, M.L., Barr, E.A., Ballew, S.H., & Gruber-Baldini, A.L. 2011. Cluster-Randomized Trial of a Mobile Phone Personalized Behavioral Intervention for Blood Glucose Control. *Diabetes Care*, 34(9): 1934-1942.
- Qurratuani. 2010. *Faktor-faktor yang berhubungan dengan terkontrolnya kadar gula darah pada pasien diabetes melitus di RSUP Fatmawati Jakarta tahun 2009*. Jakarta: UIN Syarif Hidayatullah.
- Rochana, R.N. 2009. *Evaluasi kepatuhan pasien pen-*

- gobatan obat hipoglikemik oral bagi penderita diabetes mellitus tipe 2 pada pasien rawat jalan di RSUD.dr. Moewardi Surakarta*. Master Thesis. Surakarta: Universitas Muhammadiyah Surakarta.
- Rosland, M.H., & Piette, J.D. 2012. The Impact of Family Behaviors and Communication Patterns on Chronic Illness Outcomes: A Systematic Review. *J Behav Med*, 35(2): 221–239.
- Rosyida, L., Priyandani, Y., Sulistyarini, A., & Yunita. 2015. Kepatuhan Pasien pada Penggunaan Obat Anti Diabetes dengan Metode Pill-Count dan MMAS-8 di Puskesmas Kedurus Surabaya. *Jurnal Farmasi Komunitas*, 2(2): 36-41
- Roux, M., Walsh, C., Reid, M., & Raubenheimer, J. 2018. Diabetes-Related Knowledge, Attitude and Practices (KAP) of Adult Patients with Type 2 Diabetes Mellitus in the Free State Province, South Africa. *South African Journal of Clinical Nutrition*, 2(8).
- Shugang, L., Guo., S, He, F., Zhang, M., He, J., Yan, Y., Ding, Y., Zhang, J., Liu, J., Guo, H., Xu, S., & Ma, R. 2015. Prevalence of Diabetes Mellitus and Impaired Fasting Glucose, Associated with Risk Factors in Rural Kazakh Adults in Xinjiang, China. *International Journal of Environmental Research and Public Health*, 12: 554-565.
- Zahra, Dewi, & Miftakhurrahman. 2016. Gambaran Pelayanan Konseling Gizi pada Pasien Diabetes Melitus Tipe 2 di Puskesmas Jatinangor Tahun 2015. *JSK*, 1(4) : 186-192.

Path Analysis of Blood Glucose Determinant on Diabetes Mellitus Patients through Intervening Variables of Medication Adherence

ORIGINALITY REPORT

22%

SIMILARITY INDEX

18%

INTERNET SOURCES

11%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	repo.stikmuhptk.ac.id Internet Source	3%
2	www.scilit.net Internet Source	2%
3	midwifery.iocspublisher.org Internet Source	2%
4	core.ac.uk Internet Source	1%
5	F H Tyas, J G Nikita, D K Apriyanto, Mitrayana, M N Amin. " The Performance of CO Laser Photoacoustic Spectrometer In Concentration Acetone Detection As Biomarker For Diabetes Mellitus Type 2 ", Journal of Physics: Conference Series, 2018 Publication	1%
6	amrsjournals.com Internet Source	1%

Submitted to 60556

7	Student Paper	1 %
8	Elsa Budi Sihsilya Rahmawati, Poppy Farantia Saputri. "Study the Effect of Counseling on Early Initiation of Breastfeeding with Media Breastfeeding E-books", STRADA JURNAL ILMIAH KESEHATAN, 2019 Publication	1 %
9	Submitted to Forum Perpustakaan Perguruan Tinggi Indonesia Jawa Timur Student Paper	1 %
10	Submitted to iGroup Student Paper	1 %
11	"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020 Publication	1 %
12	Saiful Gunardi, Catur Septiawan. "Pengaruh Empat Variabel Terhadap Kualitas Hidup Lansia Penderita Diabetes Melitus Tipe II", Jurnal Ilmiah Ilmu Keperawatan Indonesia, 2018 Publication	1 %
13	Submitted to Universitas Negeri Semarang Student Paper	1 %
14	journals.plos.org Internet Source	1 %

15	Tuho Konsultasi Mendrofa, Ermi Girsang, Sri Lestari Ramadhani Nasution, Chrismis Novalinda Ginting. "Prevalence and Determinants of Drug Adherence in Type 2 Diabetes Mellitus Patients", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2022 Publication	1 %
16	pingpdf.com Internet Source	1 %
17	pps.org.pk Internet Source	1 %
18	repository.stikeshangtuah-sby.ac.id Internet Source	1 %
19	Submitted to Saint Joseph's College of Maine Student Paper	1 %
20	academicjournal.yarsi.ac.id Internet Source	1 %
21	Rhein Djunaid, Eka Pratiwi Puluhulawa, Muh.Nur Syukriani Yusuf. "The correlation between bullying and students learning achievement in SMP Negeri 6 Kota Gorontalo", Asian Community Health Nursing Research, 2019 Publication	<1 %
22	Anwar Rosyadi, Indri Hapsari, Pri Iswati Utami. "The Effect of Lifestyle, Spiritual, and Family	<1 %

Support on Diabetic Mellitus Patient with Chronic Kidney Disease Complication", PHARMACY: Jurnal Farmasi Indonesia (Pharmaceutical Journal of Indonesia), 2021

Publication

23

search.oecd.org

Internet Source

<1 %

24

T. Ralineba, M.L. Netshikweta, N.H. Shilubane. "Knowledge and Practices Associated with Diabetes among Patients with Chronic Diabetes Mellitus in Rural Areas of Vhembe District, Limpopo Province, South Africa", *Journal of Human Ecology*, 2017

Publication

<1 %

25

"Review of Environmental Aspects and Community Behavior in the Determination of Filariasis Risk Vulnerability Zone", 'Universitas Negeri Semarang'

Internet Source

<1 %

26

Muhammad Hatta, Dainty Maternity, Anissa Ermasari. "The Influence Of "Aku Bangga Aku Tahu" (Abat) Campaign On Adolescent Knowledge And Attitudes About Hiv/Aids", *Jurnal Kebidanan Malahayati*, 2022

Publication

<1 %

27

www.iosrjournals.org

Internet Source

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On