

**BUKTI KORESPONDENSI**  
**ARTIKEL JURNAL INTERNASIONAL BEREPUTASI**

Judul Artikel : Comparison of Organoleptic, Protein, Lipid and Flavonoid Content of  
Commercial Starter and Isolated Culture Red Dragon Fruit Peel Yogurt  
Jurnal : Food Research 4 (3) : 920-925 (Juni 2020)  
Penulis : Mardiana, Irwan Budiono, Natalia Desy Putriningtyas

No	Perihal	Tanggal
1.	Bukti article submission to “Food Research” Journal	12 November 2019
2.	Bukti technical check	26 November 2019
3.	Bukti article in Review	12 Desember 2019
4.	Bukti article accepted	15 Januari 2020
5.	Bukti pemberitahuan pembayaran (APC)	17 Januari 2020
6.	Bukti artikel production	11 Februari 2020
7.	Bukti pemberitahuan waktu penerbitan artikel	27 Februari 2020
8.	Bukti article published online	08 Juni 2020

**1. Bukti article submission to “Food Research  
Journal “(12 Nopember 2019)**

End now Vacation Settings

Gmail

Compose

Inbox 1,198

Starred

Snoozed

Sent

Drafts 73

More

Labels +

foodresearch.my@outlook.com

MANUSCRIPT SUBMISSION

Natalia Desy Putriningtyas <nataliadesy@mail.unnes.ac.id>  
to foodresearch.my

Tue, Nov 12, 2019, 12:21 AM

Dear Professor Dr. Son Radu  
Chief Editor Food Research

I would like to submit my manuscript and i hope this manuscript will be accepted  
thanks you

Best Regards,  
Natalia Desy Putriningtyas, S.Gz, M.Gizi  
Nutrition, S1 Public Health Science  
Sport Science Faculty Universitas Negeri Semarang

3 Attachments • Scanned by Gmail

- cover letter\_Natal...
- manuscript\_subm...
- Food research\_E...

## **2. Bukti Technical check (26-11-2019)**

End now Vacation Settings

**Gmail** foodresearch.my@outlook.com Active UNNES

Compose

Inbox 1,198  
Starred  
Snoozed  
Sent  
Drafts 73  
More

Labels +

**Status of my manuscript (Manuscript ID: FR-2019-380)** Inbox x

**Natalia Desy Putriningtyas** <nataliadesy@mail.unnes.ac.id>  
to foodresearch.my  
Tue, Nov 26, 2019, 9:22 AM

Dear Professor Dr. Son Radu  
Chief Editor Food Research

I would ask you about our manuscript with title Effect of Variation Red Dragon Fruit Peel in Yoghurt on Organoleptic Properties, Protein, Lipid, and Flavonoid Content (Manuscript ID: FR-2019-380). Whether our manuscript was accepted or not?  
Thank you for your information

—  
Best Regards,  
Natalia Desy Putriningtyas, S.Gz, M.Gizi  
Nutrition, S1 Public Health Science  
Sport Science Faculty Universitas Negeri Semarang

**Food Research** <foodresearch.my@outlook.com>  
to me  
Wed, Nov 27, 2019, 12:51 AM

Dear Natalia,  
We are still waiting for the reviewers comments  
Best regards  
Son Radu  
Chief Editor

### **3. Bukti Article in Review (12-12-2019)**

mail.google.com/mail/u/0/?tab=rm&ogbl#search/foodresearch.my%40outlook.com/FMfcgwGCH8TpWkTgQFzMTfXvLMxNz

on TV Korea | Sci-Hub | Thrombo... PNP Submit an Article Try Out Uji Kompet... SIMStikes V1.0 Daftar Bahan Maka...

End now Vacation Settings

Gmail foodresearch.my@outlook.com

Compose

Inbox 1,198

Starred

Snoozed

Sent

Drafts 73

More

Labels +

FR-2019-380 Inbox x

Food Research <foodresearch.my@outlook.com> to me Thu, Dec 12, 2019, 9:05 AM

Dear Putriningtyas,

Manuscript FR-2019-380 entitled "Effect of Variation Red Dragon Fruit Peel in Yoghurt on Organoleptic Properties, Protein, Lipid, and Flavonoid Content" which you submitted to Food Research, has been reviewed. The comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s) comments and revise your manuscript. Once the revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to Food Research, your revised manuscript should be submitted by 11<sup>th</sup> January 2020. If it is not possible for you to submit your revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I look forward to receiving your revised manuscript.

Sincerely,  
Professor Dr. Son Radu  
[foodresearch.my@outlook.com](mailto:foodresearch.my@outlook.com)

Chief Editor, Food Research

**From:** Natalia Desy Putriningtyas <[nataliadesy@mail.unnes.ac.id](mailto:nataliadesy@mail.unnes.ac.id)>  
**Sent:** Friday, 10 January, 2020 10:45 PM  
**To:** [foodresearch.my@outlook.com](mailto:foodresearch.my@outlook.com) <[foodresearch.my@outlook.com](mailto:foodresearch.my@outlook.com)>  
**Subject:** Re: FR-2019-380

Dear Professor Dr. Son Radu  
Chief Editor Food Research

Good Evening,

Herewith this email, I attach our manuscript entitled "Comparison of Organoleptic, Protein, Lipid and Flavonoid Content of Commercial Starter and Isolated Culture Red Dragon Fruit Peel Yogurt" by Mardiana, Irwan Budiono and Natalia Desy Putriningtyas. We have made a revision according to the direction from reviewer. We hope this manuscript is accepted in Food Research. If there are still some parts that need to be corrected, please let us know. Do not hesitate to contact us.

Thank you in advance for your attention.

On Tue, Nov 12, 2019 at 9:21 PM Natalia Desy Putriningtyas <[nataliadesy@mail.unnes.ac.id](mailto:nataliadesy@mail.unnes.ac.id)> wrote:

Dear Professor Dr. Son Radu  
Chief Editor Food Research

Thank you for the information provided relating to my manuscript. I hope the good news about the accepted of my manuscript.

On Tue, Nov 12, 2019 at 4:21 PM Natalia Desy Putriningtyas <[nataliadesy@mail.unnes.ac.id](mailto:nataliadesy@mail.unnes.ac.id)> wrote:

Best Regards,  
Natalia Desy Putriningtyas, S.Gz, M.Gizi  
Nutrition, S1 Public Health Science Department  
Faculty of Sport Science  
Universitas Negeri Semarang

**4. Bukti Article accepted (15 Januari 2020)**



15<sup>th</sup> January 2020

Dear Dr. Natalia Desy Putriningtyas,

**ACCEPTANCE LETTER**


Food Research, is pleased to inform you that the following manuscript has been accepted for publication in Food Research journal.

Manuscript Title : Comparison of organoleptic, protein, lipid and flavonoid content of commercial starter and isolated culture red dragon fruit peel yogurt

Authors : Mardiana, Irwan Budiono and Natalia Desy Putriningtyas

We thank you for your fine contribution to the Food Research journal and encourage you to submit other articles to the Journal.

Yours sincerely,



**5. Bukti pemberitahuan pembayaran (APC)  
(17-01-2020)**

mail.google.com/mail/u/0/?tab=rm&ogbl#search/foodresearch.my%40outlook.com/QgrcJHrhxBSPrgvqqfjBxHslpStUGZdxvgG

on TV Korea | Sci-Hub | Thrombo... | Submit an Article | Try Out Uji Kompet... | SIMStikes V1.0 | Daftar Bahan Maka...

End now Vacation Settings

Gmail

Compose

Inbox 1,198

Starred

Snoozed

Sent

Drafts 73

More

Labels +

Search: foodresearch.my@outlook.com

Active

UNNES

11 of 17

Article Processing Charges (APC) FR-2019-380

Natalia Desy Putriningtyas <nataliadesy@mail.unnes.ac.id>  
to Food

Fri, Jan 17, 2020, 6:00 AM


Dear Vivian New  
Editor Food Research

Good Morning,

Herewith this email, I attach our Article Processing Charges (APC) for our manuscript entitled "**Comparison of Organoleptic, Protein, Lipid and Flavonoid Content of Commercial Starter and Isolated Culture Red Dragon Fruit Peel Yogurt**" by Mardiana, Irwan Budiono and Natalia Desy Putriningtyas. We hope our manuscript will be transferred to production. Thank you in advance for your attention.

Best Regards,  
Natalia Desy Putriningtyas, S.Gz, M.Gizi  
Nutrition, S1 Public Health Science  
Sport Science Faculty Universitas Negeri Semarang

One attachment • Scanned by Gmail



mail.google.com/mail/u/0/?tab=rm&ogbl#search/foodresearch.my%40outlook.com/FMfcgXwGCKmVIXsQWjBdp8wbwhfKTsv

on TV Korea | Sci-Hub | Thrombo... | Submit an Article | Try Out Uji Kompet... | SIMStikes V1.0 | Daftar Bahan Maka...

End now Vacation Settings

Gmail

Compose

Inbox 1,198

Starred

Snoozed

Sent

Drafts 73

More

Labels +

Search: foodresearch.my@outlook.com

Active

UNNES

10 of 17

Re: FR-2019-380 - Decision on your manuscript

Food Research <foodresearch.my@outlook.com>  
to me

Dear Dr. Natalia,

It is a pleasure to accept your manuscript for publication in Food Research journal. Please refer to the attachment for your acceptance letter.

Please note that all accepted manuscripts are subjected to Article Processing Charges (APC) as the Journal will provide full publishing services. Please fill in the article procedure form (5) working days. Once we have received the form, your article will be transferred to production.

Thank you for your fine contribution. We look forward to your continued contributions to the Journal.

Sincerely,  
Dr. Vivian New  
Editor  
Food Research

## **6. Bukti article production (11-02-2020)**

**Gmail** foodresearch.my@outlook.com Active UNNES

**Compose**

**Inbox** 1,198

Starred

Snoozed

Sent

**Drafts** 73

More

**Labels** +

Re: FR-2019-380 - Article Production inbox x

**Food Research** <foodresearch.my@outlook.com> to me Tue, Feb 11, 2020, 3:59 PM

Dear Dr Natalia,

Manuscript ID: FR-2019-380  
Manuscript Title: Comparison of organoleptic, protein, lipid and flavonoid content of commercial starter and isolated culture red dragon fruit peel yogurt

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,  
Vivian New  
Editor  
Food Research

**Comparison of organoleptic, protein, lipid and flavonoid content of commercial starter and isolated culture red dragon fruit peel yogurt**

Mardiana, Budiono, I. and Putriningtyas, N.D.

*Nutrition, Department of Public Health, Faculty of Sport Sciences, Universitas Negeri Semarang, Jl. Sekaran Gunungpati, Semarang, Indonesia*

**Article history:**

Received: 12 November 2019

Received in revised form: 10 January 2020

Accepted: 15 January 2020

\*Corresponding author: nataliadesy@mail.unnes.ac.id

Author No.1: <https://orcid.org/0000-0001-8733-8482>;

Author No.1: <https://orcid.org/0000-0001-7605-2462>;

Author No.1: <https://orcid.org/0000-0001-7626-5978>

**Abstract**

Red dragon fruit peel can be processed into yogurt. Yogurt is a coagulated milk product obtained from the lactic acid fermentation — yogurt made from various milk, whether fresh milk, skim milk, or even full cream milk. The additions of red dragon fruit peel may enhance the taste and the therapeutical values. This study was aimed to compare the organoleptic properties, protein, lipid, and flavonoid content of commercial starter and isolated culture red dragon fruit peel yogurt. Red dragon fruit peel yogurt was made with amounts of red dragon fruit peel (25% w/v), full cream milk, sucrose and, commercial starter (*L. bulgaricus*, *L. acidophilus*, *S. thermophilus*, *Bifidobacterium*) or isolated culture (*L. bulgaricus* and *S. thermophilus*). Hedonic scale test form was used for measuring organoleptic properties. Protein content was used Kjeldahl Methods, lipid content was used Soxhlet Methods, and flavonoid content was used chromatography. This research showed that protein and lipid contents in red dragon fruit peel yogurt with commercial starter were higher than red dragon fruit peel yogurt with isolated culture. Flavonoid content in the isolated culture yogurt was higher than commercial starter yogurt. Organoleptic properties between groups were significantly different ( $p < 0.05$ ). Red dragon fruit peel yogurt used isolated culture has a good chance of getting a positive reception from the public and has potential as a functional food.

**Keywords:** Yogurt, Red dragon fruit peel, Organoleptic properties, Flavonoid, *L. bulgaricus*

**7. Bukti pemberitahuan waktu penerbitan  
artikel pada vol.4 issue 3(June) (27-02-2020)**

**Gmail** foodresearch.my@outlook.com

Compose

Inbox 1,198  
Starred  
Snoozed  
Sent  
Drafts 72  
More  
Labels +

Re: FR-2019-380 - Article Published Inbox x

**Food Research** <foodresearch.my@outlook.com>  
to me

Thu, Feb 27, 2020, 9:31 AM

Dear Dr Natalia,

Thank you for the payment.  
Kindly be informed that your manuscript has been assigned to Food Research 2020, Vol. 4, Issue 3 (June). Your manuscript is currently available online and in press on our website <https://www.myfoodresearch.com>. Alternatively, you can download a copy of the manuscript by clicking on the following link:  
[https://doi.org/10.25656/fr.2017.4\(3\).380](https://doi.org/10.25656/fr.2017.4(3).380)

We encourage you to share your published work with your colleagues. Thank you for your fine contribution. We hope that you continue to submit other articles to the Journal.

Thanks & Regards,  
Dr. Vivian New  
Editor  
Food Research



**8. Bukti Article publish (08-06-2023)**

**Comparison of organoleptic, protein, lipid and flavonoid content of commercial starter and isolated culture red dragon fruit peel yogurt**

Mardiana, Budiono, I. and \*Putriningtyas, N.D.

*Nutrition, Department of Public Health, Faculty of Sport Sciences, Universitas Negeri Semarang, Jl. Sekaran Gunungpati, Semarang, Indonesia***Article history:**

Received: 12 November

2019

Received in revised form: 10

January 2020

Accepted: 15 January 2020

Available Online: 22

February 2020

**Keywords:**

Yogurt,

Red dragon fruit peel,

Organoleptic properties,

Flavonoid,

*Lactobacillus bulgaricus***DOI:**[https://doi.org/10.26656/fr.2017.4\(3\).380](https://doi.org/10.26656/fr.2017.4(3).380)**Abstract**

Red dragon fruit peel can be processed into yogurt. Yogurt is a coagulated milk product obtained from the lactic acid fermentation-yogurt made from various milk, whether fresh milk, skim milk, or even full cream milk. The additions of red dragon fruit peel may enhance the taste and the therapeutical values. This study was aimed to compare the organoleptic properties, protein, lipid, and flavonoid content of commercial starter and isolated culture red dragon fruit peel yogurt. Red dragon fruit peel yogurt was made with amounts of red dragon fruit peel (25% w/v), full cream milk, sucrose and, commercial starter (*Lactobacillus bulgaricus*, *Lactobacillus acidophilus*, *Streptococcus thermophilus*, *Bifidobacterium*) or isolated culture (*L. bulgaricus* and *S. thermophilus*). Hedonic scale test form was used for measuring organoleptic properties. Protein content was used Kjeldahl method, lipid content was used Soxhlet method, and flavonoid content was used chromatography. This research showed that protein and lipid contents in red dragon fruit peel yogurt with commercial starter were higher than red dragon fruit peel yogurt with isolated culture. Flavonoid content in the isolated culture yogurt was higher than commercial starter yogurt. Organoleptic properties between groups were significantly different ( $p < 0.05$ ). Red dragon fruit peel yogurt used isolated culture has a good chance of getting a positive reception from the public and has potential as a functional food.

**1. Introduction**

Dragon fruit plants originated from Central America (Jaya, 2010). Dragon fruit comes from tropical climates and is influenced by air humidity, temperature, soil condition, and rainfall. Dragon fruit is a fruit plant that has been cultivated in Indonesia since 2000 (Jafaar *et al.*, 2009; Jaya, 2010). Dragon fruit has good potential in terms of increasing demand with a variety of cultivation techniques that are easy to do. Dragon fruit has high productivity that affects its availability. The high availability of dragon fruit gives a lot of dragon peel waste. Dragon fruit peel has 30-35% of the weight of the fruit and has not been used optimally. Dragon fruit peel has the potential to be developed into functional food (Jafaar *et al.*, 2009).

Red dragon fruit peel contains polyphenol compounds and other active compounds, such as alkaloids, terpenoids, flavonoids, phenolic, carotene, and various other vitamins (niacin, thiamine, pyridoxine, cobalamin) (Wu *et al.*, 2006). Dragon fruit peel is a source of natural antioxidants because of the levels of polyphenols (Wu *et al.*, 2006; Nurliyana *et al.*, 2010).

The antioxidants activity in the red dragon fruit peel has higher than the antioxidant activity in the flesh of the fruit. The research showed that in 1 mg/mL, dragon fruit peels could inhibit 83.84% of free radicals. The red dragon fruit flesh can only inhibit free radicals 27.45%/1 mg. The super red dragon fruit peel has a percentage of DPPH free radical at 79.24% (Wiset *et al.*, 2012).

The red dragon fruit peel used as a food color, and even textiles and extracts were taken as a source of natural antioxidants. Another processed form of red dragon fruit peel was processed into beverages. Yogurt can be an alternative to dragon fruit peel diversification. The red dragon fruit peel used as yogurt was expected to increase the economic value and as an alternative functional food (Untari *et al.*, 2014). Yogurt can have various forms such, as semi-solid and liquid forms (Adolfsson *et al.*, 2004; Kolapo and Olubamiwa, 2012). Yogurt is a probiotic drink that has been proven to have health benefits. The red dragon fruit peel yogurt was made through the addition of lactic acid bacteria (Zainoldin and Baba, 2009; Septiana and Ardriaria, 2016; Wahdaningsih and Untari, 2016). Lactic acid bacteria

\*Corresponding author.

Email: [nataliadasy@mail.unnes.ac.id](mailto:nataliadasy@mail.unnes.ac.id)

that are commonly used are the *Lactobacillus* and *Streptococcus* groups. These function of lactic acid bacteria during the fermentation process by converting lactose to lactic acid in the manufacture of red dragon fruit peel yogurt (Kim et al., 2011; Scavuzzi et al., 2014). Taste, flavor, and texture are the primary keys to made yogurt. Several types of bacteria will affect the taste and flavor of yogurt. The distinctive taste and flavor of yogurt can be produced by mixing the bacteria. One of the objectives of this research to compare red dragon fruit peel yogurt used commercial starters (*L. bulgaricus*, *L. acidophilus*, *S. thermophilus*, *Bifidobacterium*) to isolated culture (*L. bulgaricus* and *S. thermophilus*).

## 2. Materials and methods

This research was conducted in June - August 2019. The formulation for the red dragon peel yogurt was defined in the Nutrition Laboratory of the Universitas Negeri Semarang. Laboratory of Environmental Health, Department of Public Health, Universitas Negeri Semarang as a place for making a red dragon peel yogurt starter. Proximate analysis content measurement was performed in the Chemistry Laboratory, Center for Food and Nutrition Studies, Gajah Mada University and Fisheries Product Quality Testing Laboratory, Maritime Affairs, and Fisheries Semarang. Flavonoid content measurement was performed in the Food Technology Laboratory, UNIKA Soegijapranata Semarang. Isolate *L. bulgaricus* and *S. thermophilus* obtained from Microbiology Laboratory, Center for Food and Nutrition Studies, Gajah Mada University. Ethical clearance has been obtained before the study is carried out with the number 140/KEPK/EC/2019.

### 2.1 Preparation of isolated starter culture

An isolated starter culture which contains *L. bulgaricus* and *S. thermophilus*. Pasteurized full cream milk was heated to 75-85°C for 15 minutes. *L. bulgaricus* and *S. thermophilus*, respectively, were mixed thoroughly with the preheated milk followed by incubation for 18 hrs at 42°C. The yogurt formed was stored at 4°C and used as a starter culture within 14 days.

### 2.2 Preparation of red dragon fruit peel yogurt

Yogurt A was made by a commercial starter. Pasteurized full cream milk was heated to 75-85°C for 15 mins. A mixture of probiotic by Diamond Cold Storage (which contains *L. bulgaricus*, *L. acidophilus*, *S. thermophilus*, *Bifidobacterium*). Yogurt was made by mixing 10 mL commercial starter with 100 mL of preheated milk. Red dragon fruit peel was steamed for 15 mins to remove the skin's distinctive aroma and reduce the content of pectin in the peel's red dragon fruit. Red

dragon fruit peel (25% w/v) was made by adding 25 g of gently mashed fruit into 100 mL of preheated milk. The sucrose for the yogurt was corrected by adding 10 g for every 100 mL preheated milk used. After 7 hrs of incubation at 42°C, the yogurt formed was stored at 4°C.

Yogurt B was made by mixing 5 mL of *L. bulgaricus* starter culture and 5 mL of *S. thermophilus* starter culture with 100 mL of preheated milk. Fresh red dragon fruit peel is separated from the flesh of the fruit and cleaned using running water. Red dragon fruit peel was steamed for 15 mins to remove the skin's distinctive aroma and reduce the content of pectin in the peel's red dragon fruit. Red dragon fruit peel (25% w/v) was made by adding 25 g of gently mashed fruit into 100 mL of preheated milk. The sucrose for the yogurt was corrected by adding 10 g for every 100 mL preheated milk used. After 7 hours of incubation at 42°C, the yogurt formed was stored at 4°C.

### 2.3 Sample preparations

Yogurt sample was mixed with 2.5 mL distilled water then incubated at 45°C for 10 mins.

### 2.4 Flavonoid content

Homogenized yogurt water extract (1 mL) was transferred into a test tube and mixed with 1 mL of 95% ethanol and 5 mL of distilled water. To each sample, 0.5 mL of 50% (v/v) Folin-Ciocalteu reagent was added and mixed. After 5 mins, 1 mL of 5% Na<sub>2</sub>CO<sub>3</sub> was added, and the reaction mixture was allowed to stand for 60 mins. The absorbance was read at 725 nm, and the values were converted to total phenolic, expressed in micrograms.

### 2.5 Organoleptic properties

Organoleptic properties were assessed by a panel of untrained individuals, as much as twenty people, in the Nutrition Laboratory and Chemistry Laboratory Food and Nutrition Centre at Gajah Mada University. They consist of eight females and twelve males. All criteria evaluated were given a score of 1-4 points. Organoleptic tests used a hedonic scale from dislike extremely to like extremely. Criteria included aroma, taste, body texture, color, and visual appearance.

### 2.6 Water content

Water content determination was the gravimetric method with oven drying. This method involved weighing a homogenized yogurt, oven drying 100-105°C for 3 hrs, reweighing, and calculating the mass of water lost as a percentage of the mass of the dried sample. Results are usually reported as % sample water on a dry-



mass basis.

### 2.7 Ash content

Ash refers to the inorganic residue remaining after either ignition or complete oxidation of organic matter in the food sample. Determining the ash content of a food is part of proximate analysis for nutritional evaluation, and it is an important quality attribute for some food ingredients.

The gravimetric method was made for the determination of ash content on red dragon fruit peel yogurt. The sample is ashed at 600 - 650°C to constant weight and, the ash is determined by weighing. This method used the dry ashing technique with a muffle furnace to determine the ash content of the sample.

### 2.8 Protein content

Protein content in red dragon fruit peel yogurt was assessed by the Kjeldahl method. In the Kjeldahl procedure, after digestion in concentrated sulfuric acid, the total organic nitrogen is converted to ammonium sulfate. Ammonia is formed and distilled into a boric acid solution under alkaline conditions. The borate anions formed are titrated with standardized hydrochloric acid, which is calculated as the content of nitrogen representing the amount of crude protein in the sample.

### 2.9 Fat content

The soxhlet method described for crude fat analysis on red dragon fruit peel yogurt. The organoleptic properties were analyzed using the Mann-Whitney test and p-value of less than 0.05 were considered statistically significant.

## 3. Results and discussion

Table 1 shows the organoleptic properties of yogurt from two formulations. Panelists preferred group B

compared to group A. The intended organoleptic properties include aroma, taste, body texture, color, and visual appearance. The bright pink color is more visible in group B with a thick texture typical of yogurt. The aroma and taste produced by yogurt in group B are not very flavorful and sour. In this research, the red dragon fruit peel was steamed 15 mins. Chan and Choo (2013) found that low temperatures might insufficient for the hydrolysis of protopectin (the insoluble form of pectin) by acids, which leads to a lower yield of pectin. Pectin consists essentially of galacturonic acid residues linked by bonds  $\alpha$  (1-4) partially acetylated or esterified by methyl groups, which confer gelling, thickening, stabilizing properties, and has a great water holding (Fishman *et al.*, 2000; Arioui *et al.*, 2017).

Table 2 shows the differences between group A and group B. These differences include aroma, taste, body texture, color, and visual appearance flavor. Research conducted by Settachaimongkon *et al.* (2014) reported that proto-cooperation between *S. thermophilus* is one of the important factors that play a role in the fermentation process and the quality of yogurt. The association between the two yogurt lactic acid bacteria (LAB) *S. thermophilus* and *L. bulgaricus* is regarded as a proto-cooperation because it is beneficial for both species, but each bacterium can grow alone in milk (Tamime and Robinson, 1999; Rul, 2017) This proto-cooperation has industrial importance because it can improve yogurt's properties, such as the texture via exopolysaccharide production and the flavor via the production of aromatic compounds (Rul, 2017). Yogurt is a form of milk processing product that utilizes microbes in the fermentation process of fresh milk into a form of semi-solid emulsion products with a more acidic taste. Lactic acid bacteria have the potential to be anticholesterol due to exopolysaccharide (EPS) (Horiuchi and Sasaki, 2012). Hydrolysis of lactose to lactic acid by microbial activity increases milk acidity resulting in the coagulation of milk protein and yogurt having an acidic taste.

Table 1. Frequency distribution of organoleptic properties red dragon peel yogurt

Groups	Frequency Distribution of Organoleptic Properties				
	Aroma	Taste	Body Texture	Color	Overall
<b>A</b>					
Dislike extremely	4 (20%)	0 (0%)	1 (5%)	0 (0%)	0 (0%)
Dislike	13 (65%)	9 (45%)	13 (65%)	7 (35%)	12 (60%)
Like	3 (15%)	11 (55%)	5 (25%)	10 (50%)	8 (40%)
Like extremely	0 (0%)	0 (0%)	1 (5%)	3 (15%)	0 (0%)
<b>B</b>					
Dislike extremely	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (5%)
Dislike	1 (5%)	3 (15%)	3 (15%)	2 (10%)	1 (5%)
Like	10 (50%)	9 (45%)	9 (45%)	11 (55%)	10 (50%)
Like extremely	9 (45%)	8 (40%)	8 (40%)	7 (35%)	8 (40%)

A: red dragon peel steamed, full cream milk, sucrose, commercial starter; B: red dragon peel steamed, full cream milk, sucrose, isolated culture

Table 2. Difference mean

Groups	Mean Rank				
	Aroma	Taste	Body Texture	Color	Overall
A	11.73	15.3	14.15	17.08	14.2
B	29.28	25.7	26.85	23.93	26.8
<i>p</i>	0.001*	0.002*	0.001*	0.042*	0.001*

A: red dragon peel steamed, full cream milk, sucrose, commercial starter; B: red dragon peel steamed, full cream milk, sucrose, isolated culture, *p*\* significant, mann-whitney test

The texture of yogurt is strongly influenced by various solids, which are the result of the fermentation process (Widiyaningsih, 2011; Scavuzzi et al., 2014). Red dragon fruit peel yogurt in this study used *L. bulgaricus* and *S. thermophilus* due to a symbiotic relationship between the two lactic acid bacteria and can reduce fermentation time. The growth and stimulation of *L. bulgaricus* can be stimulated by the production of formic acid, pyruvic acid, and carbon dioxide produced by *S. thermophilus*. The growth of *S. thermophilus* is stimulated by free amino acids and peptides derived from milk production by *L. bulgaricus* (Settachaimongkon et al., 2014). This is said to be the symbiotic fermentation relationship. The second growth of lactic acid bacteria is homofermentative so that the acid can be produced faster. The homofermentative bacteria, *L. bulgaricus* and *S. thermophilus*, which produce lactic acid starting from lactose milk (Arioui et al., 2017). The fermentation process results in increased microbial activity, a decrease in pH, and an increase in acid levels in the fermentation product. The fermentation process can improve organoleptic quality and nutritional value, including improving the quality of protein and its digestive value (Horiuchi and Sasaki, 2012; Untari et al., 2014; Settachaimongkon et al., 2014).

Yogurt is viewed as a highly nutritious dairy food product due to levels of protein, lipid, and flavonoid (Widiyaningsih, 2011; Bintari et al., 2017). Table 3 shows the analysis component yogurt A and yogurt B. The content of ash, lipid and protein in yogurt A were greater than yogurt B. The ash, lipid and protein content in yogurt A, respectively 0.95%; 4.37%; 2.85%. Yogurt has highly attractive nutritional properties- it is low in calories but contains enough macro and micronutrients (proteins, fatty acids, calcium, phosphorus, and vitamins) to cover a person's daily needs (Rul, 2017). Yogurt's lactic acid bacterial cultures can also improve digestive conditions, including gut microflora growth, bowel transit time, and immune response in the alimentary tract (Bintari et al., 2017; Putriningtyas and Wahyuningsih, 2017). Flavonoid content showed an increased level in group B. Table 3 shows the flavonoid content in yogurt

B has 16.6% and yogurt A has 15%.

Flavonoid is bioactive phenols commonly found in fruits, vegetables, and parts of plants. Phenolic and polyphenolic compounds constitute the main class of natural antioxidants present in plants (Wu et al., 2006). Fermentation increases antioxidant activity and, thus, the functional value of the foodstuff. Table 3 shows that group B has higher flavonoid content than group A. Polyphenolic flavonoids display strong antioxidant activity. The flavonoid and lactic acid bacteria content in yogurt gives it potential as a functional food product. Flavonoids have antioxidant activity that can help protect against free radicals and prevent clinical complications from metabolic diseases, such as diabetes mellitus, dyslipidemia, and metabolic syndrome (Galleano et al., 2012; Babu et al., 2013; Jakobsdottir et al., 2014). The addition of dragon fruit increased the total phenolics content in yogurt compared to plain yogurt based on previous research (Zainoldin and Baba, 2009). This might be due to the addition of dragon fruit that contains high phenolic compounds such as phenolic acids and polyphenols, which are commonly known to be found in plants.

Table 3. Analysis of components in yogurt

Groups	Nutrition				
	Water (%)	Ash (%)	Lipid (%)	Protein (%)	Flavonoid (%)
A	79.81	0.95	4.37	2.85	15
B	80.65	0.63	2.11	2	16.6

A: red dragon peel steamed, full cream milk, sucrose, commercial starter; B: red dragon peel steamed, full cream milk, sucrose, isolated culture

This study shows that yogurt can be used to create a nutritious and healthy beverage. The red dragon fruit peel yogurt is suitable for consumption not only by healthy people but also by those who have metabolic diseases, such as dyslipidemia and diabetes as it contains flavonoid, fiber, and lactic acid bacteria that could help decrease blood sugar, cholesterol and triglyceride levels (Babu et al., 2013). This study has many limitations that could be improved by further research, especially regarding the safety and recommended dosages for certain health conditions.

#### 4. Conclusion

Red dragon fruit peel yogurt used isolated culture has a good chance of getting a positive reception from the public. Red dragon fruit peel yogurt used isolated culture has potential as a functional food because of its high flavonoid content and low lipid content.



**Conflict of Interest**

The authors declare no conflict of interest.

**Acknowledgments**

This study received grant funding from Lembaga Penelitian dan Pengabdian (LP2M) Universitas Negeri Semarang budget in 2019.

**References**

- Adolfsson, O., Meydani, S.N. and Rusell, R.M. (2004). Yogurt and gut function. *American Journal of Clinical Nutrition*, 80(2), 245-256. <https://doi.org/10.1093/ajcn/80.2.245>
- Arioui F., Saada, D.A. and Cheriguene, A. (2017). Physicochemical and sensory quality of yogurt incorporated with pectin from peel of Citrus sinensis. *Food Science and Nutrition*, 5(2), 358-364. <https://doi.org/10.1002/fsn3.400>
- Babu, P.V.A., Liu, D. and Gilbert, E.R. (2013). Recent advances in understanding the anti-diabetic actions of dietary flavonoids. *Journal of Nutritional Biochemistry*, 24(11), 1777-1789. <https://doi.org/10.1016/j.jnutbio.2013.06.003>
- Bintari, S.H., Widyastiti, N.S., Putriningtyas, N.D., Hapsari, R. and Nugraheni, K. (2017). Development and properties of tegurt, a yogurt-like tempe product. *Pakistan Journal of Nutrition*, 16(4), 221-226. <https://scialert.net/abstract/?doi=pjn.2017.221.226>
- Chan, S.Y. and Choo, W.S. (2013). Effect of extraction conditions on the yield and chemical properties of pectin from cocoa husks. *Food Chemistry*, 141(4), 3752-3758. <https://doi.org/10.1016/j.foodchem.2013.06.097>
- Fishman, M.L., Chau, H.K., Hoagland, P. and Ayyad, K. (2000). Characterization of pectin, flash-extracted from orange albedo by microwave heating, under pressure. *Carbohydrate Research*, 323(1-4):126-138. [https://doi.org/10.1016/s0008-6215\(99\)00244-x](https://doi.org/10.1016/s0008-6215(99)00244-x)
- Galleano, M., Calabro, V., Prince, P.D., Litterio, M.C., Piotrkowski, B., Vazquez-Prieto, M.A., Miatello, R.M., Oteiza, P.I. and Fraga, C.G. (2012). Flavonoids and metabolic syndrome. *Annals of the New York Academy of Sciences*, 1259, 87-94. <https://doi.org/10.1111/j.1749-6632.2012.06511.x>
- Horiuchi, H. and Sasaki, Y. (2012). Short communication: Effect of oxygen on symbiosis between *Lactobacillus bulgaricus* and *Streptococcus thermophilus*. *Journal of Dairy Science*, 95(6), 2904-2909. <https://doi.org/10.3168/jds.2011-5147>
- Jafaar, A.R., Abdul Rahman, A.R., Mahmod, N.Z.C. and Vasudevan, R. (2009). Proximate analysis of dragon fruit (*Hylocereus polyrhizus*). *American Journal of Applied Sciences*, 6(7), 1341-1346. <https://doi.org/10.3844/ajassp.2009.1341.1346>
- Jakobsdottir, G., Nyman, M. and Fak, F. (2014). Designing future prebiotic fiber to target metabolic syndrome. *Nutrition*, 30(5), 497-502. <https://doi.org/10.1016/j.nut.2013.08.013>
- Jaya, I.K.D. (2010). Morfologi dan fisiologi buah naga dan prospek masa depannya di Indonesia. *Crop Argo*, 3(1), 44-50. <https://cropagro.unram.ac.id/index.php/caj/article/view/60/47>
- Kim, H., Choi, H.K., Moon, J.Y., Kim, Y.S., Mosaddik, A. and Cho, S.K. (2011). Comparative antioxidant and antiproliferative activities of red and white pitayas and their correlation with flavonoid and polyphenol content. *Journal of Food Science*, 76(1), C38-45. <https://doi.org/10.1111/j.1750-3841.2010.01908.x>
- Kolapo, A.L. and Olubamiwa, A.O. (2012). Effect of different concentration of coconut milk on the chemical and sensory properties of soy-coconut milk based yogurt. *Food and Public Health*, 2(4), 85-91. <https://doi.org/10.5923/j.fph.20120204.01>
- Nurliyana, R., Syed Zahir, I., Mustapha Suleiman, K., Aisyah, M.R. and Kamarul Rahim, K. (2010). Antioxidant study of pulps and peels of dragon fruit: A comparative study. *International Food Research Journal*, 17(2), 367-375. [http://www.ifrj.upm.edu.my/17%20\(02\)%202010/IFRJ-2010-367-375%20Kamarul%20IU\[1\].pdf](http://www.ifrj.upm.edu.my/17%20(02)%202010/IFRJ-2010-367-375%20Kamarul%20IU[1].pdf)
- Putriningtyas, N.D. and Wahyuningsih, S. (2017). Potensi yogurt kacang merah (*Phaseolus vulgaris* L) ditinjau dari sifat organoleptik, kandungan protein, lemak dan flavonoid. *The Indonesian Journal of Nutrition*, 6(1), 37-43. <https://doi.org/10.14710/jgi.6.1.37-43>
- Rul, F. (2017). Yogurt: microbiology, organoleptic properties and probiotic potential. In Ray, R.C. and Montet, D. (Eds.) *Fermented Foods, Part II: Technological Interventions*, p. 525. United States: CRC Press.
- Scavuzzi, B.M., Henrique, F.C., Miglioranza, L.H.S., Simao, A.N.C. and Dichi, I. (2014). Impact of prebiotics, probiotics and synbiotics on components of the metabolic syndrome. *Annals of Nutritional Disorders and Therapy*, 1(2), 1009. <https://www.austinpublishinggroup.com/nutritional-disorders/fulltext/and-t-v1-id1009.php>
- Septiana, W.C. and Ardiaria, M. (2016). Efek pemberian seduhan kulit buah naga merah (*Hylocereus polyrhizus*) terhadap kadar malondialdehyde (MDA) tikus sprague dawley dislipidemia. *Journal of*

- Nutrition College*, 5(4), 344-352. <https://doi.org/10.14710/jnc.v5i4.16434>. [In Bahasa Indonesia].
- Settachaimongkon, S., Nout, M.J., Antunes Fernandes, E.C., Hettinga, K.A., Vervoort, J.M., van Hooijdonk, T.C.M., Zwietering, M.H., Smid, E.J. and van Valenberg, H.J.F. (2014). Influence of different proteolytic strains of *Streptococcus thermophilus* in co-culture with *Lactobacillus delbrueckii* subsp. *bulgaricus* on the metabolite profile of set-yogurt. *International Journal of Food Microbiology*, 177, 29-36. <https://doi.org/10.1016/j.ijfoodmicro.2014.02.008>
- Tamime, A.Y. and Robinson, R.K. (1999). *Tamime and Robinson's Yoghurt: Science and technology*. 2<sup>nd</sup> ed. Boca Raton, Florida, USA: CRC Press.
- Untari, E.K., Wahdaningsih, S. and Damayanti, A. (2014). Efek fraksi n-heksana kulit *Hylocereus polyrhizus* terhadap aktivitas katalase tikus stress oksidatif. *Pharmaceutical Sciences and Research*, 1 (3), 141-153. <https://doi.org/10.7454/psr.v1i3.3489>. [In Bahasa Indonesia].
- Wahdaningsih, S. and Untari, E.K (2016). Pengaruh pemberian fraksi metanol kulit buah naga merah (*Hylocereus polyrhizus*) terhadap kadar malondialdehid pada tikus (*Rattus novergicus*) wistar yang mengalami stress oksidatif. *Jurnal Pharmascience*, 3(1), 45-55. <https://doi.org/10.20527/jps.v3i1.5834>. [In Bahasa Indonesia].
- Widiyaningsih, E.N. (2011). Peran probiotik untuk kesehatan. *Jurnal Kesehatan*, 4(1), 14-20. [In Bahasa Indonesia].
- Wiset, N., Poomsa-ad, N. and Srilaong, V. (2012). Comparisons of antioxidant activity and bioactive compounds of dragon fruit peel from various drying methods. *World Academy of Science, Engineering and Technology*, 6, 10-29. <https://doi.org/10.5281/zenodo.1062480>
- Wu, L., Hsu, H., Chen, Y., Chiu, C., Lin, Y. and Ho, J. (2006). Antioxidant and antiproliferative activities of red pitaya. *Food Chemistry*, 95(2), 319-327. <https://doi.org/10.1016/j.foodchem.2005.01.002>
- Zainoldin, K. and Baba, A.S. (2009). The effect of *Hylocereus polyrhizus* and *Hylocereus undatus* on physicochemical, proteolysis and antioxidant activity in yogurt. *International Journal of Nutrition and Food Engineering*, 3(12), 585-590. <https://www.researchgate.net/publication/235697248>