

The dynamics of the basic model for the zika epidemic

by Adi Cahyono

Submission date: 14-Nov-2022 10:58AM (UTC+0700)

Submission ID: 1953127617

File name: 27_The_dynamics_of_the_basic_model_for_the_zika.pdf (547.04K)

Word count: 2899

Character count: 11857

PAPER • OPEN ACCESS

The dynamics of the basic model for the zika epidemic

To cite this article: M Kharis and A N Cahyono 2020 *J. Phys.: Conf. Ser.* **1567** 022085

View the [article online](#) for updates and enhancements.

You may also like

- [Graphene oxide-based molecular diagnostic biosensor for simultaneous detection of Zika and dengue viruses](#)
Ji-Seon Lee, Jung-ho Kim, Hojeong Shin et al.
- [Construction of an Impedimetric Biosensor to Identify Zika VIRUS through a Platform Modified with Poly 4-ABA](#)
Filipe Soares da Cruz, Tais Aparecida Reis, Ricardo Augusto Corrêa et al.
- [Qualitative and Quantitative study of Zika virus epidemic model under Caputo's fractional differential operator](#)
Faiz Muhammad Khan, Amjad Ali, Zia Ullah Khan et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Extended abstract submission deadline: April 22, 2022

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD



Submit your abstract



The dynamics of the basic model for the zika epidemic

M Kharis and A N Cahyono

Department of Mathematics, Universitas Negeri Semarang, Indonesia

*Corresponding author: kharis.mat@mail.unnes.ac.id

Abstract. Zika disease began to get the world spotlight when in 2016 around 1-1.5 million people were infected with the Zika virus in Brazil and 4000 babies were born Microcephaly. Indonesia is a region that is potentially attacked by Zika virus outbreaks. This study discusses the mathematical model for the spread of Zika disease in humans is associated with the population of spreading vectors which in this case are mosquitoes. The model used is a deterministic model in the form of a compartment model in sub-population classes. The purpose of this research is to build a mathematical model, analyze the stability point, and arrange a simulation of a mathematical model.

1. Introduction

Zika disease is an infectious disease caused by the Zika virus which is derived from a type of flavivirus that is similar to the dengue virus. This virus is transmitted through the bite of the *Aedes* sp. Zika virus (ZIKV) is originally discovered in a captive monkey living in the Zika Forest of Uganda, Africa, in 1947 [1]. Symptoms arising from this viral infection include fever, freckled skin, headaches, joint pain, muscle aches, and inflammation of the conjunctiva. Symptoms of this disease appear for 2 to 7 days [2]. The Zika virus became a worldwide concern after Brazilian health authorities discovered a link between pregnant women infected with the Zika virus and the birth of a microcephaly baby. Microcephaly is a condition in which babies have small heads and incomplete brain development [3]. In addition, the Zika virus is indicated to cause Guillain-Barre syndrome which is an acute inflammation that can cause nerve cell damage.

The spread of the Zika virus was first identified in Uganda [4]. The Zika virus was reported to have spread in Yap, Micronesia and French Polynesia in 2007 [5-6]. In October 2013, the Zika virus spread in Polynesia, France [7]. Early in 2015, the Zika virus was reportedly detected in Brazilian territory [8]. In 2016 the Zika virus has also spread in Singapore.

In Indonesia, the Eijkman Molecular Biology Institute has reported the presence of the Zika virus to the Ministry of Health. The agency noted that there were five cases of the Zika virus in Indonesia, namely: (1) in 1981 there was reported one patient at Tegalyoso Hospital, Klaten, Central Java; (2) in 1983 there were reported six out of 71 samples in Lombok, NTB; (3) in 2013 it was reported that a female tourist from Australia tested positive for the Zika virus after nine days living in Jakarta; (4) in 2015 it was reported that a tourist from Australia was infected with the Zika virus after being bitten by a monkey in Bali; and (5) in 2015-2016 a patient in Jambi Province tested positive for the Zika virus. The spread of the Zika virus in Indonesia is still relatively low, but the potential spread of the Zika disease needs to be watched out and needs to be anticipated. At present, a vaccine to cure the Zika virus has not been found. Treatment that can be done is still supportive such as adequate rest, consuming enough water to prevent dehydration, and taking fever or pain relievers [2].



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

2. Methods

The research started by doing literacy study. In this step, the fact and some assumptions from various scientific literacy are collected. After that, some assumptions added to complete the facts. The next step was building and analyzing the mathematics model. In this step, the mathematics model was constructed and analyzed to determine the equilibrium points and their stability. The next step was making simulations with parameters value got from another paper. In this paper, we didn't give simulations.

3. Result and Discussion

In this paper, we used recruitment-death to the Dynamic of human and mosquito population. Transfer diagram of Zika epidemic was given in Figure 1.




Figure 1. Transfer diagram of Zika epidemic

Where N, S, I, R respectively are the human population, the number of susceptible human, the number of infected human, and the number of recovered human. For mosquito, N_v, S_v, I_v respectively are the mosquito population, the number of susceptible mosquito, and the number of infected mosquito.

The meaning of parameters: B means recruitment rate of human population, μ means the natural death rate in human, β means the probability of virus transfer from infected mosquito to susceptible human, c means the recovery rate of infected human, q means the recovery rate of treated human, and m means the death rate caused of infected. μ_v means the birth rate which is assumed same with the natural death rate in mosquito, β_v means the probability of virus transfer from infected human to susceptible mosquito. From Fig. 1, we construct the System (1).

$$\begin{aligned}
 \frac{dS_v}{dt} &= \mu_v N_v - \mu_v S_v - \beta_v \frac{I}{N} S_v \\
 \frac{dI_v}{dt} &= \beta_v \frac{I}{N} S_v - \mu_v I_v \\
 S_v + I_v &= N_v \\
 \frac{dS}{dt} &= B - \mu S - \beta \frac{S}{N} I_v \\
 \frac{dI}{dt} &= \beta \frac{S}{N} I_v - (\mu + m + c) I \\
 \frac{dR}{dt} &= cl - \mu R \\
 S + I + T + R &= N
 \end{aligned}
 \tag{1}$$

From System (1), we get $\frac{dN_v}{dt} = 0 \Leftrightarrow N_v = K, K \in \mathbb{R}$ and $\frac{dN}{dt} = \frac{dS}{dt} + \frac{dI}{dt} + \frac{dR}{dt} = B - \mu N - mI$.

Hence, we get System (2).

$$\begin{aligned} \frac{dI_v}{dt} &= \beta_v \frac{I}{N} (K - I_v) - \mu_v I_v \\ \frac{dN}{dt} &= B - \mu N - mI \\ \frac{dI}{dt} &= \beta \frac{N-I-R}{N} I_v - (\mu + m + c)I \\ \frac{dR}{dt} &= cI - \mu R \end{aligned} \quad (2)$$

Clear that $\frac{dN}{dt} = 0 \Leftrightarrow N = \frac{B-mI}{\mu} \leq \frac{B}{\mu}$.

Hence, we can determine the domain of System (2) on region

$D = \{(I_v, N, I, R) \in R_4^+ | 0 \leq I_v \leq K, 0 \leq I + T \leq N \leq \frac{B}{\mu}, N > 0\}$ where R_4^+ is non negative region subset R_4 .

Lemma 1

- 1) If $\frac{\beta K}{B} > 1$ then $R_0 = \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu+m+c}$.
- 2) If $\frac{\beta K}{B} < 1$ then $R_0 = \frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu+m+c}$.

Lemma 1 will be used in determining the existence of equilibrium points such that in theorem 2.

Theorem 2

Let R_0 in Lemma 1 and $P_0 = (I_v, N, I, R) = (0, \frac{B}{\mu}, 0, 0)$ dan $P_1 = (\hat{I}_v, \hat{N}, \hat{I}, \hat{R})$ dengan

$$\hat{I}_v = \frac{\beta_v K I}{\frac{\mu_v(B-mI)}{\mu} + \beta_v I}, \hat{N} = \frac{B-mI}{\mu}, \hat{R} = \frac{cI}{\mu}, \text{ and } \hat{I} = \frac{-a_1 - \sqrt{a_1^2 - 4a_0 a_2}}{2a_0}$$

where $a_0 = \mu_v^2 m^2 (\mu + m + c) \left(\frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} - 1 \right)$,

$a_1 = B\mu_v^2 m (\mu + m + c) \left(1 - \frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu+m+c} \right) + B\mu_v^2 m (\mu + m + c) \left(1 - \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} \right) - \beta \cdot \beta_v \cdot K \mu_v \cdot \mu (\mu + c)$, and

$a_2 = B^2 \mu_v^2 (\mu + m + c) \left(\frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu+m+c} - 1 \right)$.

- 1) If $R_0 < 1$ then System (2) only have one equilibrium point P_0 .
- 2) If $R_0 > 1$ then System (2) has two equilibrium points i.e. P_0 and P_1 .

Proof:

The equilibrium points were determined by making $\frac{dI_v}{dt}$, $\frac{dN}{dt}$, $\frac{dI}{dt}$, and $\frac{dR}{dt}$ in System (2) equal to zero. We got the System (3)

$$\begin{aligned} \beta_v \frac{I}{N} (K - I_v) - \mu_v I_v &= 0 \\ B - \mu N - mI &= 0 \\ \beta \frac{N-I-R}{N} I_v - (\mu + m + c)I &= 0 \\ cI - \mu R &= 0 \end{aligned} \quad (3)$$

From the second and fourth equations, we got $N = \frac{B-mI}{\mu}$ dan $R = \frac{cI}{\mu}$. Both were substituted to the first equation then we got

$$I_v = \frac{\beta_v \cdot K \cdot I}{\mu_v \cdot N + \beta_v \cdot I} = \frac{\beta_v \cdot K \cdot I}{\mu_v(B - m \cdot I) + \beta_v \cdot I} = \frac{\beta_v \cdot \mu \cdot K \cdot I}{\mu_v(B - m \cdot I) + \beta_v \cdot \mu \cdot I}$$

Substitute all value to the third equation, then we got

$$\frac{I}{\mu_v(B-mI)[\mu_v B + \mu \cdot I(\beta_v - \mu_v)]} [a_0 \cdot I^2 + a_1 \cdot I + a_2] = 0 \quad (4)$$

where

$$a_0 = \mu_v^2 m^2 (\mu + m + c) \left(\frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} - 1 \right),$$

$$a_1 = B \mu_v^2 m (\mu + m + c) \left(1 - \frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} \right) + B \mu_v^2 m (\mu + m + c) \left(1 - \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} \right) - \beta \cdot \beta_v \cdot K \cdot \mu_v \cdot \mu (\mu + c)$$

$$a_2 = B^2 \mu_v^2 (\mu + m + c) \left(\frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} - 1 \right)$$

The equation (4) gave solutions $I = 0$ and $I \neq 0$.

If $I = 0$ then we get free disease equilibrium point $P_0 = (I_v, N, I, R) = \left(0, \frac{B}{\mu}, 0, 0 \right)$.

For $I \neq 0$, we will find $I > 0$.

Clear that if $N > 0$ and $I \neq \frac{\mu_v B}{\mu_v m - \beta_v \mu}$, then $\mu_v(B - mI)[\mu_v B + \mu \cdot I(\beta_v - \mu_v)] \neq 0$. Hence, to determine I in equation (4), we can solve only the equation $a_0 \cdot I^2 + a_1 \cdot I + a_2 = 0$.

By Lemma 1, we got that if $R_0 > 1$ then $a_0 > 0$, $a_1 < 0$, dan $a_2 > 0$.

Hence, $a_0 \cdot I^2 + a_1 \cdot I + a_2 = 0$ has two solutions i.e.

$$I_1 = \frac{-a_1 - \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}}{2 \cdot a_0} \text{ and } I_2 = \frac{-a_1 + \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}}{2 \cdot a_0}$$

Clear that $a_1^2 - 4 \cdot a_0 \cdot a_2$

$$= B^2 \mu_v^4 m^2 (\mu + m + c)^2 \left(\frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} - \frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} \right)^2 + [\beta \cdot \beta_v \cdot K \cdot \mu_v \cdot \mu (\mu + c)]^2 + 2 \cdot \beta \cdot \beta_v \cdot K \cdot \mu_v \cdot \mu (\mu + c) \cdot B \mu_v^2 m (\mu + m + c) \left[\left(\frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} - 1 \right) + \left(\frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} - 1 \right) \right]$$

Hence $a_1^2 - 4 \cdot a_0 \cdot a_2 > 0$ if $R_0 > 1$ where R_0 in Lemma 1.

Hence $I_1 = \frac{-a_1 - \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}}{2 \cdot a_0} > 0$ and $I_2 = \frac{-a_1 + \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}}{2 \cdot a_0} > 0$.

We will show that $I_1 < \frac{B}{m}$.

$$\text{Clear that } \frac{B}{m} - I_1 = \frac{B}{m} - \frac{-a_1 - \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}}{2 \cdot a_0} = \frac{2 \cdot B \cdot a_0 + m \cdot a_1 + m \cdot \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}}{2 \cdot m \cdot a_0}$$

To determine the sign of $\frac{B}{m} - I_1$, we exactly can check the sign of $2 \cdot B \cdot a_0 + m \cdot a_1 + m \cdot \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2}$.

$$\text{Clear that } 2 \cdot B \cdot a_0 + m \cdot a_1 + m \cdot \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2} = -B \cdot \mu_v^2 m^2 (\mu + m + c) \left(\frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} - \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} \right) - \beta \cdot \beta_v \cdot K \cdot \mu_v \cdot \mu \cdot m (\mu + c) + \left[B^2 \cdot \mu_v^4 m^4 (\mu + m + c)^2 \left(\frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} - \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} \right)^2 + \beta^2 \cdot \beta_v^2 \cdot K^2 \cdot \mu_v^2 \cdot \mu^2 \cdot m^2 (\mu + c)^2 \right]$$

$$+2 \cdot \beta \cdot \beta_v \cdot K \cdot \mu_v \cdot \mu(\mu + c) \cdot B \mu_v^2 m^3 (\mu + m + c) \left[\left(\frac{\beta K}{B} \cdot \frac{\beta_v}{\mu_v} \cdot \frac{\mu}{\mu + m + c} - 1 \right) + \left(\frac{\beta_v}{\mu_v} \cdot \frac{\mu}{m} - 1 \right) \right]^{\frac{1}{2}}$$

From the calculating process above we get $2 \cdot B \cdot a_0 + m \cdot a_1 + m \cdot \sqrt{a_1^2 - 4 \cdot a_0 \cdot a_2} > 0$

So $\frac{B}{m} - I_1 > 0$ or $I_1 < \frac{B}{m}$. So $\frac{B}{m} - I_1 > 0 \Leftrightarrow B - mI_1 > 0 \Leftrightarrow \frac{B - mI_1}{\mu} > 0 \Leftrightarrow N > 0$.

Hence $I_1 \in D$.

With a similar process, we can show $I_2 > \frac{B}{m}$. We get $B - mI_2 < 0 \Leftrightarrow \frac{B - mI_2}{\mu} < 0 \Leftrightarrow N < 0$.

Hence $I_2 \notin D$.

The stability theorem of the free disease equilibrium point was given in Theorem 3.

Theorem 3

Let R_0 in Lemma 1 and $P_0 = (I_v, N, I, R) = \left(0, \frac{B}{\mu}, 0, 0\right)$

- 1) If $\frac{\beta K}{B} < 1$ and $R_0 < 1$ the P_0 is locally asymptotical stable.
- 2) if $R_0 > 1$ then P_0 is unstable.

The stability theorem of the endemic equilibrium point was given in Theorem 4.

Theorem 4

Let R_0 in Lemma 1 and $P_1 = (I_v, N, I, R)$ was defined in Theorem 2.

if $\frac{B\mu_v}{\mu(\mu_v + \mu + c) + \mu_v(\mu + m + c)} < 1$, $[(\mu + m + c) + (\mu + \mu_v)][(\mu + m + c)(\mu + \mu_v) + \mu\mu_v] \geq 1$, and

$R_0 > 1$ then P_1 is locally asymptotically stable.

Proof:

The characteristic equation of jacobian matrices of P_1 is

$$\frac{(x + \mu_v)(x + \mu)}{N_2^3 (\mu_v N_2 + \beta_v \hat{I} \mu)} (c_0 x^3 + c_1 x^2 + c_2 x + c_3) = 0 \quad (5)$$

Where

$$N_2 = B - m\hat{I}$$

$$c_0 = c_3 = N_2^3 \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu)$$

$$c_1 = N_2^3 \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) [(\mu + m + c) + (\mu + \mu_v)] + \beta_v \hat{I} \mu \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) + \beta \beta_v K \mu_v \hat{I} \mu^2 N_2^2$$

$$c_2 = N_2^3 \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) [(\mu + m + c)(\mu + \mu_v) + \mu\mu_v] + N_2^2 \mu_v \beta_v \hat{I} \mu (\mu_v N_2 + \beta_v \hat{I} \mu) (2\mu + m + c) + N_2 \beta \mu^2 \beta_v K \mu_v \hat{I}^2 [m(\mu + c) + \beta_v \mu] + N_2^2 \beta \mu \beta_v K \mu_v [\hat{I} (\mu + \mu_v)(\mu + c) + \hat{I} \mu \mu_v - N_2 \mu_v]$$

From equation (5) we got two eigen values i.e. $x_1 = -\mu_v < 0$ and $x_2 = -\mu < 0$.

We will show that $c_0 x^3 + c_1 x^2 + c_2 x + c_3 = 0$ has roots with negative real part by referring

Routh Hurwitz criteria i.e.

$$(1) c_0 > 0, c_1 > 0, c_2 > 0, \text{ and } c_3 > 0$$

$$(2) c_1 c_2 - c_0 c_3 > 0$$

Clear that $c_0 = c_3 > 0$ dan $c_1 > 0$.

Clear that $c_2 > 0$ if and only if $\hat{I} (\mu + \mu_v)(\mu + c) + \hat{I} \mu \mu_v - N_2 \mu_v > 0$.

Clear that $\hat{I} (\mu + \mu_v)(\mu + c) + \hat{I} \mu \mu_v - (B - m\hat{I})\mu_v > 0$

$$\Leftrightarrow \hat{I} > \frac{B\mu_v}{\mu(\mu_v + \mu + c) + \mu_v(\mu + m + c)}$$

So $c_2 > 0$ if and only if $\hat{I} > \frac{B\mu_v}{\mu(\mu_v+\mu+c)+\mu_v(\mu+m+c)}$.

Because \hat{I} means the number of infected humans then $\hat{I} > 1$ Hence $c_2 > 0$ if and only if $\frac{B\mu_v}{\mu(\mu_v+\mu+c)+\mu_v(\mu+m+c)} < 1$.

Clear that $c_1c_2 - c_0c_3 =$

$$= (N_2^3 \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu))^2 \left\{ \left[[(\mu + m + c) + (\mu + \mu_v)] [(\mu + m + c)(\mu + \mu_v) + \mu \mu_v] - 1 \right] \right.$$

$$+ \left\{ N_2^3 \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) [(\mu + m + c)(\mu + \mu_v) + \mu \mu_v] \right\} \left\{ \beta_v \hat{I} \mu \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) + \beta \beta_v K \mu_v \hat{I} \mu^2 N_2^2 \right\}$$

$$\left. \left\{ [N_2^3 \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) [(\mu + m + c) + (\mu + \mu_v)] + \beta_v \hat{I} \mu \mu_v (\mu_v N_2 + \beta_v \hat{I} \mu) + \beta \beta_v K \mu_v \hat{I} \mu^2 N_2^2 \right\} \right.$$

$$\left. \left\{ N_2^2 \mu_v \beta_v \hat{I} \mu (\mu_v N_2 + \beta_v \hat{I} \mu) (2\mu + m + c) + N_2 \beta \mu^2 \beta_v K \mu_v \hat{I}^2 [m(\mu + c) + \beta_v \mu] \right. \right.$$

$$\left. \left. + N_2^2 \beta \mu \beta_v K \mu_v [\hat{I} (\mu + \mu_v)(\mu + c) + \hat{I} \mu \mu_v - N_2 \mu_v] \right\} \right\}$$

Hence $c_1c_2 - c_0c_3 > 0$ if $\left[[(\mu + m + c) + (\mu + \mu_v)] [(\mu + m + c)(\mu + \mu_v) + \mu \mu_v] \geq 1 \right.$

Hence, using Routh Hurwitz criteria we can conclude that $c_0x^3 + c_1x^2 + c_2x + c_3 = 0$ has roots with a negative real part.

4. Conclusion

From the analysis above, we get the dynamic of the mathematics model of Zika epidemic. These dynamic of model are connected with Reproduction number. The reproduction number can be used to determine whether the epidemic outbreak or vanish.

Acknowledgments

This research was funded by DRPM Ministry of Research, Technology and Higher Education, Indonesia especially in the scheme of Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) in 2019.

References

- [1] Buechler C R, Bailey A L, Weiler A M, Barry G L, Breitbach M E, Stewart L M, Jasinska A J, Freimer N B, Apetrei C, Phillips-Conroy J E, Jolly C J, Rogers J, Friedrich T C, O'Connor D H. 2017. *mSphere* **2**: e00392-16.
- [2] World Health Organisation (WHO). 2016. *Western Pacific Region*. Zika Virus.
- [3] European Centre for Disease Prevention and Control (ECDC), *Zika virus*. Online, April 2016.
- [4] Dick G W, Kitchen S F and Hadow A J. 1952. *Trans. R. Soc. Trop. Med. Hyg.* **46** 509.
- [5] Duffy M R, Chen T H, Hancock W T, Powers A M, Kool J L, Lanciotti R S, Pretrick M, Marfel M, Holzbauer S, Dubray C, Guillaumot L, Griggs A, Bel M, Lambert A J, Laven J, Kosoy O, Panella A, Biggerstaff B J, Fischer M, and Hayes E B. 2009. *New England Journal of Medicine* **360** 2536.
- [6] Caminade C, Turner J, Metelmann S, Hesson J C, Blagrove M S C, Solomon T, Morse A P, and Baylis M. 2017. *Proceedings of the National Academy of Sciences* **114** 119.
- [7] Cao-Lormeau V M, Roche C, Teissier A, Robin E, Berry AI, Mallet H P, Sall A A, and Musso D. 2014. *Emerging Infectious Disease* **20** 1085.
- [8] Zancula C, Mello V C, Mosimann A L, Santos G I, Santos C N, and Luz K. 2015. *Memorias do Instituto Oswaldo Cruz* **110** 569.

The dynamics of the basic model for the zika epidemic

ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	earchive.tpu.ru Internet Source	2%
2	G W Airlangga, I B G Rama Wisesa. "A review: microbiology perspective of zika virus", <i>Journal of Physics: Conference Series</i> , 2020 Publication	1%
3	cds.cern.ch Internet Source	1%
4	Guobao Feng, Lu Liu, Wanzhao Cui, Fang Wang. "Electron beam irradiation on novel coronavirus (COVID-19): A Monte-Carlo simulation*", <i>Chinese Physics B</i> , 2020 Publication	1%
5	www.ajbasweb.com Internet Source	1%
6	Anna Chadidjah, I G N M Jaya. "Forecasting dengue incidence in Bali by means latent Bayesian count data model", <i>Journal of Physics: Conference Series</i> , 2021 Publication	1%

7	Eleoui, M.. "Micromagnetic model of magnetization reversal of magnetically hard ultrathin dots and stripes", Journal of Magnetism and Magnetic Materials, 200411 Publication	1 %
8	hdl.handle.net Internet Source	1 %
9	Cai, L.M.. "Analysis of a SEIV epidemic model with a nonlinear incidence rate", Applied Mathematical Modelling, 200907 Publication	1 %
10	www.science.gov Internet Source	1 %
11	spravochnick.ru Internet Source	1 %
12	Submitted to UC, Boulder Student Paper	<1 %
13	Ana R. M. Carvalho, Carla M. A. Pinto. "Within-host and synaptic transmissions: contributions to the spread of HIV infection", Mathematical Methods in the Applied Sciences, 2017 Publication	<1 %
14	Dinita Rahmalia, Teguh Herlambang, Awawin Mustana Rohmah, Abdul Muhith. "Weights optimization using Firefly Algorithm on	<1 %

optimal control of zika disease from dengue symptoms by vaccination", Journal of Physics: Conference Series, 2020

Publication

15

time.com

Internet Source

<1 %

16

Jiraporn Lamwong, Puntani Pongsumpun. "The Zika Virus Transmission Model", International Journal of Bioscience, Biochemistry and Bioinformatics, 2017

Publication

<1 %

17

Pascal James Imperato. "The Convergence of a Virus, Mosquitoes, and Human Travel in Globalizing the Zika Epidemic", Journal of Community Health, 2016

Publication

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On

The dynamics of the basic model for the zika epidemic

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
