

# Masturi

*by Masturi Masturi*

---

**Submission date:** 19-Mar-2023 10:49AM (UTC+0700)

**Submission ID:** 2040339993

**File name:** Masturi\_2021\_\_AJC.pdf (599.46K)

**Word count:** 5326

**Character count:** 26916



13

Asian Journal of Chemistry; Vol. 33, No. 8 (2021), 1729-1735

## ASIAN JOURNAL OF CHEMISTRY

<https://doi.org/10.14233/ajchem.2021.23029>

## Determination of Total Phenol and Flavonoid Contents and Antioxidant Activity from Extract Fraction of Sappan Wood (*Caesalpinia sappan* L.) by Liquid-Liquid Extraction and Vacuum Liquid Chromatography

MASTURI<sup>1,\*</sup>, D. ALIGHIRI<sup>2</sup>, S.S. EDIE<sup>1</sup>, U. HANISYIFA<sup>2</sup> and A. DRASTISANTI<sup>3</sup><sup>1</sup>Department of Physics, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Semarang, Indonesia<sup>2</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang, Semarang, Indonesia<sup>3</sup>Department of Chemistry Education, Faculty of Sciences and Technology, Universitas Islam Negeri Walisongo Semarang, Semarang, Indonesia

\*Corresponding author: E-mail: masturi@mail.unnes.ac.id

Received: 16 October 2020;

Accepted: 20 April 2021;

Published online: 26 July 2021;

AJC-20420

This study reported the antioxidant activity of sappan wood (*Caesalpinia sappan* L.) extract and their extract fraction from the liquid-liquid extraction and vacuum liquid chromatography. The study comprises with phytochemical assessments, evaluation of total phenolic content (TPC) and total flavonoid content (TFC). The evaluation of antioxidant activity was carried out by using the DPPH method. The effectiveness of antioxidant activity in sappan wood compared to quercetin. Extract of sappan wood (ESW) yields of 5.40%, extract fraction from liquid-liquid extraction (FE LLE) yields of 52.15% and extract fraction from vacuum liquid chromatography (FE VLC) with silica gel and eluent sequential from *n*-hexane, 5% *n*-hexane/95% ethyl acetate, ethyl acetate and 5% ethyl acetate/45% methanol was produced 60 of fractions extract. The thin-layer chromatography (TLC) was analyzed using silica gel as a stationary phase with the mobile phase of *n*-hexane:ethyl acetate (1:0.55) obtained 10 of fractions extract in fractions of 6 and 7, which showed the most positive results. The phytochemical test indicated the presence of phenolic and flavonoids compounds. The highest values for TPC were FE LLE (216.67 mg GAE/g), FE VLC (18.33 mg GAE/g) and ESW (18.33 mg GAE/g), respectively. The highest values for TFC were FE VLC (152.48 mg QE/g), FE LLE (66.2 mg QE/g) and ESW (48.4 mg QE/g), respectively. Then, the highest antioxidants activity were FE VLC (5.42 µg/mL), FE LLE (6.41 µg/mL), quercetin (6.52 µg/mL) and ESW (7.82 µg/mL), respectively. The results showed that the highest of TFC was related to the antioxidant activity. However, the highest value of TPC was not showed the highest antioxidant activity. Both ESW, FE LLE and FE VLC with quercetin as control exhibited powerful antioxidants.

**Keywords:** *Caesalpinia sappan*, Antioxidant, Total phenolic content, Total flavonoid content.

### INTRODUCTION

Oxidative pressure is described as an imbalance of oxygen free radicals or reactive oxygen species (ROS) with the organism's capacity to counteract the oxidation process [1]. ROS is a highly reactive molecule, such as peroxy, hydroxyl radicals, superoxide anion, hydrogen peroxide and singlet oxygen, which are produced as a byproduct of normal metabolism under aerobic cell condition [2,3]. In general, ROS are produced in medium concentration to maintain normal physiological function in an organism. However, excess production of ROS causes the high concentration, so that it readily reacts with DNA, protein

and lipids that cause oxidative damage and cause many chronic diseases [4-6]. Also, oxidative pressure is responsible for decreasing physiological function and food quality, that causes food became toxic, rancid smell and damage crucial biochemical component of food [7].

Antioxidants can prevent chronic disease and protect food ingredients by cleaning free radicals that will inhibit the oxidation reaction process [8,9]. The body naturally obtains antioxidants from food ingredients [10-12], containing vitamins or phytochemicals. Generally, vitamins that function as antioxidants are vitamin A (retinol), C (ascorbic acid) and E ( $\alpha$ -tocopherol) [13,14]. Examples of phytochemical compounds known as

This is an open access journal, and articles are distributed under the terms of the Attribution 4.0 International (CC BY 4.0) License. This license lets others distribute, remix, tweak, and build upon your work, even commercially, as long as they credit the author for the original creation. You must give appropriate credit, provide a link to the license, and indicate if changes were made.

antioxidants are phenol or polyphenol and flavonoids [15-17]. There are also synthetic antioxidants which are often used in food additives, namely butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA) and tetrahydroxyquinone (TBHQ). The use of synthetic antioxidants is due to their low price and effectively prevents oxidation reactions in foodstuffs. However, excessive use of synthetic materials will harm health [18].

One of Indonesia's abundant and cheap materials is secang (sappan) wood (*Caesalpinia sappan* L.), which is widely used sappan plants in overcoming diseases [19,20]. Sappan wood consists of flavonoid, brazilin, alkaloid, saponin, tannin, phenyl propane and terpenoid [21]. Traditionally, sappan wood is used as a colouring agent in food [22]. Several studies explained that phenolic compounds, polyphenols and flavonoids, show potential antioxidant activity. Previous research [23] has proven that sappan bark has xanthine oxidase inhibitory activity. A methanol extract of sappan wood has also been shown to deal with the xanthine oxidase enzyme activity [24,25]. However, research on the extract's antioxidant activity, especially as an antioxidant, is still limited. This research was conducted to study the antioxidant activity of sappan wood extract and its extract fraction from the liquid-liquid extraction and vacuum liquid chromatography.

## EXPERIMENTAL

The chemicals used in this study viz. *n*-hexane, ethyl acetate, chloroform, acetic anhydride, H<sub>2</sub>SO<sub>4</sub>, methanol, ethanol, FeCl<sub>3</sub> and Na<sub>2</sub>CO<sub>3</sub> (Merck, Germany) were of analytical grade. Silica gel 60 (0.040-0.063 mm, 300-400 mesh), Dragendorf reagent, Mayer reagent, Folin-Ciocalteu reagent, gallic acid, quercetin and DPPH reagent were procured from Sigma-Aldrich, USA.

Sappan wood was obtained from the Kendal area, Central Java and its identification was carried out at the Biology Laboratory, Universitas Negeri Semarang. Sappan wood (*Caesalpinia sappan* L.) was cleaned and dried in the open air. The dry sample was blended and weighed as much as 2 kg.

**Extraction of sappan wood:** Sappan wood dry powder (2 kg) was extracted by maceration method using *n*-hexane solvent for 5 × 24 h, then filtered. The obtained residue was macerated again with methanol for 7 × 24 h and filtered. The filtrate was concentrated using a rotary evaporator for the extraction process.

**Sappan wood fractionation by liquid-liquid extraction (LLE):** The liquid-liquid extraction (LLE) was carried out by dissolving methanolic extract with 400 mL of *n*-hexane with constant stirring and then, transferred to a separating funnel containing 400 mL of water. The mixture was then shaken several times and obtained the *n*-hexane fraction. The remaining water fraction obtained was re-partitioned using ethyl acetate to obtain the ethyl acetate fraction. The *n*-hexane and ethyl acetate fractions were then concentrated using a rotary evaporator.

**Sappan wood fractionation by vacuum liquid chromatography (VLC):** The ethyl acetate extract fraction obtained from LLE was fractionated by vacuum liquid chromatography (VLC). The sample mixed with silica gel 60 (0.040-0.063 mm,

230-400 mesh) until homogenous. Then, put in vacuum liquid column and compacted. Samples were eluted by *n*-hexane 100%, ethyl acetate (5-95%), ethyl acetate 100% and ethyl acetate: methanol (5:45%). The fraction obtained from separation by VLC was collected in a glass bottle and then concentrated by a rotary evaporator. The fraction obtained was analyzed by the separation pattern and its components using the TLC.

## Phytochemical analysis of sappan wood

**Alkaloid compound test:** The alkaloid test was carried out with Dragendorf reagent. Ethanol (5 mL) was added in the sappan wood extract (1 g) followed by the addition of 2 drops of Dragendorf reagent resulted in the formation of reddish-orange precipitates [26].

**Test for steroid and terpenoid:** Sappan wood extract (1 g) was added with 5 mL ethanol, 2 mL chloroform and 2 drops of acetic acid anhydride. Then, 2 drops of conc. H<sub>2</sub>SO<sub>4</sub> were added through the walls of the test tube. The formation of green or blue colour indicated the presence of steroids. Meanwhile, if a reddish-purple colour is formed, the extract positively contains terpenoids [26].

**Flavonoid test:** Sappan wood extract (1 g) was added with 5 mL of ethanol. The solution was added with 1 mL of hot methanol and magnesium metal on the spatula's tip and a 0.5 mL of conc. HCl. The formation of a red or orange solution indicated the presence of flavonoid [26].

**Phenolic test:** Sappan wood extract (1 g) was mixed with 5 mL of ethanol followed by the addition of 1 mL of FeCl<sub>3</sub> solution. A purple-blue colour formation confirmed the presence of phenolic compounds [26].

**Analysis of extract fractions from vacuum liquid chromatography (VLC) by thin layer chromatography (TLC):** The fractions of VLC were analyzed for their components by TLC. Fractions had similar spots or stains and then the TLC analysis was carried out again. The retention factor (R<sub>f</sub>) value was calculated.

**Determination of total phenolic content (TPC):** The maximum wavelength was determined using a UV-vis spectrophotometer using gallic acid as the standard solution. Separately, 1 mL of gallic acid and 0.5 mL of sample solution was added to 5.9 mL of Folin-Ciocalteu reagent, 30 mL of distilled water and stirred for 5 min. These two solutions were added with 7.5 mL of 7.5% (w/v) Na<sub>2</sub>CO<sub>3</sub> solution and distilled water until the volume became 50 mL. The mixture was stirred and incubated for 30 min in the dark at room temperature (28-30 °C). After 30 min, the sample solution was subjected to a wavelength scan at 600-800 nm and showed the maximum wavelength absorption at 765 nm.

**Preparation of standard calibration curves for gallic acid for the determination of TPC:** Gallic acid (50 mg) was dissolved in ethanol and made up the final volume of 50 mL. Then gallic acid solution was taken from 0.25 to 5 mL and diluted with distilled water to a final volume of 10 mL that the concentration of the gallic acid solution was 25-500 µg/mL. Each 2.5 mL of standard gallic acid solution (50-250 µg/mL) was mixed with 2.5 mL of Folin-Ciocalteu reagent in a 50 mL volumetric flask. After mixing for 5 min, 7.5 mL of 7.5%

(w/v) sodium carbonate solution was added and distilled water until the volume was 50 mL. The mixture was stirred and incubated for 30 min in the dark at room temperature (28-30 °C). The absorbance was measured at a maximum wavelength of 765 nm against the blank solution using a UV-vis spectrophotometer. The calibration curve obtained was expressed in the linear regression equation at concentrations of 25-500 µg/mL. The regression equation is  $y = 0.001x + 0.011$ ;  $R^2 = 0.986$ . The phenol concentration was expressed in µg/mL on a standard curve. The TPC obtained was expressed as mg GAE (gallic acid equivalent)/g of a sample. All the determinations were made three times replication.

**Determination of total flavonoid content (TFC):** Determination of the maximum wavelength was carried out using a UV-vis spectrophotometer using quercetin as standard solution. Quercetin solution (2 mL) and the sample solution (1 mL) were added separately to 3 mL of 0.1 mol/mL solution of  $AlCl_3$  solution. Then, the two solutions were mixed and diluted with 70% ethanol solution. After 0.5 h, the sample solution was scanned at 400-500 nm and showed the maximum wavelength absorption at 425 nm [27].

**Preparation of quercetin standard calibration curves for the determination of TFC:** Quercetin (50 mg) was dissolved in ethanol to a final volume of 50 mL. The quercetin solution was taken 0.25-2 mL and diluted with distilled water to a final volume of 10 mL, hence the concentration of the quercetin solution became 40-180 µg/mL. Each concentration of quercetin solution (2 mL) was added to 2 mL of 2%  $AlCl_3$  solution and 2 mL of 20 mM potassium acetate solution. Samples were incubated for 1 h at room temperature (28-30 °C). The absorbance was measured at a maximum wavelength of 425 nm against the blank solution using a UV-vis spectrophotometer. The calibration curve was expressed in a linear regression equation at a 25-200 µg/mL [27]. The regression equation is  $y = 0.005x - 0.025$ ;  $R^2 = 0.992$ . Flavonoid concentrations are expressed in µg/mL on a standard curve. The TFC obtained was expressed as mg QE (quercetin equivalent)/g of a sample. All the determinations were made by replication three times.

**Antioxidant activity:** Determination of antioxidant activity was conducted using free radical scavenging activity based on 1,1-diphenyl-2-picrylhydrazyl (DPPH) method. DPPH solution (1 mL) was added with 2 mL of methanol and left for 30 min in the dark at room temperature (28-30 °C) and the absorption was measured with a UV-vis spectrophotometer at a wavelength ranging from 510-520 nm to obtain the maximum wavelength. The absorbance of DPPH was determined before-hand as control at a maximum wavelength of 517 nm.

Each sample (2 mg) of ESW, LLE and VLC and quercetin were weighed and dissolved separately using 10 mL methanol. The solution was mixed in a 100 mL volumetric flask and methanol was added to the limit mark so that a solution concentration of 1000 µg/mL was obtained. Antioxidant testing was carried out by taking 3 mL of sample test solution and adding 1 mL of DPPH solution in methanol and diluting it with methanol so that the concentration becomes 10-50 µg/mL with 1 mL of methanol as a blank solution. The mixture was left for 0.5 h in a dark place at room temperature (28-30 °C). Each antioxidant

activity test was carried out three times. The absorbance measurement results have analyzed the percentage of their antioxidant activity using eqn. 1:

$$\text{Inhibition (\%)} = \frac{\text{Blank absorbance} - \text{Sample absorbance}}{\text{Blank absorbance}} \times 100 \quad (1)$$

The linear regression line equation obtained was used to evaluate the  $IC_{50}$  value.

## RESULTS AND DISCUSSION

The extraction process was carried out to take the chemical compounds contained in the sample. The principle of extraction is based on the mass transfer of the dissolved substance components into the solvent. There is a displacement of the interface layer and diffuses into the solvent-extraction of sappan wood (*C. sappan* L.) using the maceration method with *n*-hexane and methanol solvents. The maceration method was chosen since, it is easy and does not use high temperatures, which might damage chemical compounds with antioxidant activity in the sappan wood simplicia. The macerated filtrate was concentrated with a rotary evaporator to obtain 2.1436 g thick yellow *n*-hexane extract with a 0.10718% yield and 108.0855 g thick brown methanol extract with 5.404275% yield.

**Fractionation of secang wood extract with liquid-liquid extraction (LLE):** The thick extract of sappan wood was then fractionated using the liquid-liquid partition extraction method. The fractionation of the liquid-liquid partition extraction was carried out using *n*-hexane (non-polar), ethyl acetate (semi-polar) and the remaining solvent in water (polar). The purpose of this method is to separate chemical compounds based on their level of polarity. This liquid-liquid extraction process is repeated until clear *n*-hexane and ethyl acetate solvents are obtained, indicating no more sample components dissolved in *n*-hexane ethyl acetate solvents. The results of the extraction of the liquid-liquid partition were then concentrated with a rotary evaporator so that the *n*-hexane fraction was 0.0161 g and the ethyl acetate fraction was 56.37 g with a yield of 52.15% from the initial wood methanol extract.

**Fractionation of secang wood extract by vacuum liquid chromatography (VLC):** The ethyl acetate fraction was separated by vacuum liquid chromatography (VLC) method. The working principle of VLC is the partition and adsorption of compound components in the extracted sample whose separation is assisted by pressure from a vacuum device. Separation with VLC aims to determine which subfraction has antioxidant activity. The VLC results obtained 60 fractions, which were then analyzed by TLC to see the chromatographic profile.

The fractions that have similar spots are put together and 10 subfractions are obtained. The spot profile on TLC of the 10 subfractions obtained is seen using UV lamps at 366 and 254 nm (Fig. 1). The  $R_f$  value for each subfraction is shown in Table-1.

Table-1 shows that the  $R_f$  value of the subfraction 1-10 of the VLC yield varies from the sappan wood extract. However, fractions, F6 and F7 have the same  $R_f$  values: spot 1 = 0.2, 2 = 0.3, 3 = 0.35, 4 = 0.5 and 5 = 0.8. According to Rusnaeni & Fitria [19], two or more compounds can be considered identical




Fig. 1. Thin-layer chromatography (TLC) profile of subfractions 1-10 of sappan wood (*Caesalpinia sappan* L.) extract resulted from vacuum liquid chromatography (VLC) using (a) 366 mm and (b) 254 mm UV lamp

TABLE-1  
R<sub>f</sub> VALUE, NUMBER OF SPOTS AND SPOT COLOUR IN SUBFRACTIONS 1-10 FROM SAPPAN WOOD (*Caesalpinia sappan* L.) EXTRACT RESULTED FROM VACUUM LIQUID CHROMATOGRAPHY (VLC) USING THE ELUENT RATIO OF *n*-HEXANE AND ETHYL ACETATE (1:0.55)

Subfraction	Number of spots	Retention factor (R <sub>f</sub> )	Colour
F1	1	0.9	Purple
F2	1	0.9	Blue
F3	3	0.6	Purple
		0.7	Blue
		0.9	Blue
F4	7	0.2	Yellow
		0.3	Yellow
		0.4	Purple
		0.5	Purple
		0.6	Purple
		0.7	Blue
		0.9	Blue
F5	5	0.2	Orange
		0.3	Yellow
		0.4	Purple
		0.5	Purple
		0.8	Blue
F6	5	0.2	Brown
		0.3	Green
		0.35	Yellow
		0.5	Purple
		0.8	Blue
F7	5	0.2	Brown
		0.3	Green
		0.35	Yellow
		0.5	Purple
		0.8	Blue
F8	3	0.2	Purple
		0.3	Orange
		0.4	Yellow
F9	3	0.2	Blue
		0.3	Orange
		0.4	Green
F10	2	0.2	Blue
		0.4	Green

if they have the same R<sub>f</sub> values with the same TLC conditions. Then, F6 and F7 are taken together to test its antioxidant activity because of its similarity to quercetin, which is likely to contain numerous flavonoids.

Phytochemical analysis of sappan wood extract revealed that sappan wood (*C. sappan* L.) can detect secondary metabolites in alkaloid, terpenoid, phenolic and flavonoid compounds. However, it cannot detect steroid compounds. Table-1 presents the results of the phytochemical assay on the detected secondary metabolites inside sappan wood.

The presence of phenolic and flavonoid compounds in sappan wood indicates that it can have antioxidant activities, and can be further analyzed quantitatively for phenolic and flavonoid compounds through the total phenolic and flavonoid contents by using the Folin-Ciocalteu and AlCl<sub>3</sub> methods, respectively. Table-2 presents the results of the phytochemical assay of sappan wood.

TABEL-2  
PHYTOCHEMICAL ASSAY OF SAPPAN WOOD (*Caesalpinia sappan* L.)

Compound	Phytochemical assay*			Reaction result
	ESW	FE LLE	FE VLC	
Alkaloid	+	+	+	Orange sediment
Steroid	-	-	-	Red
Terpenoid	+	+	+	Red
Phenolic	+	+	+	Blackish blue
Flavonoid	+	+	+	Red

\*+: Positive/Detected; -: Negative/Undetected

ESW = Extract of sappan wood; FE LLE = Fraction of sappan wood extract from liquid-liquid extraction; FE VLC = Fraction of sappan wood extract from vacuum liquid chromatography.

For the evaluation of total phenolic content (TPC), phenol compounds were quantitatively analyzed using a UV-visible spectrophotometer to determine the phenol content in the sappan wood extract. The extract fraction from the liquid-liquid partition resulted from vacuum liquid chromatography.

In this study, the Folin-Ciocalteu method was used to determine the TPC of the sample as a standard solution with a concentration series of 50, 100, 150, 200, and 250 µg/mL. The results showed that the maximum wavelength of quercetin is 765 nm. The maximum wavelength used to measure the absorption curve of gallic acid calibration is shown in Fig. 2. These results suggested that the higher the concentration used, the higher the absorbance value obtained. The result of the standard curve of gallic acid obtained is plotted between the level and the absorbance, so that the linear regression equation was obtained ( $y = 0.001x + 0.011$ ) with the  $R^2$  value of 0.986. The gallic acid calibration curve equation can be used as a comparison to determine the total phenolic compound concentration in the sample extract.




Fig. 2. Calibration curve of gallic acid at a maximum wavelength of 765 nm

Table-3 shows that the highest TPC was found in the extract fraction from the liquid-liquid extraction. The lowest TPC was noted in the extract fraction from VLC. The result indicates that the liquid-liquid extraction process could separate components other than phenolic compounds. Compounds such as polysaccharides from amylose, cellulose, hemicellulose and lignin were not present in the extract fraction from VLC. Then, the TPC in the extract fraction from VLC showed the lowest result, possibly because only certain compounds according to their polarity present in the extract fraction.

Sample	TPC (mg GAE/g)
ESW	183.33
FE LLE	216.67
FE VL	160.83

ESW = Extract of sappan wood; FE EPC = Fraction of sappan wood extract by liquid-liquid extraction; FE VLC = Fraction of extract sappan wood by vacuum liquid chromatography.

**Evaluation of total flavonoid content (TFC):** The quantitative analysis of flavonoids was conducted using a UV-visible

spectrophotometer using  $AlCl_3$  method. This method can be used to determine the flavonoid levels in the sappan wood extract, the extract fraction from the LLE and the extract fraction from the results of VLC. In this study, quercetin was used to determine the TFC in the sample as a standard solution with a concentration series of 40, 60, 70, 80, 100, 120, 140, 160 and 180 µg/mL. The results showed that the maximum wavelength of the standard quercetin is 425 nm. The maximum wavelength used to measure the absorption of the wood extract sample, the extract fraction from the LLE, and the extract fraction from the VLC are shown in Fig. 3, which indicate that the higher the absorbance was obtained at higher concentration. The quercetin calibration curve equation can be used as a comparison to determine the TFC in the sample extract.




Fig. 3. Quercetin calibration curve at a maximum wavelength of 425 nm

Quantitative analysis with a UV-visible spectrophotometer used a blank solution as control, which functions as a blank to multiply zero compounds that do not need to be analyzed. In TFC measurement, the sample solution was added to  $AlCl_3$ , which can form a complex, resulting in a shift in the wavelength towards the visible light, marked with the solution producing a yellow colour. The addition of potassium acetate aims to maintain the wavelength in the visible area. Incubation treatment for 1 h before the measurement was intended to stabilize the resulting colour intensity.

Table-4 shows that the highest yield was found in the extract fraction from the result of VLC. At the same time, the lowest yield was found in the extract. The polarity level of the sample will affect the TFC. These results indicate that the VLC process has most likely succeeded in separating flavonoids based on the polarity degree of other sappan wood components. The TFC results of the extract fraction of the LLE still contained the largest flavonoids other than phenolic compounds.

**Antioxidant activity:** In this study, the antioxidant activity was performed using the DPPH method. A quantitative antioxidant activity testing was performed for sappan wood extract samples, extract fraction from LLE and extract fraction (F6-F7) from VLC, along with positive quercetin control at a wave-

TABLE-4  
EVALUATION OF TOTAL FLAVONOID CONTENT  
IN SAPPAN WOOD (*Caesalpinia sappan* L.)

Sample	Total flavonoid content (mg QE/g)
ESW	48.40
FE LLE	66.20
FE VL	152.48

ESW = Extract of sappan wood; FE EPC = Fraction of sappan wood extract by liquid-liquid extraction; FE VLC = Fraction of extract sappan wood by vacuum liquid chromatography.

length of 517 nm. The antioxidant activity results of sappan wood extract and extract fraction from LLE and VLC with quercetin compared with TPC and TFC are shown in Table-5.

TABLE-5  
COMPARISON OF TOTAL PHENOLIC CONTENT (TPC)  
AND TOTAL FLAVONOID CONTENT (TFC) WITH THE  
ANTIOXIDANT ACTIVITY OF EXTRACTS AND EXTRACT  
FRACTIONS FROM SAPPAN WOOD (*Caesalpinia sappan* L.)  
WITH QUERCETIN AS A COMPARISON INDICATED BY  
THE VALUE OF IC<sub>50</sub> (THE HALF MAXIMAL  
INHIBITORY CONCENTRATION)

Sample*	TPC** (mg GAE/g)	TFC*** (mg QE/g)	IC <sub>50</sub> (µg/mL)
ESW	183.33	48.4	7.82
FE LLE	216.67	66.2	6.41
FE VLC	160.83	152.48	5.42
Quercetin	–	–	6.52

\*ESW = Ekstrak sappan wood; FE LLE = Fraction of sappan wood extract by liquid-liquid extraction; FE VLC = Fraction of sappan wood by liquid vacuum chromatography; \*\*TPC = Total phenolic content which obtained and expressed as mg QE (quercetin equivalent)/g sample; \*\*\*TFC = Total flavonoid content which obtained and expressed as mg QE (quercetin equivalent)/g sample.

Table-5 shows that the IC<sub>50</sub> was the lowest at 5.42 µg/mL concentration of the sappan wood extract VLC, whereas the lowest was observed at 7.82 µg/mL concentration of the sappan wood extract. A lower IC<sub>50</sub> value indicated a higher radical scavenging effect. The smaller the IC<sub>50</sub> value, the higher the antioxidant activity. The sample has powerful, strong, moderate, or weak antioxidants depending on whether its IC<sub>50</sub> value is < 50, 50-100, 100-150 or 150-200 µg/mL, respectively. The samples with the strongest to the weakest antioxidants are FE VLC (5.42 µg/mL), FE LLE (6.41 µg/mL), quercetin (6.52 µg/mL), and ESW (7.82 µg/mL).

Additionally, present study showed that the higher the TFC, the higher the antioxidant activity (Table-5). The FE LLE sample, which had a TFC of 152.48 mg QE/g, showed the highest antioxidant activity with an IC<sub>50</sub> value of 5.42 µg/mL. Meanwhile, a high TPC does not necessarily indicate a high antioxidant activity. The FE LLE sample, with a TPC of 216.67 mg GAE/g, had antioxidant activity of 6.41 µg/mL, which was lower than that of FE VLC. In this study, flavonoid compounds in sappan wood are found to be more effective than phenols as antioxidants in inhibiting the free radical activity by up to 50%.

## Conclusion

Sappan wood extract fractions of liquid-liquid extraction (LLE) and vacuum liquid chromatography (VLC) were analyzed

for antioxidant activity by using the DPPH method. Both FE VLC (5.42 µg/mL) and FE LLE (6.41 µg/mL) fractionates consisted of potent antioxidant activities. The sappan wood extract of TPC was higher than TFC whereas, TFC has a higher antioxidant activity. That means a high TPC count does not necessarily indicate a high antioxidant activity. Moreover, this study showed that sappan wood flavonoid compounds are more effective than phenols as antioxidants in inhibiting free radical activity by 50%.

## ACKNOWLEDGEMENTS

The authors thank the Directorate General of Higher Education, Indonesia for the research through DRPM 2020-2021.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interests regarding the publication of this article.

## REFERENCES

- Z.K. Abbas, S. Saggi, M.I. Sakeran, N. Zidan, H. Rehman and A.A. Ansari, *Saudi J. Biol. Sci.*, **22**, 322 (2015); <https://doi.org/10.1016/j.sjbs.2014.11.015>
- N. Chattopadhyay, T. Ghosh, S. Sinha, K. Chattopadhyay, P. Karmakar and B. Ray, *Food Chem.*, **118**, 823 (2010); <https://doi.org/10.1016/j.foodchem.2009.05.069>
- D.Y. Choi, Y.J. Lee, J.T. Hong and H.J. Lee, *Brain Res. Bull.*, **87**, 144 (2012); <https://doi.org/10.1016/j.brainresbull.2011.11.014>
- J. Dai and R.J. Mumper, *Molecules*, **15**, 7313 (2010); <https://doi.org/10.3390/molecules15107313>
- M.H. Dicko, H. Gruppen, A.S. Traore, W.J.H. van Berkel and A.G.J. Voragen, *J. Agric. Food Chem.*, **53**, 2581 (2005); <https://doi.org/10.1021/jf0501847>
- H. Jabeen, S. Saleemi, H. Razzaq, A. Yaqub, S. Shakoor and R. Qureshi, *J. Photochem. Photobiol. B*, **180**, 268 (2018); <https://doi.org/10.1016/j.jphotobiol.2018.02.006>
- H. Kikuzaki, M. Hisamoto, K. Hirose, K. Akiyama and H. Taniguchi, *J. Agric. Food Chem.*, **50**, 2161 (2002); <https://doi.org/10.1021/jf011348w>
- J. Liu, H. Sun, F. Dong, Q. Xue, G. Wang, S. Qin and Z. Guo, *Carbohydr. Polym.*, **78**, 439 (2009); <https://doi.org/10.1016/j.carbpol.2009.04.030>
- M. Hasanah, B. Maharani and E. Munarsih, *Indo. J. Pharm. Sci. Technol.*, **4**, 42 (2017); <https://doi.org/10.15416/ijpst.v4i2.10456>
- J.P. Maro, A.H. Alimuddin and Harlia, *J. Kimia Khatulistiwa*, **4**, 35 (2015).
- Masturi, D. Alighiri, S.S. Edie, A. Drastisianti, U. Khasanah, K.A. Tanti, Susilawati, R.Z. Maghfiroh, K.G.C. Kirana and F. Choirunnisa, *J. Phys. Conf. Ser.*, **1567**, 042084 (2020); <https://doi.org/10.1088/1742-6596/1567/4/042084>
- Masturi, D. Alighiri, K. Nuzulina, M. Rodhiyah and A. Drastisianti, *J. Phys. Conf. Ser.*, **1321**, 022041 (2019); <https://doi.org/10.1088/1742-6596/1321/2/022041>
- P. Molyneux, *Songklanakarin J. Sci. Technol.*, **26**, 211 (2004).
- M.T. Nguyen, S. Awale, Y. Tezuka, Q.L. Tran, H. Watanabe and S. Kadota, *Biol. Pharm. Bull.*, **27**, 1414 (2004); <https://doi.org/10.1248/bpb.27.1414>
- G.R. Nimade, S.D. Agus and A. Komang, *J. Ilmu Teknol. Pangan*, **8**, 216 (2019).
- P. Ohama and N. Tumpat, *Int. J. Mater. Textile Eng.*, **8**, 432 (2014); <https://doi.org/10.5281/zenodo.1092830>
- T. Persson, B.O. Popescu and A. Cedazo-Minguez, *Oxid. Med. Cell. Longev.*, **2014**, 1 (2014); <https://doi.org/10.1155/2014/427318>

18. H.M. Pertamawati and M. Hardhiyuna, *Kartika J. Ilmiah Farm.*, **3**, 12 (2015); <https://doi.org/10.26874/kjif.v3i2.99>
19. D.I.S. Rusnaeni and L. Fitria, *J. Pharm. (Caino)*, **13**, 84 (2016).
20. K. Sugamura and J.F. Keaney Jr., *Free Radic. Biol. Med.*, **51**, 978 (2011); <https://doi.org/10.1016/j.freeradbiomed.2011.05.004>
21. W. Tan, Q. Li, T. Zhou, Q. Chen, G. Wang, F. Dong and Z. Guo, *Bioorg. Chem.*, **74**, 66 (2017); <https://doi.org/10.1016/j.bioorg.2017.07.005>
22. S.P. Umasari, H.M. Ana and R. Ayu, Prosiding Seminar Nasional Mahasiswa Unimus, vol. 1, p. 283 (2018) (In Indonesian).
23. M. Valko, D. Leibfritz, J. Moncol, M.T. Cronin, M. Mazur and J. Telser, *Int. J. Biochem. Cell Biol.*, **39**, 44 (2007); <https://doi.org/10.1016/j.biocel.2006.07.001>
24. J.A. Vinson, X. Su, L. Zubik and P. Bose, *J. Agric. Food Chem.*, **49**, 5315 (2001); <https://doi.org/10.1021/jf0009293>
25. B. Xu, S. Guan, B. Wang, S. Wang, Y. Wang, C. Sun, X. Ma and T. Liu, *Int. J. Biol. Macromol.*, **109**, 1 (2018); <https://doi.org/10.1016/j.ijbiomac.2017.12.019>
26. J.L.B. Zanin, B.A. de Carvalho, P. Salles Martineli, M.H. dos Santos, J.H.G. Lago, P. Sartorelli, C. Viegas Jr. and M.G. Soares, *Molecules*, **17**, 7887 (2012); <https://doi.org/10.3390/molecules17077887>
27. C. Zeng, M. Su, Q. Chen, Q. Chang, W. Wang and H. Li, *Carbohydr. Polym.*, **153**, 391 (2016); <https://doi.org/10.1016/j.carbpol.2016.07.067>



# Masturi

---

## ORIGINALITY REPORT

---

22%

SIMILARITY INDEX

15%

INTERNET SOURCES

16%

PUBLICATIONS

7%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	<a href="http://oaji.net">oaji.net</a> Internet Source	2%
2	<a href="http://www.ubu.ac.th">www.ubu.ac.th</a> Internet Source	1%
3	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	1%
4	<a href="http://academic-accelerator.com">academic-accelerator.com</a> Internet Source	1%
5	<a href="http://talenta.usu.ac.id">talenta.usu.ac.id</a> Internet Source	1%
6	Loudfy Eka Widyawaty, I Nyoman Ehrich Lister, Sukirman Lie. "Antioxidant Activity, Total Phenol, and Total Flavonoid of Syzygium Polyanthum", 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce), 2021 Publication	1%
7	<a href="http://www.orientjchem.org">www.orientjchem.org</a> Internet Source	1%

---

8	Submitted to University of Newcastle upon Tyne Student Paper	<1 %
9	Submitted to Ganpat University Student Paper	<1 %
10	I. Wayan Sutapa, Mirella Fonda Maahury, Rahayu, Marthinus J. Saptenno, Dewi Sri, Adriani Bandjar. "Mayang fruit juice innovation for rheumatism medicine and non dairy source of calsium", AIP Publishing, 2023 Publication	<1 %
11	mdpi-res.com Internet Source	<1 %
12	Submitted to Rochester Institute of Technology Student Paper	<1 %
13	eprints.ums.edu.my Internet Source	<1 %
14	Hao Zhang, Lu Jiang, Shu Ye, Yubin Ye, Fazheng Ren. "Systematic evaluation of antioxidant capacities of the ethanolic extract of different tissues of jujube (Ziziphus jujuba Mill.) from China", Food and Chemical Toxicology, 2010 Publication	<1 %
15	Akiri SVC Rao, Sareddy G Reddy, Phanithi P Babu, Attipalli R Reddy. "The antioxidant and	<1 %

antiproliferative activities of methanolic extracts from Njavara rice bran", BMC Complementary and Alternative Medicine, 2010

Publication

16

"Ethnobotany of the Mountain Regions of Southeast Asia", Springer Science and Business Media LLC, 2021

Publication

<1 %

17

Edible Medicinal and Non Medicinal Plants, 2015.

Publication

<1 %

18

N. J. George, I. B. Obot, A. N. Ikot, A. E. Akpan, N. O. Obi-Egbedi. " Phytochemical and Antimicrobial Properties of Leaves of ", E-Journal of Chemistry, 2010

Publication

<1 %

19

[www.preinvestigo.biblioteca.uvigo.es](http://www.preinvestigo.biblioteca.uvigo.es)

Internet Source

<1 %

20

[www.worldresearchersassociations.com](http://www.worldresearchersassociations.com)

Internet Source

<1 %

21

Pooja D. Motikar, Pavankumar R. More, Shalini S. Arya. "A novel, green environment-friendly cloud point extraction of polyphenols from pomegranate peels: a comparative assessment with ultrasound and microwave-

<1 %

assisted extraction", Separation Science and Technology, 2020

Publication

---

22

Hasim ., Dimas Andrianto, Wira Islamiati, Annisa Farhiah Wahdah Ham, Didah Nur Farida. "Antioxidant Activity of Ethanol Extract of Red Yeast Rice and its Fractionation Products", Research Journal of Phytochemistry, 2018

Publication

---

<1 %

23

Submitted to Arkansas Tech University

Student Paper

---

<1 %

24

Muhammad Deni Anugerah, Adawiah Daud, Hermin Sulistyarti, Akhmad Sabarudin, Rurini Retnowati. "Method development for the determination of aluminum by  $\mu$ PAD using sappan wood extract (Caesalpinia sappan)", AIP Publishing, 2022

Publication

---

<1 %

25

Submitted to The Robert Gordon University

Student Paper

---

<1 %

26

Submitted to University College London

Student Paper

---

<1 %

27

Wenqiang Tan, Qing Li, Tingting Zhou, Qiuhong Chen, Gang Wang, Fang Dong, Zhanyong Guo. "Synthesis and antioxidant

<1 %

ability of 6,6'-diamino-6,6'-dideoxytrehalose",  
Bioorganic Chemistry, 2017

Publication

28

Muthulisi Siwela, John R.N. Taylor, Walter A.J. de Milliano, Kwaku G. Duodu. "Influence of phenolics in finger millet on grain and malt fungal load, and malt quality", Food Chemistry, 2010

Publication

<1 %

29

Submitted to University of the Philippines Diliman

Student Paper

<1 %

30

[journals.plos.org](http://journals.plos.org)

Internet Source

<1 %

31

[s3.eu-central-1.amazonaws.com](http://s3.eu-central-1.amazonaws.com)

Internet Source

<1 %

32

[www.phytojournal.com](http://www.phytojournal.com)

Internet Source

<1 %

33

[1library.net](http://1library.net)

Internet Source

<1 %

34

R Susanti, Nur Rahayu Utami, Nur Rahayu Utami, Margareta Rahayuningsih, Fidia Fibriana, Margareta Rahayuningsih, Fidia Fibriana. "Identification of Avian Influenza Genetic Resistance Gene Marker in Chickens", Makara Journal of Science, 2017

Publication

<1 %

- 35 Sunitha Dontha, Kamurthy Hemalatha, Mantripragada Bhagavan Raju. "Phytochemical Evaluation and Screening of in vitro Antioxidant Potentiality of Extracts of Ixora chinensis Lam Leaves", Asian Journal of Chemistry, 2016  
Publication <1 %
- 
- 36 [bcc.bas.bg](http://bcc.bas.bg)  
Internet Source <1 %
- 
- 37 [spiral.imperial.ac.uk](http://spiral.imperial.ac.uk)  
Internet Source <1 %
- 
- 38 Habibi, Maryam, Mohammad-Taghi Golmakani, Asgar Farahnaky, Gholamreza Mesbahi, and Mahsa Majzooobi. "NaOH-free debittering of table olives using power ultrasound", Food Chemistry, 2016.  
Publication <1 %
- 
- 39 Submitted to Higher Education Commission Pakistan  
Student Paper <1 %
- 
- 40 [sciencepubco.com](http://sciencepubco.com)  
Internet Source <1 %
- 
- 41 [vestnik-bees.kemsu.ru](http://vestnik-bees.kemsu.ru)  
Internet Source <1 %
- 
- 42 Ali Ghasemzadeh, Hawa Z. E. Jaafar, Mohamad Fhaizal Mohamad Bukhori, Mohd <1 %

Hafizad Rahmat, Asmah Rahmat.  
"Assessment and comparison of  
phytochemical constituents and biological  
activities of bitter bean (*Parkia speciosa*  
Hassk.) collected from different locations in  
Malaysia", *Chemistry Central Journal*, 2018

Publication

---

43

I.M.C. Brighente, M. Dias, L.G. Verdi, M.G.  
Pizzolatti. "Antioxidant Activity and Total  
Phenolic Content of Some Brazilian Species",  
*Pharmaceutical Biology*, 2008

<1 %

Publication

---

44

Meihua Chen, Xu He, Hui Sun, Yue Sun, Li Li,  
Junyi Zhu, Guangqing Xia, Xin Guo, Hao Zang.  
"Phytochemical analysis, UPLC-ESI-Orbitrap-  
MS analysis, biological activity, and toxicity of  
extracts from *Tripleurospermum limosum*  
(Maxim.) Pobed", *Arabian Journal of  
Chemistry*, 2022

<1 %

Publication

---

45

Tayyaba Afsar, Suhail Razak, Ali Almajwal,  
Maria Shabbir, Khushbukhat Khan, Janeen  
Trembley, Nawaf W. Alruwaili. "Bioassay-  
guided isolation and characterization of lead  
antimicrobial compounds from *Acacia  
hydasypica* plant extract", *AMB Express*, 2022

<1 %

Publication

---

46

[file.scirp.org](http://file.scirp.org)

Internet Source

<1 %

47

[hydra.hull.ac.uk](http://hydra.hull.ac.uk)

Internet Source

<1 %

48

[researcherslinks.com](http://researcherslinks.com)

Internet Source

<1 %

49

[www.hindawi.com](http://www.hindawi.com)

Internet Source

<1 %

50

Ayana Fekadu, Yetenayet B. Tola, Addisalem Hailu Taye, Ebisa Olika Keyata. "Effect of Oromo Dinich (*Plectranthus edulis*) flour supplemented on quality characteristics of teff-maize composite injera", *Heliyon*, 2022

Publication

<1 %

51

Sechene S. Gololo, Denga Nthai, Motetelo A. Mogale. "Phytochemical Contents and Antioxidant Capacity of the Leaves of *Aloe greatheadii* var. *davyana* from Four South African Provinces", *Asian Journal of Chemistry*, 2019

Publication

<1 %

52

Shiquan Qian, Jing Zhou, Xiangyu Chen, Wei Ji, Luyao Zhang, Weicheng Hu, Zhaoxin Lu. "Evaluation of an efficient fed-batch enzymatic hydrolysis strategy to improve production of functional xylooligosaccharides

<1 %



# from maize straws", Industrial Crops and Products, 2020

Publication

---

53	<a href="http://bpkimi1.kemenperin.go.id">bpkimi1.kemenperin.go.id</a> Internet Source	<1 %
54	<a href="http://conference.cusat.ac.in">conference.cusat.ac.in</a> Internet Source	<1 %
55	<a href="http://mail.scialert.net">mail.scialert.net</a> Internet Source	<1 %
56	<a href="http://repository.up.ac.za">repository.up.ac.za</a> Internet Source	<1 %
57	<a href="http://scidar.kg.ac.rs">scidar.kg.ac.rs</a> Internet Source	<1 %
58	<a href="http://stp-mataram.e-journal.id">stp-mataram.e-journal.id</a> Internet Source	<1 %
59	<a href="http://sure.su.ac.th">sure.su.ac.th</a> Internet Source	<1 %
60	<a href="http://wjst.wu.ac.th">wjst.wu.ac.th</a> Internet Source	<1 %
61	<a href="http://www.jpsr.pharmainfo.in">www.jpsr.pharmainfo.in</a> Internet Source	<1 %
62	<a href="http://www.science.gov">www.science.gov</a> Internet Source	<1 %
63	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	<1 %

---

64

A.A. Karaçelik, H. Sahin. "DETERMINATION OF ENZYME INHIBITION AND ANTIOXIDANT ACTIVITY IN SOME CHESTNUT HONEYS", *Foods and Raw materials*, 2018

Publication

<1 %

65

Nayana Monção, Bruno Araújo, Jurandy Silva, Daisy Lima, Paulo Ferreira, Flavia Airoidi, Cláudia Pessoa, Antonia Citó. "Assessing Chemical Constituents of *Mimosa caesalpinifolia* Stem Bark: Possible Bioactive Components Accountable for the Cytotoxic Effect of *M. caesalpinifolia* on Human Tumour Cell Lines", *Molecules*, 2015

Publication

<1 %

66

F. C. Pontes, V. C. P. Abdalla, M. Imatomi, L. F. G. Fuentes, S. C. J. Gualtieri. "Antifungal and antioxidant activities of mature leaves of *Myrcia splendens* (Sw.) DC.", *Brazilian Journal of Biology*, 2018

Publication

<1 %

67

M. Arivalagan, T.K. Roy, A.M. Yasmeeen, K.C. Pavithra, P.N. Jwala, K.S. Shivasankara, M.R. Manikantan, K.B. Hebbar, S.R. Kanade. "Extraction of phenolic compounds with antioxidant potential from coconut (*Cocos nucifera* L.) testa and identification of phenolic acids and flavonoids using UPLC coupled with TQD-MS/MS", *LWT*, 2018

<1 %

68

Van Hoang Lan, Le Thi Phuong Hoa.  
"SECONDARY METABOLITES AND IN VITRO  
BIOLOGICAL ACTIVITIES OF FRACTIONS FROM  
Stemona collinsae Craib ROOT EXTRACT", BÁO  
CÁO KHOA HỌC VỀ NGHIÊN CỨU VÀ GIẢNG  
DẠY SINH HỌC Ở VIỆT NAM - PROCEEDING  
OF THE 4TH NATIONAL SCIENTIFIC  
CONFERENCE ON BIOLOGICAL RESEARCH  
AND TEACHING IN VIETNAM, 2020

Publication

<1 %

69

[jwoodscience.springeropen.com](http://jwoodscience.springeropen.com)  
Internet Source

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On