

On total edge irregularity strength of tadpole chain graph $Tr(6, n)$

by Mulyono Matematika

Submission date: 31-Aug-2022 08:18AM (UTC+0700)

Submission ID: 1889724063

File name: Nurdini_2020_J._Phys._Conf._Ser._1538_012001_dg_Bu_Isnaini.pdf (1.21M)

Word count: 3020

Character count: 11160

PAPER · OPEN ACCESS

4

On total edge irregularity strength of tadpole chain graph $Tr(6, n)$

To cite this article: E Nurdini *et al* 2020 *J. Phys.: Conf. Ser.* **1538** 012001

View the [article online](#) for updates and enhancements.



IOP ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

On total edge irregularity strength of tadpole chain graph $T_r(6, n)$

E Nurdini, I Rosyida, and Mulyono

Mathematics Department, Universitas Negeri Semarang, Semarang, Indonesia

E-mail: eka.nurdini51@gmail.com; iisnaini@gmail.com; mulyono.mat@mail.unnes.ac.id

Abstract. Given a graph $G(V, E)$ with a non-empty set of vertices V and a set of edges E . A total labelling $f: V \cup E \rightarrow \{1, 2, \dots, k\}$ is called an edge irregular total labeling if the weight of every edge is distinct. The weight of an edge e , under the total labeling f , is the sum of label of edge and all labels of vertices that are incident to e . In other words, $wt(xy) = f(xy) + f(x) + f(y)$. The total edge irregularity strength of G , denoted by $tes(G)$ is the minimum k used to label graph G with the edge irregular total labeling. A tadpole chain graph of length r , denoted as $T_r(6, n)$, is a chain graph that consists of tadpole graph $T(6, n)$ on each block. In this paper, we get $tes(T_r(6, n)) = \left\lceil \frac{(6+n)r+2}{3} \right\rceil$ and construct an algorithm to find it.

1. Introduction

Given a simple, connected and undirected graph $G = (V(G), E(G))$. A labelling of G is a function that assigns a set of elements of G into a set of positive integers [14]. A labelling f on G is said to be a total labeling if its domain is union $V(G) \cup E(G)$. Bača et al. [3] defined an edge irregular total k -labelling as a function $f: V(G) \cup E(G) \rightarrow \{1, 2, \dots, k\}$ which has the weights $wt(uv) \neq wt(xy)$ for every two different edges uv and xy , where $wt(e) = wt(uv) = f(u) + f(v) + f(e)$. Further, a total edge irregularity strength of G , symbolized by $tes(G)$, is a minimum number k in edge irregular total k -labelling.

The bounds for tes of any graph G was given by Bača et al. [3] as the following:

$$\left\lceil \frac{|E(G)|+2}{3} \right\rceil \leq tes(G) \leq |E|. \quad (1)$$

Meanwhile, Ivančo and Jendrol [8] found a conjecture for tes of graph G :

$$tes(G) = \max \left\{ \left\lceil \frac{|E(G)|+2}{3} \right\rceil, \left\lceil \frac{\Delta(G)+1}{2} \right\rceil \right\} \quad (2)$$

where $\Delta(G)$ is a maximum degree of all vertices of G .

The proof of Conjecture (2) has been revealed by some researchers for some special graphs, such as: Jendrol et al. [9] verified tes of complete and complete bipartite graphs; Ivančo and Jendrol [8] gave tes of any tree. Furthermore, tes of some graph classes has been investigated by many researchers as well as presented in Gallian [4]. Mushayt and Ahmad investigated tes of hexagonal grid graphs [10]. Indriati et al. ([6],[7]) found tes of generalized helm and generalized web graphs. Nurdini and Rosyida [11] found tes of dovetail graph with some pendant vertices and related graph. Rosyida and Indriati [13] provided tes of C_3 and C_4 cactus chain graphs with pendant vertices. The readers can find more results on es , tes , and tvs of graphs in [5].

The authors were encouraged by the results in Rosyida et al. [12] that determined tvs of $T_r(4, 1)$ tadpole chain graph. The authors were also interested to the result in [5] that gave an edge irregularity



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

strength (es) of cycle chain graphs and results from Ahmad et al. [1] which proposed es of several chain graphs. The problem investigated in this paper is different to the result in [12], [5] and [1]. We verify tes of tadpole chain graph $T_r(6, n)$ and construct an algorithm to find it.

2. Main Results

In the following investigation, we discuss an exact value of the total edge irregularity strength of $T_r(6, n)$ tadpole chain graph as presented in Theorem 2.1. We refer the concept of $T_r(6, n)$ tadpole chain graph from [2] and [12].

Definition 1 A tadpole graph $T_{(k,n)}$, is the graph created by concatenating an edge from any vertex of C_k with a pendant of P_n for integers $k \geq 3$ and $n \geq 1$. The tadpole graph contains $m + n$ vertices and $m + n$ edges.

Definition 2 Given a connected graph $G(V, E)$. A block cut vertex graph of G is a graph in which the vertices are the blocks and cut vertices of G . A chain graph is a graph which contains some blocks B_1, B_2, \dots, B_r so that each pair of block B_i, B_{i+1} has at most one common cut vertex such that the block cut vertex is a path. A chain graph which each block is tadpole graph is called tadpole chain graph.

In a $T_r(6, n)$ tadpole chain graph, each hexagon has cut vertices at most two, each of two hexagons has one common cut vertex, and a path P_n concatenated in each hexagon. The length of the chain is indicated by the number r on $T_r(6, n)$ tadpole chain graph. The notation $T_r(6, n)$ stands for a $T_r(6, n)$ tadpole chain graph with length r . The formula for tes of $T_r(6, n)$ is presented in Theorem 2.3.

Theorem 1 Given a tadpole chain graph $T_r(6, n)$ with length r and P_n concatenated in each hexagon. The total edge irregularity strength of $T_r(6, n)$ is

$$tes(T_r(6, n)) = \left\lceil \frac{(6+n)r+2}{3} \right\rceil.$$

Proof. Tadpole chain graph $T_r(6, n)$ consists of $(6+n)r$ edges. Let $u_{2i-1}, u_{2i}, v_i, x_{2i-1}, x_{2i}$ be vertices located on each hexagon. Let u_{2i-1}, u_{2i} be the two vertices on the top of hexagon for $i = 1, 2, \dots, 2r$, let v_i be the cut vertices for $i = 1, 2, \dots, r$, and x_{2i-1}, x_{2i} be vertices located on the bottom of hexagon for $i = 1, 2, \dots, 2r$. Let x_{2i-1} be the vertex that concatenated with y_i^i which is the part of y_i^j for $i = 1, 2, \dots, r$ and $j = 1, 2, \dots, n$. The lower bound for tes of the graph $T_r(6, n)$ is as follows [3]:

$$\left\lceil \frac{(6+n)r+2}{3} \right\rceil \leq tes(T_r(6, n)) \leq (6+n)r.$$

Further, we show the upper bound of $tes(T_r(6, n)) \leq \left\lceil \frac{(6+n)r+2}{3} \right\rceil$ by constructing a total k -labeling $f: V \cup E \rightarrow \{1, 2, \dots, k\}$ where $k = \left\lceil \frac{(6+n)r+2}{3} \right\rceil$ as follows.

Case 1. For $n = 3 \pmod 3$:

Labels of vertices are defined as the following:

$$\begin{aligned} f(U_{2i-1}) &= f(U_{2i}) = \left\lceil \frac{(6+n)i+2}{3} \right\rceil, i = 1, 2, \dots, r \\ f(X_{2i-1}) &= f(X_{2i}) = \left\lceil \frac{(6+n)i+2}{3} \right\rceil - 1, i = 1, 2, \dots, r \\ f(V_i) &= \left\lceil \frac{(6+n)i+2}{3} \right\rceil - 3, i = 1, 2, \dots, r \end{aligned}$$

$$f(Y_i^j) = \frac{(6+n)i - n + \left(3 \left\lfloor \frac{i}{3} \right\rfloor - 6\right)}{3}, i = 1, 2, \dots, r; j = 1, 2, \dots, n$$

Meanwhile, labels of edges are:

$$f(U_{2i-1}U_{2i}) = \left\lfloor \frac{(6+n)i + 2}{3} \right\rfloor - 3, i = 1, 2, \dots, r$$

$$f(U_{2i-1}V_i) = f(V_iX_{2i-1}) = f(X_{2i-1}X_{2i}) = \left\lfloor \frac{(6+n)i + 2}{3} \right\rfloor - 2, i = 1, 2, \dots, r$$

$$f(U_{2i-2}V_i) = f(V_iX_{2i-2}) = \frac{(6+n)i - n + 3}{3}, i = 1, 2, \dots, r$$

$$f(Y_i^j Y_i^{j+1}) = \frac{(6+n)i - n + \left(3 \left\lfloor \frac{j+2}{3} \right\rfloor - 6\right)}{3}, i = 1, 2, \dots, r; j = 1, 2, \dots, n$$

$$f(X_{2i-1}Y_i^n) = \left\lfloor \frac{(6+n)i - 8}{3} \right\rfloor - 3, i = 1, 2, \dots, r$$

Case 2. For $n \neq 3 \pmod 3$:

Labels of vertices are defined as follows:

$$f(U_{2i-1}) = f(U_{2i}) = \begin{cases} 3, & \text{if } i = 1; n = 2 \\ \left\lfloor \frac{(6+n)i + 2}{3} \right\rfloor, & i = 1, 2, \dots, r \end{cases}$$

$$f(X_{2i-1}) = \begin{cases} 1, & \text{if } i = 1; n = 1 \\ 2, & \text{if } i = 1; n = 2 \\ \left\lfloor \frac{(6+n)i + 2}{3} \right\rfloor - 1, & i = 1, 2, \dots, r \end{cases}$$

$$f(X_{2i}) = \begin{cases} 2, & \text{if } i = 1; n = 2 \\ \left\lfloor \frac{(6+n)i + 2}{3} \right\rfloor - 1, & i = 1, 2, \dots, r \end{cases}$$

$$f(V_i) = \begin{cases} 1, & \text{if } i = 1 \text{ and } n = 1 \\ \left\lfloor \frac{(6+n)i + 2}{3} \right\rfloor - 3, & i = 1, 2, \dots, r \end{cases}$$

$$f(Y_i^j) = \left\lfloor \frac{(6+n)i - n + j - 6}{3} \right\rfloor, i = 1, 2, \dots, r; j = 1, 2, \dots, n$$

Meanwhile, labels of edges are:

$$f(U_{2i-1}U_{2i}) = \begin{cases} 2, & \text{if } i = 1, n = 2 \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 4, & \text{if } ((6+n)i) \bmod 3 = 2, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 2, & \text{if } ((6+n)i) \bmod 3 = 1, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 3, & \text{if } ((6+n)i) \bmod 3 = 0, i = 1, 2, \dots, r \end{cases}$$

$$f(U_{2i-1}V_i) = \begin{cases} 1, & \text{if } i = 1, n = 1 \\ 2, & \text{if } i = 1, n = 2 \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 3, & \text{if } ((6+n)i) \bmod 3 = 2, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 1, & \text{if } ((6+n)i) \bmod 3 = 1, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 2, & \text{if } ((6+n)i) \bmod 3 = 0, i = 1, 2, \dots, r \end{cases}$$

$$f(V_iX_{2i-1}) = \begin{cases} \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 3, & \text{if } ((6+n)i) \bmod 3 = 2, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 1, & \text{if } ((6+n)i) \bmod 3 = 1, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 2, & \text{if } ((6+n)i) \bmod 3 = 0, i = 1, 2, \dots, r \\ 3, & \text{if } i = 1, n = 1, 2 \end{cases}$$

$$f(X_{2i-1}X_{2i}) = \begin{cases} \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 3, & \text{if } ((6+n)i) \bmod 3 = 2, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 1, & \text{if } ((6+n)i) \bmod 3 = 1, i = 1, 2, \dots, r \\ \left\lfloor \frac{(6+n)i+2}{3} \right\rfloor - 2, & \text{if } ((6+n)i) \bmod 3 = 0, i = 1, 2, \dots, r \end{cases}$$

$$f(U_{2i-2}V_i) = f(V_iX_{2i-2}) = \begin{cases} 4, & \text{if } i = 1 \text{ and } n = 2 \\ \left\lfloor \frac{(6+n)i - (n-2)}{3} \right\rfloor, & i = 1, 2, \dots, r \end{cases}$$

$$f(Y_i^j Y_i^{j+1}) = \left\lfloor \frac{(6+n)i - n + j - 4}{3} \right\rfloor, i = 1, 2, \dots, r; j = 1, 2, \dots, n$$

$$f(X_{2i-1}Y_i^n) = \begin{cases} 1, & i = 1, n = 1, 2 \\ \left\lfloor \frac{(6+n)i - 8}{3} \right\rfloor, & i = 1, 2, \dots, r \end{cases}$$

Since we get the labels of vertices and edges are less than or equal to $k = \left\lfloor \frac{(6+n)r+2}{3} \right\rfloor$, then the labeling f is ak -total labeling. [8](#)

Further, we verify that the weights of edges are distinct under the function f as follows:

$$wt(U_{2i-1}U_{2i}) = (6+n)i, i = 1, 2, \dots, r$$

$$wt(U_{2i-1}V_i) = (6+n)i - 2, i = 1, 2, \dots, r$$

$$wt(V_iX_{2i-1}) = (6+n)i - 3, i = 1, 2, \dots, r$$

$$wt(X_{2i-1}X_{2i}) = (6+n)i - 1, i = 1, 2, \dots, r$$

$$\begin{aligned} wt(U_{2i-2}V_i) &= (6+n)i+2, i=1,2,\dots,r \\ wt(V_iX_{2i-2}) &= (6+n)i+1, i=1,2,\dots,r \\ wt(Y_i^j Y_i^{j+1}) &= (6+n)i-n+j-4, i=1,2,\dots,r, j=1,2,\dots,n \\ wt(X_{2i-1}Y_i^n) &= wt(Y_i^{n-1}Y_i^n) + 1, i=1,2,\dots,r \end{aligned}$$

It is clear that the weights of all edges are distinct and we obtain upper bound $tes(T_r(6,n)) \leq \left\lceil \frac{(6+n)r+2}{3} \right\rceil$. Thus, we show that tes of tadpole chain graph $T_r(6,n)$ as follows:

$$tes(T_r(6,n)) = \left\lceil \frac{(6+n)r+2}{3} \right\rceil.$$

3. Computational results

In this section, we present computational result of tes of $T_r(6,n)$ graph. A computer program by using Matlab R2016a is constructed based on an algorithm in Table 1.

Table 1. Algorithm to determine tes of $T_r(6,n)$ tadpole chain graph.

```

Commands:
1  input r           %Length of chain graph
2  input n           %Number of vertices in path Pn
3  for i=1 to r      % assign labels to vertices of G
4       $f(U_{2i-1}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right); f(U_{2i}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right)$ 
5       $f(X_{2i-1}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 1; f(X_{2i}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 1$ 
6           $f(V_i) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 3$ 
7
8      for j=1 to n
9           $f(Y_i^j) = \frac{(6+n)i - n + \left(3\text{ceil}\left(\frac{i}{3}\right) - 6\right)}{3}$ 
10     end
11 end
12 for i=1 to r      % assign labels to edges and determine the
    weights
13      $f(U_{2i-1}U_{2i}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 3;$ 
14          $wt(U_{2i-1}U_{2i}) = (6+n)i;$ 
15          $f(U_{2i-1}V_i) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 2$ 
16          $wt(U_{2i-1}V_i) = (6+n)i - 2$ 
17          $f(V_iX_{2i-1}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 2$ 
18          $wt(V_iX_{2i-1}) = (6+n)i - 3$ 
19          $f(X_{2i-1}X_{2i}) = \text{ceil}\left(\frac{(6+n)i+2}{3}\right) - 2$ 
         $wt(X_{2i-1}X_{2i}) = (6+n)i - 1$ 

```

Commands :

```

20       $f(U_{2i}V_{i+1}) = \frac{(6+n)i-n+3}{3}$ 
21       $wt(U_{2i}V_{i+1}) = (6+n)i + 2,$ 
22       $f(V_{i+1}X_{2i}) = \frac{(6+n)i-n+3}{3}$ 
23       $wt(V_{i+1}X_{2i}) = (6+n)i + 1,$ 
24       $f(X_{2i-1}Y_i^n) = \text{ceil}\left(\frac{(6+n)i-8}{3}\right) - 3$ 
25      for j=1 to n-1
26       $(Y_i^j Y_i^{j+1}) = \frac{(6+n)i-n + (3\text{ceil}(\frac{j+2}{3}) - 6)}{3}$ 
27       $wt(Y_i^j Y_i^{j+1}) = (6+n)i - n + j - 4$ 
28      end
29      end
30      Print [f(edges) wt(edges)]%display edges and the weighs
    
```

As a simulation, we give an illustration of the edge irregular total 13-labeling of $T_4(6,3)$ in Figure 1. The weight of each edge is printed in the red color. By using the algorithm, the labeling output and tes of $T_r(6,3)$ from computer program is given in Figure 2.




Figure 1.The total 13-labeling of $T_4(6,3)$.

Matlab output for determining *tes* of $T_4(6,3)$ is presented in Figure 2.

Label_u(2i)	Label_x(2i)	Label_y1	Label_y2	Label_y3	Label_y4	Label_v
0	0					1
4	3	1	4	7	10	4
0	0					7
7	6	1	4	7	10	10
0	0					13
10	9	1	4	7	10	
0	0					
13	12					

Edges_u(2i-1) u(2i)	Label	Weight	Edges_u(2i-1) v(i)	Label	Weight
1 2	1	9	1 1	2	7
3 4	4	18	3 2	5	16
5 6	7	27	5 3	8	25
7 8	10	36	7 4	11	34

Edges_x(2i-1) x(2i)	Label	Weight	Edges_x(2i) v(i)	Label	Weight
1 2	2	8	2 2	3	10
3 4	5	17	4 3	6	19
5 6	8	26	6 4	9	28
7 8	11	35	8 5	12	37

Edges_x(2i-1) y(i)	Label	Weight	Edges_y1_y2	Label	Weight
1 1	1	5	1 2	1	3
3 2	4	14	2 3	2	4
5 3	7	23			
7 4	10	32			

Edges_y5_y6	Label	Weight	Edges_y5_y6	Label	Weight
1 2	7	21	1 2	7	21
2 3	8	22	2 3	8	22

Figure 2. The labeling output of total 13-labeling of $T_4(6,3)$ by Matlab.

4. Conclusions

In this paper, we have invented and proved tes oftadpole chain graph $T_r(6, n)$. We found that $tes(T_r(6, n)) = \left\lfloor \frac{(6+n)r+2}{3} \right\rfloor$ and an algorithm to find the tes is also constructed. In upcoming work, we will investigate tes of generalized tadpole chain and generalized cactus chain graphs. Moreover, we present an open problem for further research.

Open Problems ² The total vertex irregularity strength of generalized tadpole chain and generalized cactus chain graphs

Acknowledgment

This paper is supported by Basic Research Grant, Ristekdikti, with contract number 192/SP2H/LT/BRPM2019.

References

- [1] Ahmad A, Gupta A, Simanjuntak R 2018 Computing the edge irregularity strengths of chain graphs and the join of two graphs *Electronic Journal of Graph Theory and Applications* **6**(1) 201-07
- [2] Arockiamary S T 2016 Total edge irregularity strength of diamond snake and dove *IJPAM* **109** 125-132
- [3] Bača M, Jendrol S, Miller M and Ryan J 2007 On irregular total labeling *Discrete Math* **307** 1378-1388
- [4] Gallian J A 2017 A dynamic survey of graph labeling *The Electronic Journal of Combinatorics* **19** #DS6
- [5] Imran M, Aslam A, Zafar S and Nazeer W 2017 Further results on edge irregularity strength of graphs *Indonesian Journal of Combinatorics* **1**(2) 82-91
- [6] Indriati D, Widodo, Wijayanti I E and Sugeng K A 2013 On the total edge irregularity strength of generalized helm *AKCE International Journal of Graphs and Combinatorics* **10**(2) 147-55
- [7] Indriati D, Widodo, Wijayanti I E, Sugeng K A and Baca M 2015 On total edge irregularity strength of generalized web graphs and related graphs *Mathematics in Computer Science* **9** 161-67
- [8] Ivančo J and Jendrol S 2006 Total edge irregularity strength of trees *Discussiones Math Graph Theory* **26** 449-456
- [9] Jendrol S, Miskuf J and Sotak R 2010 Total edge irregularity strength of complete graphs and complete bipartite graphs *Discrete Mathematics* **310**(3) 400-07
- [10] Mushayt O A and Ahmad A 2012 On the total edge irregularity strength of hexagonal grid graphs *Australasian Journal of Combinatorics* **53** 263–271
- [11] Nurdini E, Rosyida I 2018 On total edge irregularity strength of dove tail graph with pendant vertices and its subdivision *J of Phys: Conference Series* **1217** 012064
- [12] Rosyida I, Mulyono and Indriati D 2019 Determining total vertex irregularity strength of $T_r(4, 1)$ tadpole chain graph and its computation *Procedia Computer Science* **157** 699–706
- [13] Rosyida I, Indriati D 2018 On total edge irregularity strength of some cactus chain graphs with pendant vertices *J of Phys: Conference Series* **1211** 012016
- [14] Wallis W D 2001 *Magic Graphs* Birkhauser Boston

On total edge irregularity strength of tadpole chain graph $Tr(6, n)$

ORIGINALITY REPORT

10%

SIMILARITY INDEX

8%

INTERNET SOURCES

6%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	earchive.tpu.ru Internet Source	2%
2	www.ejgta.org Internet Source	2%
3	fs.unm.edu Internet Source	1%
4	F. Salama. "On total edge irregularity strength of polar grid graph", Journal of Taibah University for Science, 2019 Publication	1%
5	Béla Bollobás. "Chapter 3 Flows, Connectivity and Matching", Springer Science and Business Media LLC, 1979 Publication	1%
6	Ika H. Agustin, Liliek Susilowati, Dafik, Ismail N. Cangul, N. Mohanapriya. "On the vertex irregular reflexive labeling of several regular and regular-like graphs", Journal of Discrete	1%

Mathematical Sciences and Cryptography, 2022

Publication

7	Nurdin Hinding, Dian Firmayasari, Hasmawati Basir, Martin Bača, Andrea Semaničová-Feňovčíková. "On irregularity strength of diamond network", AKCE International Journal of Graphs and Combinatorics, 2017 Publication	1 %
8	repository.unhas.ac.id Internet Source	1 %
9	Submitted to Higher Education Commission Pakistan Student Paper	1 %
10	Muhammad Kamran Siddiqui, Deeba Afzal, Muhammad Ramzan Faisal. "Total edge irregularity strength of accordion graphs", Journal of Combinatorial Optimization, 2016 Publication	<1 %
11	Submitted to University of Cambridge Student Paper	<1 %
12	ijmsi.ir Internet Source	<1 %
13	acikerisim.isikun.edu.tr Internet Source	<1 %
14	archive.org Internet Source	<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On

On total edge irregularity strength of tadpole chain graph $Tr(6, n)$

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
