

The total irregularity strength of caterpillars with odd number of internal vertices of degree three

by Mulyono Matematika

Submission date: 31-Aug-2022 10:59AM (UTC+0700)

Submission ID: 1889819482

File name: pillars_with_odd_number_of_internal_vertices_of_degree_three.pdf (810.97K)

Word count: 3055

Character count: 12760

PAPER · OPEN ACCESS

The total irregularity strength of caterpillars with odd number of internal vertices of degree three

To cite this article: M R Maulana *et al* 2021 *J. Phys.: Conf. Ser.* **1918** 042034

View the [article online](#) for updates and enhancements.



IOP ebooksTM

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

5

The total irregularity strength of caterpillars with odd number of internal vertices of degree three

M R Maulana, I Rosyida* and Mulyono

Department of Mathematics, Faculty of Mathematics and Natural Sciences,
Universitas Negeri Semarang, Indonesia.

*Corresponding author: isnaini@mail.unnes.ac.id

Abstract. Given a graph G consisting of vertex set V and edge set E , respectively. Assume G is simple, connected, and the edges do not have direction. A function λ that maps $V \cup E$ into a set of k -integers is named a totally irregular total k -labelling if no vertices have the same weight and also the edges of G get distinct weights. We call the minimum number k for which G has totally irregular total k -labelling as total irregularity strength of G , $ts(G)$. In this article, we construct labels of vertices and edges of caterpillar graphs which have q internal vertices of degree 3 where q is 5, 7, and 9. We obtain the exact values of ts in the following: $n + 2$ if the caterpillars have $q=5$ internal vertices, $n + 3$ for $q=7$, and $n + 4$ for $q=9$.

1. Introduction

The notion of graph labelling was introduced in Alexander Rosa's paper in 1967. The definition is as follows: "graph labelling is a function that has elements of $G(V, E)$ as its domain and a set of positive integers as its co-domain. If the function assigns $V(G)$ to the co-domain, then it is called a *vertex labelling*. Meanwhile, if the function maps $E(G)$ to the set of positive integers, then it is named an *edge labelling*. Moreover, the function is mentioned as a *total labelling* when it has $V \cup E$ as its domain" [1].

Further, the definition of *edge irregular total labelling* was given in Baca et al. as follows: "a total k -labelling $f: V \cup E \rightarrow \{1, 2, \dots, k\}$ is named as an *edge irregular total k -labelling* when $w(e_1) \neq w(e_2)$ for each different pair of edges e_1, e_2 in $E(G)$. If $e = xy$, then the weight $w(e) = f(x) + f(y) + f(xy)$. The minimum number k in such a way G has such labelling as a total edge irregularity strength of G , indicated by $tes(G)$ " [2]. Whereas, the tes of a tree $T(V, E)$ with a maximum degree $\Delta(T)$ was proved in [3]:

$$tes(T) = \max \left\{ \left\lceil \frac{|E(T)| + 2}{3} \right\rceil, \left\lceil \frac{\Delta(T) + 1}{2} \right\rceil \right\} \quad (1)$$

Note that the symbol $\lceil x \rceil$ means the least integer greater than or equal to x .

The definition of *vertex irregular total k -labelling* was also given in Baca et al. as follows: "a total k -labelling $f: V \cup E \rightarrow \{1, 2, \dots, k\}$ is named as an *vertex irregular total k -labelling* if the weights $w(v_1) \neq w(v_2)$ whenever $v_1 \neq v_2$ in $V(G)$, where $w(v) = f(v) + \sum_{v \in E} f(vy)$. We call the minimum positive integer k for which G has such labelling as a total vertex irregularity strength of G , symbolized



by $tv_s(G)$ " [1]. The tv_s of a tree $T(V, E)$ that consists of n pendant vertices and it does not have vertices of degree 2 was proved in [4]:

$$tv_s(T) = \left\lceil \frac{n+1}{2} \right\rceil \tag{2}$$

Furthermore, the notion of *totally irregular total labelling* was initiated in [5]: "the function f is called out as a totally irregular total k -labelling if both of the vertex-weights and the edge-weights are distinct. The minimum integer k such that G has totally irregular total k -labelling is named as total irregularity strength of G , denoted by $ts(G)$ ". The lower bound for ts of any graph was also given in [5]:

$$ts(G) \geq \max\{tes(G), tv_s(G)\} \tag{3}$$

Some researchers have found tes and tv_s of any graph classes as in [6], [7], [8-13], and [14]. Whereas, several exact values of ts of any graph have also been found, such as in [14-18]. A *caterpillar* graph is a *tree* such that removing its pendant edges will form a path [19]. The exact values of ts of some classes of caterpillars are still unknown [15]. Therefore, we investigate ts of caterpillar graphs where the number of internal vertices of degree 3 is odd.

2. Methods

According to the method in [2],[15-18], we summarize the steps used in this paper as follows:

- a. Defining caterpillars which have q internal vertices of degree 3 where $q = 5,7,9$. The caterpillar is denoted by $S_{n, \underbrace{3,3, \dots, 3}_q, n}$.
- b. Calculating tes of the caterpillars based on eq.(1).
- c. Calculating tv_s of the caterpillars based on eq.(2).
- d. Determining lower bound of ts of $G = S_{n, \underbrace{3,3, \dots, 3}_q, n}$ according to inequality (3).
- e. Setting the lower bound $k = \max\{tes(G), tv_s(G)\}$ where $tes(G)$ is given in (1) and $tv_s(G)$ is provided in (2), respectively.
- f. Proving that ts of the caterpillars is least than or equal to k by constructing a totally irregular total k -labelling f (trial and error process) on the caterpillars and the process is continued until we obtain a fixed pattern of the labelling;
- g. Formulating vertex and edge-labels and formulating the ts of the caterpillars.
- h. Proving the formula of the weights and showing that the weights are different.
- i. Obtaining the $ts = k$.

3. Results and Discussion

We present the results of ts of caterpillar graphs having odd number of internal vertices of degree 3.

3.1. Caterpillar graphs with 5 internal vertices of degree 3

We present the concept of the caterpillars in Definition 3.1.1.

Definition 3.1.1. Caterpillar $S_{n, \underbrace{3,3, \dots, 3}_5, n}$ is tree which is formed from double star $S_{n,n}$ by inserting five vertices (v^2, v^3, v^4, v^5, v^6) on the bridge that connects the centrals of the stars (v^1 and v^7) and the five vertices are incident to pendant edges ($v^j | 2 \leq j \leq 6$). Whereas, the vertices of the double stars are $\{v_i^1: 1 \leq i \leq n - 1\}$ and $\{v_i^7: 1 \leq i \leq n - 1\}$. The caterpillar has $2n + 10$ vertices, $2n + 9$ edges, and $2n + 3$ pendant vertices. It has maximum degree is $\Delta = n$.

An illustration of caterpillar *caterpillar* with 5 internal vertices of degree 3 is presented in Figure 1.




Figure 1. Caterpillar graph $S_{n,3,3,\dots,3,n}$

The ts of the caterpillars as in Definition 3.1.1 is presented in Theorem 3.1.1.

Theorem 3.1.1. If $S_{n,3,3,\dots,3,n}, n \geq 5$ is the caterpillar as in Definition 3.1.1, then

$$ts(S_{n,3,3,\dots,3,n}) = \left\lfloor \frac{2n+4}{2} \right\rfloor = n+2.$$

Proof. Based on eq. (1):

$$tes(S_{n,3,3,\dots,3,n}) = \max \left\{ \left\lfloor \frac{|E|+2}{3} \right\rfloor, \left\lfloor \frac{d+1}{2} \right\rfloor \right\} = \max \left\{ \left\lfloor \frac{2n+9+2}{3} \right\rfloor, \left\lfloor \frac{n+1}{2} \right\rfloor \right\} = \max \left\{ \left\lfloor \frac{2n+11}{3} \right\rfloor, \left\lfloor \frac{n+1}{2} \right\rfloor \right\} = \left\lfloor \frac{2n+11}{3} \right\rfloor$$

Further, based on eq. (2): $tvs(S_{n,3,3,\dots,3,n}) = \left\lfloor \frac{n+1}{2} \right\rfloor = \left\lfloor \frac{2n+3+1}{2} \right\rfloor = \left\lfloor \frac{2n+4}{2} \right\rfloor.$

The lower bound of ts is obtained from eq. (3):

$$ts(S_{n,3,3,\dots,3,n}) \geq \max \left\{ tes(S_{n,3,3,\dots,3,n}), tvs(S_{n,3,3,\dots,3,n}) \right\} = \max \left\{ \left\lfloor \frac{2n+11}{3} \right\rfloor, \left\lfloor \frac{2n+4}{2} \right\rfloor \right\} = \left\lfloor \frac{2n+4}{2} \right\rfloor = n+2, n \geq 5.$$

Let $k = \left\lfloor \frac{2n+4}{2} \right\rfloor$. We will prove that $ts(S_{n,3,3,\dots,3,n}) \leq k$ by constructing a totally irregular total k -labelling $h: V \cup E \rightarrow \{1, 2, \dots, k\}$ with $k = \left\lfloor \frac{2n+4}{2} \right\rfloor = n+2$. Meanwhile, vertex and edges-labels are given in Table 1.

Table 1. Labels of elements of the caterpillars with 5 internal vertices

	Vertex-labels	Edge-labels
$h(v_i^1)$	1, $i = 1$ $i - 1, \forall 2 \leq i \leq n - 1$	$h(v^1 v_i^1)$ 1, $i = 1$ 2, $\forall 2 \leq i \leq n - 1$
$h(v_i^7)$	$\left\lfloor \frac{2n+4}{2} \right\rfloor - 3, \forall 1 \leq i \leq n - 4$ $i + 2, \forall n - 3 \leq i \leq n - 1$	$h(v^7 v_i^7)$ $i + 1, \forall 1 \leq i \leq n - 4$ $\left\lfloor \frac{2n+4}{2} \right\rfloor - 4, \forall n - 3 \leq i \leq n - 1$
$h(v^j)$	1, $j = 1, 7$ $\left\lfloor \frac{2n+4}{2} \right\rfloor + (j - 6), \forall 2 \leq j \leq 6$	$h(v^j v^{j+1})$ $\left\lfloor \frac{2n+4}{2} \right\rfloor, \forall 1 \leq j \leq 5$ $\left\lfloor \frac{2n+4}{2} \right\rfloor - 3, j = 6$
$h(v_1^j)$	$\left\lfloor \frac{2n+4}{2} \right\rfloor, \forall 2 \leq j \leq 6$	$h(v^j v_1^j)$ $\left\lfloor \frac{2n+4}{2} \right\rfloor + (j - 6), \forall 2 \leq j \leq 6$

We calculate the weights of elements of the caterpillars in Table 2.

Table 2. Weights of elements of the caterpillars with 5 internal vertices

Weights of vertices		Weights of edges	
$\forall 1 \leq i \leq n - 1$		$\forall 1 \leq i \leq n - 1$	
$w(v^j)$	$3n$, for $j = 1$	$w(v^j v^{j+1})$	$2n + 1$, for $j = 1$
	$4n + (2j - 4)$, $\forall 2 \leq j \leq 5$		$3n + (2j - 5)$, $\forall 2 \leq j \leq 5$
	$4n + (j - 1)$, for $j = 6$		$2n + 2$, for $j = 6$
$w(v_i^j)$	$\frac{n(n+3)}{2} - 4$, for $j = 7$	$w(v^j v_i^j)$	$3n + (2j - 6)$, $\forall 2 \leq j \leq 6$
	$1 + i$, $j = 1$		$2 + i$, for $j = 1$
	$n + i$, $j = 7$		$1 + n + i$, for $j = 7$
$w(v_i^j)$	$2n + (j - 2)$, $\forall 2 \leq j \leq 6$		

It is shown that the labels of elements of the caterpillars are not more than $k = \lfloor \frac{2n+4}{2} \rfloor$. Further, no vertices have a same weight and also all edges have distinct weights. It proves $ts(S_{n,3,3,\dots,3,n}) \leq k = \lfloor \frac{2n+4}{2} \rfloor$. Thus, $ts(S_{n,3,3,\dots,3,n}) = \lfloor \frac{2n+4}{2} \rfloor = n + 2$ for $n \geq 5$.

3.2. Caterpillar graphs with 7 internal vertices of degree 3

The notion of a caterpillar with 7 internal vertices of degree 3 is presented in Definition 3.2.1.

Definition 3.2.1. Caterpillar graphs $(S_{n,3,3,\dots,3,n})$ are obtained from double star $S_{n,n}$ by inserting seven vertices $(v^2, v^3, v^4, v^5, v^6, v^7, v^8)$ on the bridge between the two centrals $(v^1$ and $v^9)$ and the seven vertices are incident to pendant edges $(v_i^j | 2 \leq j \leq 8)$. Whereas, the vertices of the double stars are $\{v_i^1 : 1 \leq i \leq n - 1\}$ and $\{v_i^9 : 1 \leq i \leq n - 1\}$. The caterpillar has $2n + 14$ vertices, $2n + 13$ edges, and $2n + 5$ pendant vertices. It has maximum degree is $\Delta = n$.

The ts of the caterpillars as in Definition 3.2.1 is provided in Theorem 3.2.1.

Theorem 3.2.1. If $S_{n,3,3,\dots,3,n}$, $n \geq 6$ is the caterpillar as in Definition 3.2.1, then

$$ts(S_{n,3,3,\dots,3,n}) = \lfloor \frac{2n+6}{2} \rfloor = n + 3.$$

Proof. The lower bound of ts is as follows:

$$ts(S_{n,3,3,\dots,3,n}) \geq \max \left\{ tes(S_{n,3,3,\dots,3,n}), tvs(S_{n,3,3,\dots,3,n}) \right\} = \max \left\{ \lfloor \frac{2n+15}{3} \rfloor, \lfloor \frac{2n+6}{2} \rfloor \right\} = \lfloor \frac{2n+6}{2} \rfloor = n + 3, n \geq 6.$$

Let $k = \lfloor \frac{2n+6}{2} \rfloor$. We will prove $ts(S_{n,3,3,\dots,3,n}) \leq k$ by constructing a totally irregular total k -labelling $g : V \cup E \rightarrow \{1, 2, \dots, k\}$ with $k = \lfloor \frac{2n+6}{2} \rfloor = n + 3$. Labels of elements of the caterpillar are given in Table 3.

Table 3. Labels of elements of the caterpillars with 7 internal vertices

Vertex-labels		Edge-labels	
$g(v_i^1)$	$1, i = 1$ $i - 1, \forall 2 \leq i \leq n - 1$	$g(v^1 v_i^1)$	$1, i = 1$ $2, \forall 2 \leq i \leq n - 1$

$$\begin{array}{l}
 g(v_i^9) \begin{cases} \lfloor \frac{2n+6}{2} \rfloor - 5, \forall 1 \leq i \leq n-5 \\ i+3, \forall n-4 \leq i \leq n-1 \end{cases} & g(v^9 v_i^9) \begin{cases} i+2, \forall 1 \leq i \leq n-5 \\ \lfloor \frac{2n+6}{2} \rfloor - 6, \forall n-4 \leq i \leq n-1 \end{cases} \\
 g(v^j) \begin{cases} 1, j=1,9 \\ \lfloor \frac{2n+6}{2} \rfloor + (j-8), \forall 2 \leq j \leq 8 \end{cases} & g(v^j v^{j+1}) \begin{cases} \lfloor \frac{2n+6}{2} \rfloor, \forall 1 \leq j \leq 7 \\ \lfloor \frac{2n+6}{2} \rfloor - 5, j=8 \end{cases} \\
 g(v_1^j) \begin{cases} \lfloor \frac{2n+6}{2} \rfloor, \forall 2 \leq j \leq 8 \end{cases} & g(v^j v_1^j) \begin{cases} \lfloor \frac{2n+6}{2} \rfloor + (j-8), \forall 2 \leq j \leq 8 \end{cases}
 \end{array}$$

It is obvious that the labels of elements of the caterpillar is not more than $k = \lfloor \frac{2n+6}{2} \rfloor$. Moreover, we evaluate the weights in Table 4.

Table 4. Weights of elements of the caterpillars with 7 internal vertices

Weights of vertices		Weights of edges	
	$\forall 1 \leq i \leq n-1$		$\forall 1 \leq i \leq n-1$
	$3n+1$, for $j=1$		$2n+1$, for $j=1$
$w(v^j)$	$4n+(2j-4)$, $\forall 2 \leq j \leq 7$	$w(v^j v^{j+1})$	$3n+(2j-6)$, $\forall 2 \leq j \leq 7$
	$4n+(j-1)$, for $j=8$		$2n+2$, for $j=8$
	$\frac{n(n+5)}{2} - 13$, $j=9$	$w(v^j v_1^j)$	$3n+(2j-7)$, $\forall 2 \leq j \leq 8$
$w(v_i^j)$	$1+i$, $j=1$		$2+i$, $j=1$
	$n+i$, $j=9$	$w(v^j v_i^j)$	$1+n+i$, $j=9$
$w(v_1^j)$	$2n+(j-2)$, $\forall 2 \leq j \leq 8$		

Based on the above calculation, we can see that the vertices have different weights and no edges have a same weight. Therefore, the upper bound is obtained and $ts(S_{n,3,3,\dots,3,n}) = \lfloor \frac{2n+6}{2} \rfloor = n+3$ for $n \geq 6$.

An illustration of labelling on the caterpillar $S_{n,3,3,\dots,3,n}$ is shown in Figure 2. The green colors show vertex-labels and the blue colors denote labels of edges.




Figure 2. The totally irregular total 9-labelling on the Caterpillar $S_{n,3,3,\dots,3,n}$

3.3. Caterpillar graphs with 9 internal vertices of degree 3

Definition 3.3.1. Caterpillar $S_{n,3,3,\dots,3,n}$ is a graph which is obtained from double star $S_{n,n}$ by inserting nine vertices ($v^2, v^3, v^4, v^5, v^6, v^7, v^8, v^9, v^{10}$) on the bridge connecting the two centres (v^1 and v^{11}) and the nine vertices are incident to pendant edges ($v_1^j | 2 \leq j \leq 10$). Meanwhile, the vertices of the double stars are $\{v_i^1: 1 \leq i \leq n-1\}$ and $\{v_i^{11}: 1 \leq i \leq n-1\}$. The caterpillar has $2n+18$ vertices, $2n+17$ edges, and $2n+7$ pendant vertices. Its maximum degree is $\Delta = n$.

The ts of the *caterpillars* that contain nine internal vertices of degree 3 is proved in Theorem 3.3.1.

Theorem 3.3.1. If $S_{n,3,3,\dots,3,n}$, $n \geq 7$ is the *caterpillar* as in Definition 3.3.1, then

$$ts(S_{n,3,3,\dots,3,n}) = \left\lceil \frac{2n+8}{2} \right\rceil = n+4.$$

Proof. It is similar to the proof of Theorem 3.1.1 and Theorem 3.2.1, we get a lower bound as follows:

$$ts(S_{n,3,3,\dots,3,n}) \geq \max \left\{ tes(S_{n,3,3,\dots,3,n}), tvs(S_{n,3,3,\dots,3,n}) \right\} = \max \left\{ \left\lceil \frac{2n+19}{3} \right\rceil, \left\lceil \frac{2n+8}{2} \right\rceil \right\} = \left\lceil \frac{2n+8}{2} \right\rceil = n+4, n \geq 7.$$

Let $k = \left\lceil \frac{2n+8}{2} \right\rceil$. We should show $ts(S_{n,3,3,\dots,3,n}) \leq k$ by constructing a totally irregular total k -labelling

$p: V \cup E \rightarrow \{1, 2, \dots, k\}$ with $k = \left\lceil \frac{2n+8}{2} \right\rceil = n+4$. We define labels for elements of the caterpillar to Table 5.

Table 5. Labels of vertices and edges in the caterpillars with 9 internal vertices

$p(v)$ For all $v \in V(G)$	$p(e)$ For all $e \in E(G)$
$p(v_i^1)$ 1, $i = 1$ $i - 1, \forall 2 \leq i \leq n - 1$	$p(v^1 v_i^1)$ 1, $i = 1$ 2, $\forall 2 \leq i \leq n - 1$
$p(v_i^{11})$ $\left\lceil \frac{2n+8}{2} \right\rceil - 7, \forall 1 \leq i \leq n - 6$ $i + 4, \forall n - 5 \leq i \leq n - 1$	$p(v^{11} v_i^{11})$ $i + 3, \forall 1 \leq i \leq n - 6$ $\left\lceil \frac{2n+8}{2} \right\rceil - 8, \forall n - 5 \leq i \leq n - 1$
$p(v^j)$ 1, $j = 1, 11$ $\left\lceil \frac{2n+8}{2} \right\rceil + (j - 10), \forall 2 \leq j \leq 10$	$p(v^j v^{j+1})$ $\left\lceil \frac{2n+8}{2} \right\rceil, \forall 1 \leq j \leq 9$ $\left\lceil \frac{2n+8}{2} \right\rceil - 7, j = 10$
$p(v_1^j)$ $\left\lceil \frac{2n+8}{2} \right\rceil, \forall 2 \leq j \leq 10$	$p(v^j v_1^j)$ $\left\lceil \frac{2n+8}{2} \right\rceil + (j - 10), \forall 2 \leq j \leq 10$

It is shown above that the labels of elements of the caterpillar is less than or equal to $k = \left\lceil \frac{2n+8}{2} \right\rceil$. Furthermore, we calculate the weights to Table 6.

Table 6. Weights of vertices and edges in the caterpillars with 9 internal vertices

$w(v)$ For all $v \in V(G)$	$w(e)$ For all $e \in E(G)$
$\forall 1 \leq i \leq n - 1$ $3n + 2, j = 1$ $4n + (2j - 4), \forall 2 \leq j \leq 9$ $4n + (j - 1), j = 10$ $\frac{n(n+7)}{2} - 25, j = 11$	$\forall 1 \leq i \leq n - 1$ $2n + 1, j = 1$ $3n + (2j - 7), \forall 2 \leq j \leq 9$ $2n + 2, j = 10$ $3n + (2j - 8), \forall 2 \leq j \leq 9$
$w(v_i^j)$ $1 + i, j = 1$ $n + i, j = 11$	$w(v^j v_i^j)$ $2 + i, j = 1$
$w(v_1^j)$ $2n + (j - 2), \forall 2 \leq j \leq 10$	$1 + n + i, j = 11$

We observe that all elements of the caterpillar do not have a same weight. Therefore, we get the upper bound and $ts(S_{n,3,3,\dots,3,n}) = \left\lceil \frac{2n+8}{2} \right\rceil = n+4$ for $n \geq 7$.

Figure 3 describes labelling on the *caterpillar* $S_{n,3,3,\dots,3,n}$. The green colors indicate labels of vertices and the blue colors present edge-labels.




Figure 3. The totally irregular total 11-labeling on the Caterpillar $S_{n, \underbrace{3, 3, \dots, 3}_q, n}$

4. Conclusion

In this research, we proved that ts of $(S_{n, \underbrace{3, 3, \dots, 3}_q, n})$ is equal to: $\lfloor \frac{2n+4}{2} \rfloor = n + 2$ for $q=5$, it is equal to $\lfloor \frac{2n+6}{2} \rfloor = n + 3$ for $q=6$, and it is equal to $\lfloor \frac{2n+8}{2} \rfloor = n + 4$ for $q=9$. In upcoming research, we will investigate ts of the caterpillars which have q internal vertices of degree 3 for odd $q > 9$.

References

[1] Wallis W D and Marr A M 2013 *Magic Graphs* (New York: Birkhauser-Springer)

[2] Bača M, Jendrol' S, Kathiresan K, Muthugurupackiam K, Semaničová-Fenovcikova A 2015 *Electron. Notes Discrete Math.* **48** 19

[3] Ivančo J and Jendrol' S 2006 *Discussiones Math. Graph Theory* **26** 449

[4] Nurdin, Baskoro E T, Salman A N M, and Gaos N N 2010 *Discrete Math.* **310** 3043

[5] Marzuki C C, Salman A N M and Miller M 2013 *Far East J. Math. Sci* **82**(1) 1

[6] Indriati D, Widodo, Wijayanti I E, Sugeng K A, Bača M 2015 *Math. Comput. Sci* **9** 161

[7] Indriati D, Widodo, Wijayanti I E, Sugeng K A, Bača M, Semaničová-Fenovcikova A 2016 *Australas. J. Combin.* **65** 14

[8] Rosyida I and Indriati D 2020 *Indonesian J. Comb.* **4**(1) 53

[9] Rosyida I and Indriati D 2019 *J. Phys Conf Ser.* **1211** 012016

[10] Rosyida I, Mulyono, Indriati D 2020 *J. Discret. Math. Sci. Cryptogr.* **23**(6) 1369

[11] Rosyida I, Mulyono, Indriati D 2019 *Procedia Comput. Sci.* **157** 699

[12] Rosyida I, Ningrum E, Mulyono, and Indriati D 2020 *J. Discret. Math. Sci. Cryptogr.* **23**(6) 1335

[13] Rosyida I, Ningrum E, Setyaningrum A, and Mulyono 2020 *J. Phys Conf Ser.* **1567** 022073

[14] Gallian JA 2018 *Electron. J. Combin* **1**

[15] Indriati D, Rosyida I, Widodo 2018 *J. Phys Conf Ser.* **1008** 012042

[16] Indriati D, Widodo, Rosyida I, Sugeng K A 2017 *J. Phys Conf Ser.* **855** 012018

[17] Rosyida I, Widodo, and Indriati D 2018 *J. Phys Conf Ser.* **1008** 012046

[18] Indriati D, Widodo, Wijayanti I E, Sugeng K A, Rosyida I 2020 *Electron. J. Graph Theory Appl.* **8**(2) 247

[19] Weisstein E W 2009 Caterpillar graph *MathWorld--a Wolfram Web Resource: <https://mathworld.wolfram.com/CaterpillarGraph.html>.*

The total irregularity strength of caterpillars with odd number of internal vertices of degree three

ORIGINALITY REPORT

7%

SIMILARITY INDEX

4%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Isnaini Rosyida, Mulyono, Diari Indriati. "On totally irregular total labeling of caterpillars having even number of internal vertices with degree three", AIP Publishing, 2021
Publication 2%
- 2 Diari Indriati, Isnaini Rosyida, Widodo. "On the total irregularity strength of caterpillar with each internal vertex has degree three", Journal of Physics: Conference Series, 2018
Publication 2%
- 3 E Nurdini, I Rosyida. "On total edge irregularity strength of dove tail graph with pendant vertices and its subdivision", Journal of Physics: Conference Series, 2019
Publication 1%
- 4 hdl.handle.net
Internet Source 1%
- 5 nlist.inflibnet.ac.in
Internet Source 1%

6

Internet Source

<1 %

7

Submitted to Universitas Negeri Semarang

Student Paper

<1 %

8

Submitted to Higher Education Commission
Pakistan

Student Paper

<1 %

9

ebin.pub

Internet Source

<1 %

10

repository.iainkediri.ac.id

Internet Source

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On

The total irregularity strength of caterpillars with odd number of internal vertices of degree three

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
