

# Optimization of Drying Process for Production Red Ginger Granulated Palm Sugar Using Response Surface Methodology

*by Prima Astuti Handayani*

---

**Submission date:** 07-Sep-2022 02:18PM (UTC+0700)

**Submission ID:** 1894246675

**File name:** 36124-96023-1-PB.pdf (929.57K)

**Word count:** 5062

**Character count:** 24213



## Optimization of Drying Process for Production Red Ginger Granulated Palm Sugar Using Response Surface Methodology

Prima Astuti Handayani<sup>✉</sup>, Idama Kusuma Dewi, Ady Prasetyo

DOI: <https://doi.org/10.15294/jbat.v11i1.36124>

Chemical Engineering Department, Faculty of Engineering, Semarang State University (UNNES), Indonesia

### Article Info

Article history:

Received  
April 2022  
Accepted  
May 2022  
Published  
June 2022

Keywords:

Granulated Palm  
Sugar;  
Red Ginger;  
Water Content;  
Optimum;  
Response Surface  
Methodology

### Abstract

Palm sugar is used as a natural sweetener that is added to food and beverages. The nutrient content in palm sugar can be enriched with the addition of antioxidants derived from red ginger (*Zingiber officinale var. rubrum*). The antioxidant activity of red ginger is 75.61% higher than that of empty ginger and elephant ginger. The problem that arises during the production process of granulated palm sugar is the drying process that has not considered chemical characteristics, especially water content. The requirement for water content of palm sugar according to SNI 01-3743-1995 is a maximum of 3%. Moisture content is the main parameter that determines the quality of granulated palm sugar to long shelf life. Water content can affect other chemical characteristics such as sucrose, reducing sugar, ash content, calories, protein, fat, and carbohydrates. This optimization is using RSM (Response Surface Methodology) CCD model (Central Composite Design) on Software Statistica 10 with 20 treatments. The independent variables used were time (4-6 hours), material weight (100-300 grams), and material size (10-26 mesh). Data processing with Statistica 10 software resulted in the optimum water content condition of 2.9019%, with the drying process conditions covering 6.68 hours, material weight 368.18 grams, and material size 31.45 mesh. Validation was carried out to test the accuracy of the optimization results from Statistica 10 Software. The validation of the moisture content results obtained a value of 2.9016%, with an error percentage of 0.0003%. The chemical characteristic test was applied to the granulated palm sugar as a result of optimization so that the value of sucrose (96.5967%) was obtained, reducing sugar (6.0434%), ash content (1.8660%), calories (379.93%), protein (2.4268%), fat (0.3972%), and carbohydrates (91.5379%).

### INTRODUCTION

Granulated palm sugar is one of the natural sweeteners that is rich in benefits and is consumed by the community as a food or drink sweetener (Pratama, et al., 2020). Granulated palm sugar has a lower glycemic index value than cane sugar, which is 35% (Wilberta et al., 2021). In addition, granulated palm sugar also contains several compounds useful for the body, such as calories, carbohydrates, fat, protein, and calcium (Fatriani et al., 2019). The content of beneficial compounds in granulated palm sugar can be enriched by adding antioxidants obtained from spices, one of which is

red ginger (*Zingiber officinale var. rubrum*) (Wulandari et al., 2021). Red ginger can increase the body's response to fighting viruses. Including the chikungunya virus, pharyngitis, and COVID-19 (Assegaf et al., 2020; Kaushik et al., 2020; Sinamo & Hutabarat, 2021).

Granulated palm sugar has a shelf life of 1 year, longer than jaggery sugar, lasting for two weeks. The shelf life of granulated palm sugar is influenced by the water content (Ritonga et al., 2020). The drying process is an essential part of the production of granulated palm sugar to obtain a minimum water content. Optimum drying operating conditions will produce granulated palm

<sup>✉</sup> Corresponding author:  
E-mail: [prima@mail.unnes.ac.id](mailto:prima@mail.unnes.ac.id)

sugar with minimum moisture content (Meldayanoor et al., 2019).

In general, the conventional drying process of granulated palm sugar still use the sunlight. The disadvantages of this drying method are the poor physical quality of sugar and high water content (Muhandri et al., 2020). These weaknesses can be overcome by determining the optimum operating conditions of the granulated palm sugar drying process. Parameters that are considered to produce granulated palm sugar water content according to SNI 01-3743-1995 standards are drying time, material weight, and material size using the Response Surface Methodology (RSM) method. So that, it fulfill the water content standard according to SNI 01-3743-1995, which is a maximum of 3%. The purpose of this study was to determine the effect of optimum drying operating conditions on granulated palm sugar.

## MATERIALS AND METHODS

### Materials

The materials used in this study were palm sugar obtained from the Palm Sugar Production Center Limbangan and red ginger obtained from the Karang Ayu Market, Semarang. Materials used for analysis include 30% HCl (Merck), 45% NaOH (Merck), 30% NaOH (Merck), Luff's reagent, aquades, 20% KI (Merck), Na-thio 0.1 N, H<sub>2</sub>SO<sub>4</sub> 26,5%(Merck), starch indicator, arsenomolybdate reagent, concentrated H<sub>2</sub>SO<sub>4</sub>(Merck), 2% boric acid, 0.01 N HCl (Merck), 30% HCl (Merck).

The tools used in this study were oven, thermometer, analytical scale, wooden stirrer, frying pan, stove, coconut shell, sieve, baking sheet, grater, measuring cup, filter cloth, mesh sieve, brush, and spoon. The tools used for analysis include vochdoos, 100 mL Erlenmeyer, 100 mL volumetric flask spectrophotometer (UV Vis Shimadzu), test tube, water bath, furnace (Thermolyne 1000), and a cup.

### Methods

This research was conducted in several stages, namely the production of red ginger granulated palm sugar, optimization with RSM (Response Surface Methodology) using Statistica 10 software, and testing the chemical characteristics of red ginger granulated palm sugar.

### Red Ginger Granulated Palm Sugar Production

The red ginger that was cleaned is grated using a grater to extract the juice. Red ginger juice is separated from starch deposits through a filtering process. The red ginger juice is then cooked until it boils. The boiling red ginger juice is added with palm sugar that has been cut into pieces. Cooking is done until the sugar crystallizes. The granulated palm sugar that has been formed is sieved using a sieve to obtain granulated palm sugar based on material size. The palm sugar that passed the filter then dried using an oven to fullfil water content standard set by SNI, a maximum of 3%.

### Optimization With RSM (Response Surface Methodology)

Optimization with RSM is based on research that has been done by (Mariana 2017). The RSM method is used to obtain data based on time, material weight, and material size so the water content of granulated palm sugar samples fulfill the SNI standards is a maximum of 3% (SNI, 1995). ). The experimental design was carried out using the Central Composite Design (CCD) with 20 trials using the Statistica 10 software.

### Moisture Test

The water content during drying of granulated palm sugar is calculated based on the following gravimetric equation (Normilawati et al., 2019).

$$\% \text{moisture level} = \frac{(W1 + W) - W2}{W} \times 100\% \quad (1)$$

Where, W is the weight of granulated palm sugar sample before drying (grams), W1 is the weight of vochdoos after drying (grams) and W2 is the weight of vochdoos + granulated palm sugar sample after drying (grams).

### Sucrose Test

50 ml of the sample filtrate was put into an Erlenmeyer added with 25 ml of distilled water and 10 ml of 30% HCl. Heat the solution on a water bath at 200°C. The solution is neutralized with 45% NaOH then diluted to a specific volume so that the solution contains 15-60 mg of reducing sugar. A total of 25 ml of the solution was taken and put into an Erlenmeyer, then added 25 ml of Luff School's solution. The solution in the Erlenmeyer was

cooled rapidly, and 15 ml of 20% KI was added, then carefully added 25 ml of 26.5% H<sub>2</sub>SO<sub>4</sub> solution. The iodine is titrated with a 0.1 N N-thiosulfate solution using 2-3 ml starch indicator. Calculation of sucrose level using the Eq. (2)

$$\begin{aligned} \text{Sucrose level} = & \\ & (\text{Reducing sugar content after inversion} \\ & - \text{Reducing sugar content after inversion}) \quad (2) \\ & \times 0.95 \end{aligned}$$

#### Reducing Sugar Test

Testing for reducing sugar was carried out using the Nelson Somogyi method (Haryanti, 2020). A total of 1 ml of a solution containing 0.02-0.08 mg/mL of glucose solution and 1 ml of Nelson's reagent was put into a test tube. Heated in a water bath for 20 minutes and then cooled. The solution that has been cooled is added with 1 mL of Arsenomolybdate reagent, then shaken until the Cu<sub>2</sub>O precipitate dissolves. After the precipitate dissolves, add 7 mL of distilled water, then shake until homogeneous. Optical Density (OD) was read by spectrophotometry at 540 nm wavelength.

#### Ash Level Test

Testing the ash content is carried out at temperatures above 450 °C (Yenrina et al., 2015). 3 - 5 grams of the sample is placed on a cup and put into an ashing furnace. The heating temperature is carried out in the first stages of 400 °C then the second stage is heated to 550 °C. The sample which has become ash is cooled in a desiccator. Ash content is calculated based on the Eq. (3).

$$\text{Ash level} = \frac{C - A}{B - A} \times 100\% \quad (3)$$

Where, A is the weight of empty cup, B is the weight of cup + sample and C is the weight of cup + sample (after ashing)

#### Calorie Value Test

The caloric value is calculated using the bomb calorimeter test (Sogandi, 2019). Calculation of heat of combustion based on the Eq. (4).

$$\Delta U_T = \frac{C\Delta T - \Delta U_1 - \Delta U_2}{m} \quad (4)$$

Where,  $\Delta U_T$  is heat of combustion of the sample (cal/g) and C is heat capacity of the calorimeter (cal/°C).

#### Protein Level Test

Protein level test using Kjeldahl method to determine protein levels (Siska<sup>7</sup> Apri, 2021). 0.4 grams of granulated palm sugar sample was put into a 100 ml Kjeldahl flask, added 25 ml of concentrated H<sub>2</sub>SO<sub>4</sub>, and heated for 2 hours. Then cooled and diluted with a 100 ml volumetric flask to the limit. 5 mL of the solution was put into a distillation and added 5 mL of 30% NaOH with a few drops of phenolphthalein indicator and distilled for 10 minutes. 10 ml of the distillation solution was added with 10 ml of 2% boric acid and then titrated with 0.01N HCl solution. The protein content is calculated based on the Eq. (5)

$$\begin{aligned} \text{Protein level} = & \\ & \frac{(V_1 - V_2) \times N \times 0,014 \times f_k \times 100}{w} \quad (5) \end{aligned}$$

Where  $V_1$  and  $V_2$  are the volume of HCl used for titration of sample and blank solutions, W is the sample weight, N is the normality of HCl and  $f_k$  is the conversion factor.

#### Fat Level Test

1-2 grams<sup>6</sup> sample is weighed and put into a beaker glass. 30 ml of 25% HCl and 20 ml of water were put<sup>6</sup> into a beaker glass, and then some boiling stones were added. The beaker glass was covered with a watch glass and boiled for 15 minutes. After that, the liquid is filtered in hot conditions by washing using hot water to prevent the acid reaction from occurring again. The filter paper and its contents were dried in an oven at 100-105 degrees.

The dried samples were extracted with hexane solvent for 5-6 hours at 80°C. The extracted solution was then distilled<sup>7</sup> separate the fat extract from the hexane solvent. The fat extract was dried in an oven at 100-105, then cooled and weighed. Calculation of fat content based on the Eq. (6).

$$\% \text{fat} = \frac{\text{mass of extract}}{\text{mass of sample}} \times 100\% \quad (6)$$

#### Carbohydrate Test

1 mL of the sample filtrate was added with 25 mL of distilled water. The filtrate and distilled water mixture were heated with a water bath for 1 hour. After that, the sample was dripped with three drops of PP and NaOH with 50% to neutral<sup>3</sup> H concentration. The neutral solution was diluted into



a 100 mL volumetric flask, shaken, and filtered to obtain the filtrate. The filtrate formed was shaken 0.05 mL with a pipette and then added 0.45 mL of distilled water and 0.5 mL of Nelson's reagent. The solution was then heated for 10 minutes and cooled. The cooled solution was added with 4 mL of arsenomolybdate reagent, then vortexed and incubated for 30 minutes to read the absorbance. The absorbance value will be used to calculate the carbohydrate content. Calculation of carbohydrate content with Eq. (7)

$$\% \text{Carbohydrate} = \frac{X \left( \frac{\text{mg}}{\text{L}} \right) \times \text{fp} \times \text{Total Volume (L)}}{w \times 100} \times 100\% \quad (7)$$

Where, X is the concentration (mg/L), fp is the dilution factor and w is the initial sample weight (kg).

## RESULTS AND DISCUSSION

The influence between variables consisting of time, material weight, and material size was used to determine the optimum moisture content of the red ginger granulated palm sugar drying process. The data processing results with statistical software 10 to obtain a relationship between the observed value and predicted value.

The results of the percentage error that obtained were analyzed by MAPE (Mean Absolute Percentage Error). So that a value of 2.58% was obtained. This value indicates that the experiments carried out are very accurate. The value of the data accuracy category on MAPE is as follows:

- < 10% = very accurate
  - 10-20% = good
  - 20-50% = reasonable
  - >50% = inaccurate
- (Maricar, 2019)

Based on Table 1. the values have shown the accuracy between the predicted and observed so it can be analyzed to know equations of the mathematical model. The data that were obtained as shown in Table 2.

Based on Table 2, the  $R^2$  value is 0.956. It shows that the model used is accurate because it is close to 1. Adjust  $R^2$  of 0.917 indicates that the relationship between the independent and response variables is getting stronger. The p-value for the time factor ( $X_1$ ) is 0.8892, and the weight of the material ( $X_2$ ) is 0.4193, which shows no significant

effect on the water content response because the p-value is > 0.05. Based on the interaction effect and significance, the response equation is shown in Eq. (8).

$$Y = 2.02379 + 0.00629X_1 + 0.03774X_2 - 0.136877X_3 - 0.18784X_1^2 + 0.31622X_2^2 + 0.13226X_3^2 + 0.46205X_1X_2 + 0.48250X_1X_3 + 0.30515X_2X_3 \quad (8)$$

Equation 8 is mathematical modeling in a polynomial equation of order 2 with significant coefficients on the model as in Table 2. The above equation is not the final equation because there are still insignificant effects. The linear factors of time ( $X_1$ ) and weight of the material ( $X_2$ ) must be neglected so that Eq. (8) is simplified to Eq. (9)

$$Y = 2.02379 - 0.136877X_3 - 0.18784X_1^2 + 0.31622X_2^2 + 0.13226X_3^2 + 0.46205X_1X_2 + 0.48250X_1X_3 + 0.30515X_2X_3 \quad (8)$$

The interaction effect between the variables obtained was validated using ANOVA, as shown in Table 3.

ANOVA determines the significance value of the independent variables that affect the dependent variable. The error tolerance limit ( $\alpha$ ) used is 5% or 0.05 so that the confidence level (1-0.05) = 0.95 or 95%. Based on Table 3, the time factor ( $X_1$ ) and material weight ( $X_2$ ) have a significance value of 0.88917 and 0.41932. So, these two factors do not affect the value of the optimum water level. However, the quadratic factor of time ( $X_1^2$ ) and weight of material ( $X_2^2$ ) had a significant effect on the optimum moisture content because it had a p-value of <0.05, namely 0.00642 and 0.00064. The interaction factor between time variables on the weight of material ( $X_1X_2$ ), time on the size of the material ( $X_1X_3$ ), and weight of the material on the size of the material ( $X_2X_3$ ) has a significant effect on the optimum water level because it has a p-value <0.05, which is 0.00043; 0.00035; and 0.00284.

In addition to the significance test, it is necessary to do a model fit test (Lack of Fit). Lack of fit means a discrepancy so that if  $H_0$  is accepted, then there is no discrepancy in the model, so it means that the model is appropriate (Pertiwi, 2014)

The hypothesis used in this test is as follows:

- $H_0$  = There is no lack of fit in the model
- $H_1$  = There is a lack of fit in the model

Table 1. Value of Observed and Predicted Moisture Content

Run	Value			Response Observed	Response Predicted	% Error
	X <sub>1</sub> (hour)	X <sub>2</sub> (grams)	X <sub>3</sub> (mesh)			
1	4	100	10	2.90	2.83	2.56
2	4	100	26	1.91	1.90	0.33
3	4	300	10	2.05	2.10	2.13
4	4	300	26	1.83	1.78	2.62
5	6	100	10	1.81	1.89	4.13
6	6	100	26	1.94	1.93	0.86
7	6	300	10	2.05	2.08	1.62
8	6	300	26	2.63	2.73	3.85
9	3.32	200	18	1.69	1.75	3.75
10	6.68	200	18	1.87	1.76	5.45
11	5	31.82	18	2.41	2.44	1.09
12	5	368.18	18	2.57	2.50	2.51
13	5	200	4.55	2.36	2.33	1.41
14	5	200	31.45	2.10	2.10	0.24
15	5	200	18	2.10	2.02	3.72
16	5	200	18	2.02	2.02	0.34
17	5	200	18	2.14	2.02	5.26
18	5	200	18	1.96	2.02	3.30
19	5	200	18	1.94	2.02	4.14
20	5	200	18	1.98	2.02	2.28

Table 2. Effect Estimation for Polynomial Equations of Order 2

Factor	Effect	Std.Err.	t	p
Mean/Interc.	2.0238	0.0323	62.5862	0.00000
X <sub>1</sub>	0.0063	0.0429	0.1466	0.8892 (p>0.05)
X <sub>1</sub> <sup>2</sup>	-0.1878	0.0418	-4.4971	0.0064
X <sub>2</sub>	0.0377	0.0429	0.8796	0.4193 (p>0.05)
X <sub>2</sub> <sup>2</sup>	0.3162	0.0418	7.5704	0.0006
X <sub>3</sub>	-0.1369	0.0429	-3.1898	0.0243
X <sub>3</sub> <sup>2</sup>	0.1323	0.0418	3.1665	0.0249
X <sub>1</sub> X <sub>2</sub>	0.4621	0.0561	8.2417	0.0004
X <sub>1</sub> X <sub>3</sub>	0.4825	0.0561	8.6064	0.0003
X <sub>2</sub> X <sub>3</sub>	0.3052	0.0561	5.443	0.0028
R <sup>2</sup>	0.95662	Adjust R <sup>2</sup>	0.91758	

For the model to fit, it is necessary to reject H<sub>1</sub> so that H<sub>0</sub> is accepted. In Table 3, the F<sub>table</sub> value is 1.50524 < F count is 4.77247, and the p table value is 0.33228, which is greater than 0.05, so the conditions for H<sub>0</sub> are accepted. It means that the model is appropriate and there is no model discrepancy so that the second-order polynomial equation model can be accepted.

#### Determination of Optimum Conditions

The profile and desirability determine the optimum conditions for the drying process. We used it to predict the optimum condition of the process to obtain valuable data for the validation process. The interaction of the three variables is known for its optimum condition by drawing a straight line vertically. The optimum conditions for

Table 3. Results of ANOVA Analysis with Statistica 10 Software

Factor	SS	df	MS	F	p
X <sub>1</sub>	0.00014	1	0.00014	0.02149	0.88917 (p>0.05)
X <sub>1</sub> <sup>2</sup>	0.12713	1	0.12713	20.22358	0.00642
X <sub>2</sub>	0.00486	1	0.00486	0.77377	0.41932 (p>0.05)
X <sub>2</sub> <sup>2</sup>	0.36026	1	0.36026	57.31085	0.00064
X <sub>3</sub>	0.06396	1	0.06396	10.17489	0.02427
X <sub>3</sub> <sup>2</sup>	0.06303	1	0.06303	10.02642	0.02491
X <sub>1</sub> X <sub>2</sub>	0.42698	1	0.42698	67.92509	0.00043
X <sub>1</sub> X <sub>3</sub>	0.46561	1	0.46561	74.07078	0.00035
X <sub>2</sub> X <sub>3</sub>	0.18623	1	0.18623	29.62641	0.00284
Lack of Fit	0.04731	5	0.00946	1.50524	0.33228
Pure Error	0.03143	5	0.00629		
Total SS	18.151	19			
R <sup>2</sup>	0.95662	Adjust R <sup>2</sup>	0.91758		




Figure 1. Optimum Condition of Drying Process on Moisture Level

the drying process of granulated palm sugar are shown in Figure 1.

A desirability value of 1 indicates that the goal solution has been fulfilled. While a desirability value of 0 indicates that the response must be discarded (Nurmiah et al., 2013). The prediction model produces a moisture response of 2.9019% with an optimum condition of 6.68 hours, material weight 368.18 grams, and material size 31.45 mesh.

The surface plot shows that the optimization of time, material size and material weight can affect the moisture content of granulated palm sugar.

**Validation of Optimum Conditions for Drying Process Optimization Results**

The optimum condition of granulated palm sugar was validated by running process and testing the water level. It aims to determine the

Table 4 Validation Results of Predicted Value and Observed Value of Software Statistics

Variable	Optimum Value Based on Statistica 10 Software	Water Level of Granulated palm Sugar	
		Predicted value	Observed value
X <sub>1</sub>	6.68 hour		
X <sub>2</sub>	368.18 gram	2.9019%	2.9016%
X <sub>3</sub>	31.45 mesh		
	% Error		0.0003%

Table 5. Chemical Characteristics of Red Ginger Granulated Palm Sugar Test Results

Chemical Characteristics	Test Results	Quality qualification SNI 01-3743-1995	Information
Sucrose	96.5967%	Minimum 90%	Fulfilled
Sugar reduction	6.0434%	Minimum 6%	Fulfilled
Ash level	1.8660%	Maximum 2%	Fulfilled
Caloric	379.8951 kcal	379.93 kkal	Fulfilled
Protein	2.4268%	2.45%	Fulfilled
Fat	0.3972%	0.41%	Fulfilled
Carbohydrate	91.5379%	91.61%	Fulfilled

match between the observed values from running validation and the predicted values from optimization with RSM. The validation value of the optimization results is shown in Table 4.

Based on the data in Table 4, the difference in the response value between the prediction software and the actual data is 0.0003%. The difference in value is less than 5% so that the validation results can be accepted. It proves that the test results' water content matches the program's predicted value (Mariana, 2017).

#### Chemical Characteristics of Red Ginger Granulated Palm Sugar Optimum Validation Results

The chemical characteristics observed include sucrose, sugar reduction, ash level, calories, protein, fat, and carbohydrates are presented in Table 5.

The sucrose level of the test result showed a value of 96.5967%. This value is under the quality requirements of SNI 01-3743-1995, which is at least 90%. A significant sucrose value affects the durability of granulated palm sugar because it is associated with a low content of reducing sugar, so it is more durable during storage (Zuliana et al., 2016). It is appropriate because the reducing sugar content in palm sugar and optimization results shows a low value of 6.0434%. This value is close to the minimum condition stated in SNI 01-3743-1995, at least 6%.

Low reducing sugar levels indicate the quality of the granulated palm sugar produced is good, because the sugar becomes stable and does not melt quickly (Indahyanti et al., 2014). The ash content of the test results shows a value of 1.8660%. It follows the quality requirements of SNI 01-3743-1995, which is a maximum of 2%. The ash level is related to the preservative level added in palm sugar. The higher the ash level in palm sugar, indicates the presence of mineral additives, one of which is preservatives added during the cooking process (Haryanti, 2020). Other chemical characteristics that are not a quality requirement for palm sugar in accordance with SNI 01-3743-1995 include calories, protein, fat, and carbohydrates. The results of the test on these characteristics are calories (379.8951 kcal), protein (2.4268%), fat (0.3972%), and carbohydrates (91.5379%). This value has met the quality of the Banten Agricultural Technology Assessment Agency (Heryani, 2016).

#### CONCLUSION

The second order polynomial equation with RSM has an R<sup>2</sup> value of 0.95662 with the simplified equation. The optimum operating condition of the drying results showed a response of water content of 2.9016% with a percentage error of 0.0003%, affecting the chemical characteristics of granulated palm sugar so that it fulfill the quality requirements of SNI 01-3743-1995, sucrose (96.5967%), sugar reduction (6.0434%), ash level



(1.8660%), calories (379.8951 kcal), protein (2.4268%), fat (0.3972 %) and carbohydrates (91.5379%).

## REFERENCES

- Assegaf, S., Kawilarang, A. P., Handajani, R. 2020. Antibacterial Activity Test of Red Ginger Extract (*Zingiber officinale var. Rubrum*) Against *Streptococcus pyogenes* In vitro. *Biomolecular and Health Science Journal*. 3(1): 24–27.
- Fatriani. 2019. Karakteristik Gula Semut dari Pengaron Sebagai Pemanis Pangan Alternatif. *Prosiding Seminar Nasional Lingkungan Lahan Basah*. 4(1): 34–37.
- Haryanti, P. 2020. Evaluasi Mutu Gula Kelapa Kristal (Gula Semut) Di Kawasan Home Industri Gula Kelapa Kabupaten Banyumas. *Jurnal Agroteknologi*. 5(1): 48–61.
- Heryani, H. 2016. Keutamaan Gula Aren dan Strategi Pengembangan Produk. (Edisi Pertama). Lambung Mangkurat Universiti Press, pp. 1-157.
- Indahyanti, E., Kamulyan, B. Ismuyanto, B. 2014. Optimasi Konsentrasi Garam Bisulfit pada Pengendalian Kualitas Nira Kelapa. *Jurnal Penelitian Saintek*. 19(1): 1–8.
- Kaushik, S., Jangra, G., Kundu, V., Yadav, J. P., Kaushik, S. 2020. Anti-viral Activity of *Zingiber officinale* (Ginger) Ingredients Against the Chikungunya Virus. *Virus Disease*. 31(3): 270–276.
- Mariana. 2017. Optimasi Formulasi dan Karakterisasi Fisikokimia dalam Pembuatan Daging Restrukturisasi Menggunakan Response Surface Methodology (Konsentrasi Jamur Tiram Serta Gel Porang dan Karagenan). *Jurnal Pangan dan Agroindustri*. 5(4): 83–91.
- Maricar, A. 2019. Analisa Perbandingan Nilai Akurasi Moving Average dan Exponential Smoothing untuk Sistem Peramalan Pendapatan pada Perusahaan XYZ. *Jurnal Sistem dan Informatika*. 13(2): 36–45.
- Meldayanoor, M., Ilmannafian, A. G., Wulandari, F. 2019. Pengaruh Suhu Pengeringan Terhadap Kualitas Produk Gula Semut dari Nira. *Jurnal Teknologi Agro-Industri*. 6(1): 1.
- Muhandri, T., Rifqi, D. M., Lestari, T., Widodo, S. 2020. Pelatihan Teknis dalam Rangka Perbaikan Mutu Gula Semut di Kabupaten Tasikamaya. *Agrokreatif: Jurnal Ilmiah Pengabdian kepada Masyarakat*. 6(3): 276–280.
- Normilawati, N., Fadlilaturrahmah, F., Hadi, S., Normaidah, N. 2019. Penetapan Kadar Air dan Kadar Abu pada Biskuit Yang Beredar Di Pasar Banjarbaru. *Jurnal Ilmu Farmasi*. 10(2): 51–55.
- Nurmiah, S., Syarief, R., Sukarno, S., Peranginangin, R., Nurmata, B. 2013. Aplikasi Response Surface Methodology pada Optimalisasi Kondisi Proses Pengolahan Alkali Treated Cottonii (ATC). *Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan*. 8(1): 9.
- Pertiwi, P. 2018. Aplikasi Response Surface Methodology (RSM) untuk Meningkatkan Kuat Tekan Paving Block dengan Campuran Abu Ampas Tebu. *Teknik Sipil Universitas Mataran*. 2(1): 1–10.
- Pratama, A., Wisdaningrum, O., Nugrahani, M. P. 2020. Pendampingan dan Penerapan Teknologi untuk Peningkatan Produktivitas Usaha Mikro Gula Semut. *Dinamisia: Jurnal Pengabdian Kepada Masyarakat*. 4(2): 275–284.
- Ritonga, A. M., Masrukhi, S. 2020. Pendugaan Umur Simpan Gula Kelapa Kristal Menggunakan Metode Akselerasi Berdasarkan Pendekatan Kadar Air Kritis. *Jurnal Teknologi Pertanian*. 21(1): 11–18.
- Sinamo, K. N., Hutabarat, N. D. M. R. 2021. Red Ginger Wedang to Strengthen Immune System Against Covid-19 of Children Living in an Orphanage. *ABDIMAS TALENTA: Jurnal Pengabdian Kepada Masyarakat*. 6(1): 60–67.
- SNI. 1995. *SNI 01-3743-1995 Gula Palma*. Badan Standar Nasional.
- Sogandi, S. 2019. Identifikasi Kandungan Gizi dan Profil Asam Amino dari Ikan Seluang (*Rasbora Sp*). *The Journal of Nutrition and Food Research*. 42(2): 73–80.
- Wilberta, N., Sonya, N. T., Lydia, S. H. R. 2021. Analisis Kandungan Gula Reduksi Pada Gula Semut dari Nira Aren yang Dipengaruhi pH dan Kadar Air.

- BIOEDUKASI (Jurnal Pendidikan Biologi). 12(1): 101–107.
- Wulandari, T. S. H., Nurhaniefa, A. M., Lestari, W. A. 2021. The Content Effectiveness in Red Ginger (*Zingiber officinale* Rosc.) Variety to Increase Immunity. 1st International Conference In Education, Science And Technology. 1(1): 459–463.
- Yenrina, I. R. 2015. Metode Analisis Bahan Pangan dan Komponen Bioaktif (Edisi Pertama). Andalas University Press. Padang, Indonesia.
- Zuliana, C., Widyastuti, E., Susanto, W. H. 2016. Pembuatan Gula Semut Kelapa (Kajian pH Gula Kelapa dan Konsentrasi Natrium Bikarbonat). Jurnal Pangan dan Agroindustri. 4(1): 109–119.

# Optimization of Drying Process for Production Red Ginger Granulated Palm Sugar Using Response Surface Methodology

## ORIGINALITY REPORT

6%

SIMILARITY INDEX

2%

INTERNET SOURCES

3%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 P A Handayani, D Hartanto, W T Eden, D F Anyelir, G R H H Salsabila. "Optimization of the pH of mosquito repellent from citronella using response surface methodology", IOP Conference Series: Earth and Environmental Science, 2022  
Publication 1%
- 2 Submitted to Academic Library Consortium  
Student Paper 1%
- 3 S. V. Ting. "Fruit Juice Assay, Rapid Colormetric Methods for Simultaneous Determination of Total Reducing Sugars and Fructose in Citrus Juices", Journal of Agricultural and Food Chemistry, 2002  
Publication 1%
- 4 Submitted to Universiti Teknologi Malaysia  
Student Paper 1%
- 5 Rika Sepriani, Deswandi. "Biochemical analysis of ginger-infused water with a combination of 1%

# lemon and mint leaves", E3S Web of Conferences, 2021

Publication

6

Submitted to Universitas Pelita Harapan

Student Paper

1 %

7

W T Eden, C O Rumambarsari. "Proximate analysis of soybean and red beans cookies according to the Indonesian National Standard", Journal of Physics: Conference Series, 2020

Publication

1 %

8

[repository.uinjkt.ac.id](https://repository.uinjkt.ac.id)

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On