

**BUKTI KORESPONDENSI ARTIKEL PADA  
JURNAL NASIONAL TERAKREDITASI**



**PENGUSUL**

Maharani Kusumaningrum, S.T., M.Eng / NIDN 0627039101

**UNIVERSITAS NEGERI SEMARANG**

Yth. Penilai pada usulan PAK

Bersama dengan surat ini, saya sertakan dokumen bukti - bukti korespondensi untuk karya penelitian pada Jurnal Nasional Terakreditasi (SINTA 2) dengan judul artikel: "Turmeric Extraction (*Curcuma Longa L*) Using The *Reflux* Method And Characterization", yang dimuat pada Jurnal Bahan Alam Terbarukan (JBAT), Vol. 11 No. 1, Hal: 85-91 Desember 2022, ISSN: 2303-0623.

Kronologi bukti korespondensi terdiri dari beberapa poin dan disusun dalam tabel di bawah Ini:

No	Tanggal	Aktivitas
1	29 Oktober 2022	Submit file artikel di JBAT (Sinta 2)
2	29 Oktober 2022	Menerima balasan email dari Editor bahwa telah submit artikel
3	31 Oktober 2022	Menerima proses review pada JBAT (Sinta 2), komentar Editor ke-1
4	31 Oktober 2022	Menerima pemberitahuan review artikel oleh Editor melalui email
5	9 November 2022	Mengirim hasil revisi tahap pertama ke Editor
6	14 November 2022	Artikel dinyatakan "accepted" dari Editor
7	30 November 2022	Diminta untuk membayar biaya proses publikasi artikel
8	30 November 2022	Artikel memasuki tahapan final <i>copyediting</i>
9	7 Desember 2022	Artikel telah terbit di jurnal JBAT

Demikian kronologi proses publikasi artikel di JBAT, agar dapat menjadi bahan periksa. Terimakasih.

Semarang, 13 Juni 2024

Hormat kami,



Maharani Kusumaningrum

## **Kronologi korespondensi publikasi artikel pada jurnal nasional terakreditasi**

Judul : Turmeric Extraction (*Curcuma Longa L*) Using The *Reflux* Method  
And Characterization

Jurnal : Jurnal Bahan Alam Terbarukan (JBAT)

Volume : 11

Nomor : 1

Tanggal publikasi : 7 Desember 2022

p-ISSN : 2303 – 0623

e-ISSN : 2407 – 2370

Halaman : 85 – 91

Penerbit : Teknik kimia Fakultas Teknik, UNNES

Pengindeks jurnal : SINTA 2

Penulis : Maharani Kusumaningrum\*, Hanif Ardhiansyah, Andika Wicaksana  
Putranto , Anita Trihardini , Putri Arum Kinanti, Duwi Naely Maslahah,  
Harianingsih.

doi: <https://doi.org/10.15294/jbat.v11i2.39784>

1) Submit file artikel di JBAT (Sinta 2) pada tanggal 29 Oktober 2022.

The screenshot shows the JBAT website interface. At the top, there is a navigation bar with links for 'About', 'Editorial Team', 'Reviewer', and 'Archive', along with a user profile icon and 'User Home'. Below this is a sidebar menu with various options like 'Focus & Scope', 'Editorial Policies', and 'Online Submission Here'. The main content area displays the submission details for article #39784, including the title 'Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization', the author 'Maharani Kusumaningrum', and the submission date 'October 29, 2022 - 10:55 AM'.

Home / User / Author / Submissions / #39784 / Summary

## #39784 Summary

Summary | Review | Editing

### Submission

Authors	Maharani Kusumaningrum, Hanif Ardiansyah, Andika Wicaksana Putranto, Anita Trihardini, Putri Arum Kinanti, Duwi Naely Maslahah, Harianingsih Harianingsih
Title	Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization
Original file	<a href="#">39784-101942-1-SM.docx</a> 2022-10-29
Supp. files	<input type="checkbox"/>
Submitter	Maharani Kusumaningrum
Date submitted	October 29, 2022 - 10:55 AM
Section	Articles
Editor	Haniif Prasetiawan
Abstract Views	1614

2) Menerima balasan email dari Editor bahwa telah submit artikel pada tanggal 29 Oktober 2022.

The screenshot shows a Gmail inbox with an email from 'jurnal.bat@mail.unnes.ac.id' titled '[JBAT] Submission Acknowledgement'. The email content includes a thank you message from the editor, the manuscript URL 'https://journal.unnes.ac.id/ojs/index.php/jbat/author/submission/39784', and the username 'maharani\_kusumaningrum'. The email is dated '29 Okt 2022, 10:55'.

Telusuri dalam email

[JBAT] Submission Acknowledgement Kotak Masuk x

- jurnal.bat@mail.unnes.ac.id lewat my.unnes.id kepada saya

Terjemahkan ke Indonesia

Maharani Kusumaningrum:

Thank you for submitting the manuscript, "Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization" to Jurnal Bahan Alam Terbarukan. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL:  
<https://journal.unnes.ac.id/ojs/index.php/jbat/author/submission/39784>  
Username: maharani\_kusumaningrum

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

-  
Jurnal Bahan Alam Terbarukan

Jurnal Bahan Alam Terbarukan (Journal of Natural Resources)  
<http://journal.unnes.ac.id/ojs/index.php/jbat>

3) Menerima proses review pada JBAT (Sinta 2), komentar Editor ke-1 pada tanggal 31 Oktober 2022.

Home / User / Author / Submissions / #39784 / Review

## #39784 Review

Summary | Review | Editing

### Submission

**Authors** Maharani Kusumaningrum, Hanif Ardhiyansyah, Andika Wicaksana Putranto, Anita Trihardini, Putri Arum Kinanti, Duwi Naely Maslahah, Harianingsih Harianingsih

**Title** Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization

**Section** Articles

**Editor** Haniif Prasetiawan

### Peer Review

Round 1

Review Version	39784-101943-1-RV.docx	2022-10-29
Initiated		2022-10-31
Last modified		2022-10-31
Uploaded file	Reviewer A 39784-102077-1-RV.docx	2022-10-31

4) Menerima pemberitahuan review artikel oleh Editor melalui email pada tanggal 31 Oktober 2022.

Gmail

JBAT

[JBAT] Editor Decision

Dr. Megawati Megawati megawati@mail.unnes.ac.id

Maharani Kusumaningrum:

We have reached a decision regarding your submission to Jurnal Bahan Alam Terbarukan, "Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization".

Our decision: Minor Revision


It will be better if your article goes through a professional proofreader for better grammar and vocabulary.

Thank you,

Best Regards,  
Dr. Megawati Megawati  
Scopus ID: 36959759900 Department of Chemical Engineering, Faculty of Engineering, Universitas Negeri Semarang, Indonesia  
megawati@mail.unnes.ac.id

Jurnal Bahan Alam Terbarukan (Journal of Natural Resources)  
<http://journal.unnes.ac.id/index.php/jbat>


5) Mengirim hasil revisi tahap pertama ke Editor pada tanggal 9 November 2022.



The screenshot shows the APTÉKIM submission portal. On the left, there are logos for APTÉKIM (Asosiasi Pendidikan Tinggi Teknik Kimia Indonesia) and ETROTEX (Pustaka Publikasi Group). The main interface includes the following fields and actions:

- Notify Editor:** Editor/Author Email Record 2022-10-31
- Editor Version:** None
- Author Version:** Two entries are listed: 39784-102693-1-ED.docx (2022-11-09) and 39784-102693-2-ED.docx (2022-11-29), each with a green 'Delete' button.
- Upload Author Version:** A 'Browse...' button with the text 'No file selected.' and a green 'Upload' button below it.


6) Artikel dinyatakan “accepted” dari Editor pada tanggal 14 November 2022.



The screenshot shows the 'Editor Decision' page in the APTÉKIM submission portal. On the left is the APTÉKIM logo. The main content area displays the following information:

- Decision:** Accept Submission 2022-11-14
- Notify Editor:** Editor/Author Email Record 2022-10-31
- Editor Version:** None

7) Diminta untuk membayar biaya proses publikasi tanggal 30 November 2022



The screenshot shows an email from 'Jurnal BAT' (jurnal.bat@mail.unnes.ac.id) received on 30 Nov 2022, 13:44. The subject is 'JBAT Publication Fee'. The email content is as follows:

Dear Dr. Maharani,

The publication fee of **JBAT** will cost Rp. 1.000.000,-. The publication fee is payable to BNI with acc. number 04 581 56424 a.n. Hanif Prasatiawan. Please make a confirmation after the payment has been done.

Thank You

—  
Prof. Dr. Megawati S.T., M.T.  
Editor in Chief  
Jurnal Bahan Alam Terbarukan (Journal of Biorefinery)  
Department of Chemical Engineering, Engineering Faculty  
Semarang State University  
Gd. E1 Lt. 2 Kampus Sekaran, Gunungpati, Semarang 50229  
Telp/Fax: (024) 3508101 ext.114  
Email: [jurnal.bat@mail.unnes.ac.id](mailto:jurnal.bat@mail.unnes.ac.id)  
Web Site: <http://journal.unnes.ac.id/nju/index.php/jbat>

8) Artikel memasuki tahapan final *copyediting* pada tanggal 30 November 2022

- Editorial Board
- Reviewer Acknowledgement
- Abstracting/Indexing
- Ethics Statement
- Scopus Citation Analysis
- Subscription Form
- Open Access Policy
- Contact Us

p-ISSN 2303-0623

e-ISSN 2407-2370



COLLABORATION WITH



TOOLS



grammarly.com



Submission

**Authors** Maharani Kusumaningrum, Hanif Ardiansyah, Andika Wicaksana Putranto, Anita Trihardini, Putri Arum Kinanti, Duwi Naely Maslahah, Harianingstih Harianingstih

**Title** Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization

**Section** Articles

**Editor** Haniif Prasetiawan

Copyediting

Copyedit Instructions

Review Metadata

	Request	Underway	Complete
1. Initial Copyedit	2022-11-30	—	2022-11-30
File: <input type="text" value="39784-102862-1-CE.docx"/> 2022-11-14			
2. Author Copyedit	—	—	
File: None			
<input type="button" value="Browse..."/> No file selected.			
<input type="button" value="Upload"/>			
3. Final Copyedit	—	—	2022-11-30
File: None			

Copyedit Comments No Comments

Layout

Galley Format	File	
1. PDF <input type="button" value="View Proof"/>	<input type="text" value="39784-103765-2-PB.pdf"/> 2022-12-06	806
Supplementary Files	File	
1. Turmeric Extraction (Curcuma Longa L) Using The Reflux Method And Characterization	<input type="checkbox"/>	

Layout Comments No Comments

Proofreading

Review Metadata

	Request	Underway	Complete
1. Author	—	—	
2. Proofreader	2022-11-30	—	2022-11-30
3. Layout Editor	2022-11-30	—	2022-11-30

Proofreading Corrections No Comments

9). Artikel telah terbit di jurnal JBAT pada tanggal 7 Desember 2022

Status						
Status	Published	Vol 11, No 2 (2022): December 2022 [Nationally Accredited - Sinta 2]				
Initiated	2022-11-30					
Last modified	2022-12-07					

---

[Home](#) / [User](#) / [Author](#) / [Archive](#)

---

## Archive

[Active](#) | [Archive](#)

ID	MM-DD Submit	Sec	Authors	Title	Status
39784	10-29	ART	Kusumaningrum, Ardhiansyah, Putranto,...	Turmeric Extraction (Curcuma Longa L.) Using The Reflux...	Vol 11, No 2 (2022): December 2022 [Nationally Accredited - Sinta 2]

Hasil review artikel tahap ke-1 dari Editor



# Turmeric Extraction (*Curcuma Longa L*) Using The Reflux Method And Characterization

## Abstract

Turmeric (*Curcuma longa L*) is one of the many spices ~~groups~~ that grow on the Asian continent, especially Southeast and South Asia, which can be utilized to its full potential, especially the antioxidant compounds contained in curcuminoids. This study aims to determine the effect of different forms of turmeric, namely *simplicia* (turmeric powder), fresh (grated turmeric and pieces) with a solute/solvent ratio (w/v) (1:5). The viscous extract in the form of a paste ~~is was~~ obtained after the distillation process and solvent evaporation. The best extracts and essential oils obtained were extracts from powdered turmeric samples with yields of 8.28 (% w/w) and essential oils of 0.44 g, which were clearer than the other two samples. The analysis showed that the sample of turmeric with the highest antioxidant activity was a sample of freshly grated turmeric with an IC50 value of 114.7 ppm with a moderate level of antioxidant activity. The cut turmeric sample has an IC50 value of 158.3 ppm, which is included in the weak antioxidant activity. In comparison, the powdered turmeric sample has an IC50 value of 134.1 ppm with a moderate level of strength.

**Keywords:** Antioxidant; DPPH; Extraction; Reflux; Turmeric (*Curcuma longa L*)

## INTRODUCTION

In big cities, air pollution is a problem that always increases yearly (Abidin & Hasibuan, 2019). Even if you live in a big city, your body's oxidative

burden will be much higher than in a place with lower pollution, such as in the countryside. An environment with a relatively high pollution level triggers the emergence of free radicals. Free radicals are molecules that have unpaired

Commented [WU1]: Tolong diperjelas maksudnya?

Commented [WU2]: Berikan istilah resmi yang umum

Commented [WU3]: Hilangkan kata 'you' dan grammar

electrons in their outer orbitals, so they are reactive and unstable (Sopiah et al., 2019). These radicals tend to have a chain reaction which, if they occur in the body, can cause ongoing and continuous damage. Free radicals produced by cigarette smoke, radiation, pollutants, and other radical-triggering substances can cause chronic diseases such as heart attacks, cancer, and cataracts to decreased kidney function (Fakriah et al., 2019). Antioxidant compounds are needed to prevent free radicals. (Sopiah et al., 2019).

Antioxidants are inhibitors of the oxidation process, even at relatively small concentrations. If free radicals do not bind to antioxidants, the oxidation reaction will continue to cause cell damage (Andarina & Djauhari, 2017). One of the plants that contain antioxidants is turmeric.

Turmeric is one of the spices that are commonly found in Indonesia. Turmeric or turmeric rhizome contains curcumin and its derivatives by 3-15% (curcumin 71.5%, demethoxycurcumin 19.4% and bisdemethoxycurcumin 9.1%). The following chemical content

is phenylpropene and other phenolic components such as terpenes, monoterpenes, sesquiterpenes, diterpenes, triterpenes, alkaloids, steroids, and fatty acids (Suprihatin et al., 2020). Curcumin is insoluble in ether, soluble in oil, brownish red in alkali, and light yellow in acid (Rezki et al., 2015). The level of antioxidants in turmeric is strongly influenced by the solvent used. The more polar the solvent, the higher the antioxidant content attracted by turmeric (Pratiwi & Wardaniati, 2019).

The uptake of curcumin in a plant allows the removal of essential nutrients and biological activity during processing. One of the most effective methods is extraction (K-Sa'adah, 2019).

Turmeric can be extracted using several methods, namely the soxhlet extraction method, maceration, reflux extraction and hydrodistillation. The reflux method is more efficient and requires a shorter time (Yurleni, 2018). Turmeric can be extracted using several methods, namely the soxhlet extraction method, maceration, reflux extraction and hydrodistillation. The reflux method

is more efficient and requires a shorter time (Yurleni, 2018). Extraction by heating or reflux is a solvent-based extraction for a certain time with a limited and relatively constant amount of solvent (Susanty & Bachmid, 2016). In addition, extraction using water and acetone solvents is more suitable using the reflux method than maceration and soxhletation because the reflux method has higher yields of water solvent extract and acetone (Putra et al., 2014). Aquades is considered a solvent because it is a non-toxic stable solvent that tends to be safer, environmentally friendly, and cheaper in terms of cost (H. Sa'adah & Nurhasnawati, 2017). The extract's yield can be higher because the distilled water binds more of the extracted material due to the large number of hydrogen bonds formed (Masyitoh et al., 2016).

Turmeric has a high potential to be a source of natural antioxidants. However, to produce maximum curcumin compounds, research still needs to examine the effect of turmeric form on antioxidant activity. This study aims to determine the effect of variations in the shape of turmeric rhizome as a

solute in the extraction of curcumin from the reflux process using distilled water as a solvent. Variations in the curcumin extraction process are carried out with variations in the form of turmeric, which are powdered, grated, and cut.

#### **MATERIALS AND METHOD**

This research was conducted at the Chemical Engineering Operations Laboratory, Faculty of Engineering, ~~the State University of Semarang~~UNNES, with three types of sample treatments, namely variations in the form of turmeric rhizome (1) powder, (2) grated, and (3) pieces. Observations made include the appearance and weight of the oil and extracts produced. In addition, a UV-Vis spectrophotometer was also used to analyze DPPH (2,2-diphenyl-1-picrylhydrazil). The data obtained were analyzed with the GraphPad Prism 8 program to determine the level of strength of its antioxidant activity.

#### **Research Implementation Procedure**

The procedure for carrying out this research consisted of four stages: the sample preparation stage, the extraction

**Commented [WU5]:** Dicek lagi gramarnya, pakai passive past tense.

**Commented [WU4]:** Dicek lagi, jangan ada nama depannya

process, the antioxidant activity test, and data analysis.

### **Sample Preparation**

Dry turmeric powder (simplicia) has been sifted and weighed as much as 50 grams. Meanwhile, fresh turmeric is peeled, cleaned, and then grated using a grater and cut to a size of  $\pm 3-5$  mm using a knife and a cutting board. Grated turmeric pieces weighed as much as 50 grams.

### **Extraction Process**

The process of making turmeric extract using the reflux method. Samples weighing 50 grams and 250 mL of distilled water were put into neck flask 2. Next, neck flask two was paired with reflux extraction tools. The reflux process was carried out for 150 minutes at 100°C. After the process is complete and cooled, filtering is carried out to separate the filtrate and residue. The filtrate resulting from the reflux process was then distilled until the remaining filtrate in the distillation flask was  $\pm 25$  mL. The filtrate is then evaporated using an oven at a temperature of 100°C to evaporate the remaining solvent. Then

after a constant filtrate mass was produced during evaporation, a thick extract was produced in the form of a paste.

### **Antioxidant Activity Test**

Antioxidant activity test using DPPH (2,2-diphenyl-1-picrylhydrazil), whose absorbance was measured using a UV-Vis spectrophotometer.

### **Data Analysis**

The absorbance value data obtained from the analysis of the UV-Vis spectrophotometer were used to calculate the inhibition value (% antioxidant activity). Then the resulting data were analyzed using the GraphPad Prism 8 program.

## **RESULT AND DISCUSSION**

### **Effect of Variation in Form on Extract Results**

The observations found that the appearance was in the form of a slightly reddish brown paste (viscous extract), soluble in alcohol. The curcumin obtained can be seen in **Figure 3.1**. The

results of the qualitative analysis research are not much different from the journals reported by (Popuri & Pagala, 2013) and also (Rezki et al., 2015). Of the three samples studied, the brown color in sample 1, the dry powder sample, was more concentrated than the other two. In comparison, the colors of samples 2 and 3, namely samples of fresh turmeric in grated form and pieces of both, have almost no difference in color.



**Figure 3.1.** Concentrated extract of turmeric sample (1) powder (2) grated (3) pieces

Yield is the percentage ratio between the extract portion and the total weight of the initial sample used (Sari et al., 2020). Data on the % yield of turmeric extract in various forms of *Simplicia* (dry powder) and fresh (grated and cut) with distilled water can be seen in **Table 3.1**.

**Table 3.1.** Data on % yield of turmeric extract (*Curcuma Longa L*)

Sample shape	Initial sample weight (g)	Extract weight (g)	Yield (% w/w)
Powder	50	4,14	8,28
Grater	50	4,08	8,16
Piece	50	3,97	7,94
Average yield value			8,13

**Commented [WU8]:** Masih pakai koma

The average yield value of the extraction results was 8.13% (w/w). This value is influenced by several factors, one of which is the particle size of the sample (Wijaya et al., 2018). The size of the sample affects the amount of yield. The smaller the sample surface area will further expand the contact and increase the solvent interaction, the more yields will be produced (Sineke et al., 2016).


### The Effect of Variations in Forms on the Yield of Essential Oils

From the observations made, it was found that the essential oil was distilled by distillation. From the results obtained that the good essential oil was from a variety of powder forms because the volatile oil produced was very clear, and the second best result was produced through turmeric rhizome material with grated treatment because the more small surface area of the material affects the yield of the essential oil itself. The

**Commented [WU6]:** Cukup Tabel 1. dst

**Commented [WU7]:** Italic atau nggak?

results of these essential oils can be seen in **Figure 3.2** and **Table 3.2**.



**Figure 3.2.** The results of the turmeric essential oil sample (1) powder (2) grated (3) pieces

**Table 3.2.** Data on essential oil yields of turmeric extract (*Curcuma Longa L*)

Sample shape	Initial sample weight (g)	Oil weight (g)
Powder	50	0,44
Grater	50	0,29
Piece	50	0,20

Based on the results obtained in **Table 3.2**, it is known that the highest essential oil yield was sample 1, namely turmeric powder. The highest essential oil yield was sample 3, namely the cut sample. The result indicates that the smaller the sample surface area, the more volatile oil is produced (Sineke et al., 2016).

#### Antioxidant Analysis

Quantitative testing using a UV-Vis spectrophotometer with a

wavelength of 517 nm. The test was carried out by dividing the sample solution into five concentration variations, namely 50 ppm, 100 ppm, 150 ppm, 200 ppm, and 250 ppm. The test solution added with DPPH solution was incubated for 30 minutes at room temperature to optimize binding between antioxidant compounds in capturing unpaired electrons from free radicals. Research (Irianti et al., 2016) stated that DPPH has a stable absorbance at 30 to 40 minutes. After incubation, a color change from purple to yellowish was caused by antioxidants reacting with DPPH, stabilizing free radicals and reducing DPPH (Erviana et al., 2016). The color change of the turmeric extract solution from purple to yellow indicates that the turmeric extract has antioxidant potential. The DPPH method was chosen because it has several advantages, including being simple, easy, fast, sensitive, and requiring a small number of samples (Erviana et al., 2016).

The results of antioxidant measurements using a UV-Vis spectrophotometer were obtained in the form of absorbance values from the control and sample, which were then

used to determine the percent antioxidant activity (% inhibition) (Pratiwi & Wardaniati, 2019). The wavelength used in the UV-VIS spectrophotometer is 517 nm because DPPH provides strong absorption at a wavelength of 517 nm with a dark violet color which provides information on the reactivity of the compound tested with a stable radical (Katrin & Bendra, 2015). The following equation determines the % antioxidant activity: (Kuntorini & Astuti, 2012). The following equation determines the % antioxidant activity: (Kuntorini & Astuti, 2012).

$$\% \text{ antioxidant activity} = x \cdot 100\%$$

Description :

A<sub>0</sub> = Absorbance of control solution

A<sub>1</sub> = Absorbance of sample solution

The absorbance and % inhibition values in the sample can be seen in **Table 3.3**. Table 3.3 shows that the greater the extract concentration, the greater the value of % antioxidant activity (Mariani et al., 2018). The highest % activity value was obtained from sample 3, namely sliced turmeric with a concentration variation of 250 ppm

According to (Handayani et al., 2014), percent (%) antioxidant activity indicates the ability of an antioxidant compound to inhibit free radicals, namely the number of hydrogen atoms of an antioxidant compound that captures DPPH radicals so that they are reduced to more stable DPPH-H.

Commented [WU9]: Cek penulisan in text citation

**Table 3.3.** Data on absorbance and % inhibition

Concentration (ppm)	Absorbance			% Inhibition		
	1	2	3	1	2	3
50	0,375	0,379	0,354	52,711	52,327	55,471
100	0,339	0,352	0,327	57,250	55,723	58,867
150	0,308	0,350	0,326	61,160	55,974	58,993
200	0,295	0,337	0,276	62,799	57,610	65,283
250	0,260	0,327	0,260	67,213	58,867	67,295
Control	0,793	0,795	0,795	-	-	-

Commented [WU10]: Masih koma

Description :


Sample 1 = turmeric powder

Sample 2 = grated turmeric

Sample 3 = turmeric pieces

The value of IC<sub>50</sub> (50% Inhibitor Concentration) determines the radical scavenging activity. This value describes the concentration of the test compound

that can capture radicals by 50% (Erviana et al., 2016). The smaller the IC<sub>50</sub> value, the higher the antioxidant activity (Sumarlan et al., 2018). The IC<sub>50</sub> value using a non-linear regression equation using *GraphPad Prism 8 software*. IC<sub>50</sub> value data is shown in **Table 4.4**.



**Commented [WU11]:** Gambar nya pecah, minta dibuat ulang saja pakai origin pro

**Figure 3.3.** The curve of the relationship of % antioxidant activity to the concentration of turmeric samples (a) powder (b) grated (c) slices from *GraphPad Prism* analysis

**Table 3.4.** The IC<sub>50</sub> value of the turmeric extract sample

Sample	IC <sub>50</sub> value(ppm)
--------	-----------------------------



Tumeric powder	134,1
Grated tumeric	114,7
Cut tumeric	158,3

Based on the results obtained in **Table 3.4**, it is known that the lowest IC<sub>50</sub> value is sample 2, grated turmeric, which shows that its antioxidant activity is higher than the other two samples (Yuliani et al., 2016). The correlation between the IC<sub>50</sub> value and the strength of antioxidant activity is presented in **Table 3.5**.

**Table 3.5.** Strength Level of Antioxidant Activity (Tristantini et al., 2016)

Antioxidant Properties	IC50 value (ppm)
Very strong	<50
Strong	50-100
Medium	100-150
Weak	150-200

Table 3.4 and Table 3.5 show that the best results were obtained in samples of grated turmeric with an IC<sub>50</sub> value of 114.7 in the range of 100-150, which indicates that the antioxidant activity is

moderate (Yuliani et al., 2016). Likewise, turmeric powder samples have weak antioxidant activity with an IC<sub>50</sub> value of 134.1. At the same time, the cut sample has the largest IC<sub>50</sub> value, 158.3, in the range of 150-200, so it is included in the weak category (Yuliani et al., 2016). This difference in IC<sub>50</sub> value can be caused by the number of antioxidants in the extract. This phenomenon can occur due to antioxidant damage in the extract, which is influenced by the length of contact time between the active substance and the solvent, whose temperature increases due to prolonged heating (Tristantini et al., 2016).

The results of antioxidant activity obtained are not much different from the statement (Pratiwi Isna, 2019), namely the IC<sub>50</sub> value of dried powdered turmeric samples is lower than the IC<sub>50</sub> value of fresh-cut turmeric samples, which indicates that the antioxidant activity of powdered turmeric *Simplicia* samples is higher. However, this study

**Commented [WU12]:** Pakai nama belakang aja, referensi nya disesuaikan

also produced the best IC<sub>50</sub> value in grated turmeric samples, where the surface area was much smaller than the cut samples.

## CONCLUSION

The difference in variations in the form of simplicity and fresh turmeric has an effect, although not too large, on the results of the reflux method turmeric extract with distilled water, namely the highest yield in the powdered *Simplicia* sample, which is 8.28% (w/w) and the weight of the oil produced is 0.44 g. The antioxidant activity tested by the DPPH method was also influenced by differences in *simplicia* (dried powder) and fresh (grated and cut) with the best IC<sub>50</sub> value, namely grated fresh turmeric sample of 114.7 ppm with a medium strength category.

## REFERENCES

- Abidin, J., & Hasibuan, F. A. (2019). Pengaruh dampak pencemaran udara terhadap kesehatan untuk menambah pemahaman masyarakat awam tentang bahaya dari polusi udara. *Prosiding SNFUR-4, September*, 978–979.
- Andarina, R., & Djauhari, T. (2017). Antioksidan Dalam Dermatologi. *Jurnal Kedokteran Dan Kesehatan*, 4(1), 39–48.
- Erviana, L., Malik, A., & Najib, A. (2016). Uji Aktivitas Antiradikal Bebas Ekstrak Etanol Daun Kemangi (*Ocimum basilicum* L.) Dengan Menggunakan Metode DPPH. *Jurnal Fitofarmaka Indonesia*, 3(2), 164–168. <https://doi.org/10.33096/jffi.v3i2.217>
- Fakriah, Kurniasih, E., Adriana, & Rusydi. (2019). Sosialisasi bahaya radikal bebas dan fungsi antioksidan alami bagi kesehatan. *Jurnal Hasil-Hasil Penerapan IPTEKS Dan Pengabdian Kepada Masyarakat Sosialisasi*, 3(1), 1–7.
- Handayani, V., Ahmad, A. R., Sudir, M., Etlingera, P., & Sm, R. M. (2014). Uji Aktivitas Antioksidan Ekstrak Metanol Bunga dan Daun Patikala (*Etlingera elatior* (Jack) R. M. Sm) Menggunakan Abstrak. *Pharm Sci Res*, 1(2), 86–93.
- Irianti, T., Murti, Y. B., Kanistri, D. N.,

**Commented [WU13]:** Wahyuningtyas, S. E. P., Permana, I. D. G. M., & Wiadnyani, A. A. I. (2017) tika disitasi di text

Cek lagi semua penulisan referensi, ada beberapa yang belum ada halamannya

- Pratiwi, D. R., Kuswandi, & Kusumaningtyas, R. A. (2016). DPPH Radical Scavenging Activity Of Aqueous Fraction From Ethanolic Extract Of Talok Fruit (Muntingia calabura L.). *Traditional Medicine Journal*, 21(1), 38–47. <https://doi.org/10.22146/tradmedj.10724>
- Katrin, & Bendra, A. (2015). *Aktivitas Antioksidan Ekstrak , Fraksi dan Golongan Senyawa Kimia Daun Premna oblongata Miq . Abstrak*. 2(1).
- Kuntorini, E. M., & Astuti, M. D. (2012). Penentuan Aktivitas Antioksidan Ekstrak Ethanol Bulbus Bawang Dayak (Eleutherine americana Merr.). *Sains Dan Terapan Kimia*, 4, 15–22.
- Mariani, S., Rahman, N., & Pembahasan, H. (2018). Uji aktivitas antioksidan ekstrak buah semangka (Citrullus lanatus) Antioxidant Activity Test of Watermelon (Citrullus lanatus) Fruit Extracts. *Jurnal Akademika Kimia*, 7(May), 96–101.
- Masyitoh, Dewanti, & Setyorini. (2016). Analisis Profil Protein Ekstrak Aquades dan Etanol Daun Mimba (Azadirachta Indica A . Juss ) dengan Metode SDS-PAGE. *Pustaka Kesehatan*, 4(3), 533–539.
- Popuri, A. K., & Pagala, B. (2013). Extraction of curcumin from turmeric roots. *International Journal of Innovative Research Study*, 2(5), 289–299.
- Pratiwi, D., & Wardaniati, I. (2019). *Pengaruh Variasi Perlakuan ( Segar Dan Simplisia )*. 11(2).
- Pratiwi Isna, D. W. (2019). Pengaruh Variasi Perlakuan (Segar dan Simplisia) Rimpang Kunyit (Curcuma domestica) Terhadap Aktivitas Antioksidan dan Kadar Fenol Total. *Jurnal Farmasi Higea*, 11(Vol 11, No 2 (2019): Jurnal Farmasi Higea Vol 11. No. 2 Tahun 2019), 159–165.
- Putra, A. A. B., Bogoriani, N. W., Diantariani, N. P., & Sumadewi, N. L. U. (2014). IH on a Shoestring. *Kimia*, 8(1), 27. <https://doi.org/10.3320/1.2928419>
- Rezki, R. S., Anggoro, D., & MZ, S. (2015). Ekstraksi Multi Tahap Kurkumin dari Kunyit

- (Curcumadomestica Valet) Menggunakan Pelarut Etanol. *Jurnal Teknik Kimia USU*, 29, 29–34.
- Sa'adah, H., & Nurhasnawati, H. (2017). Perbandingan Pelarut Etanol Dan Air Pada Pembuatan Ekstrak Umbi Bawang Tiwai (*Eleutherine americana* Merr) MENGGUNAKAN METODE MASERASI. *Jurnal Ilmiah Manuntung*, 1(2), 149. <https://doi.org/10.51352/jim.v1i2.27>
- Sa'adah, K. (2019). *Uji Aktivitas Antiinflamasi Ekstrak Rimpang Temu Putih ( Curcuma zedoaria ( Christm.) Roscoe ) Dengan Metode Stabilisasi Membran Sel Darah Merah Secara Invitro.*
- Sari, M., Apriandi, A., Suhandana, M., Studi, P., Hasil, T., Ilmu, F., Dan, K., Maritim, U., & Ali, R. (2020). *Uji Toksisitas Ekstrak Daun Beruwat Laut (Scaevola taccada) Dengan Metode Brine Shrimp Lethality Test (BSLT) Toxicity Test of Beach Naupaka Leaves (Scaevola taccada) Extract Using the Brine Shrimp Lethality Test (BSLT) Method.* 03(April), 37–46.
- Sineke, F. U., Suryanto, E., & Sudewi, S. (2016). Penentuan Kandungan Fenolik Dan Sun Protection Factor (Spf) Dari Ekstrak Etanol Dari Beberapa Tongkol Jagung (*Zea mays L.*). *Pharmakon*, 5(1), 275–283. <https://doi.org/10.35799/pha.5.2016.11316>
- Sopiah, B., Muliasari, H., & Yuanita, E. (2019). Skrining Fitokimia dan Potensi aktivitas Antioksidan Ekstrak Etanol Daun Hijau dan Daun Merah Kastuba. *Jurnal Ilmu Kefarmasian Indonesia*, 17(1), 27. <https://doi.org/10.35814/jifi.v17i1.698>
- Sumarlan, S. H., Susilo, B., Mustofa, A., & Mu'nim, M. (2018). Ekstraksi Senyawa Antioksidan Dari Buah Strawberry (*Fragaria X Ananassa*) dengan Menggunakan Metode Microwave Assisted Extraction (Kajian Waktu Ekstraksi dan Rasio Bahan dengan Pelarut). *Jurnal Keteknik Pertanian Tropis Dan Biosistem*, 6(1), 40–51.
- Suprihatin, T., Rahayu, S., Rifa'i, M., & Widyarti, S. (2020). Senyawa pada

- Serbuk Rimpang Kunyit (*Curcuma longa* L.) Yang Berpotensi sebagai Antioksidan. *Buletin Anatomi Dan Fisiologi*, 5(1), 35–42. <https://doi.org/10.14710/baf.5.1.2020.35-42>
- Susanty, & Bachmid, F. (2016). Comparison Of Maceration And Reflux Extraction Methods To Phenolic Levels Of Corn Cob Extract (*Zea mays* L.). *Jurnal Konversi*, 5(2), 87.
- Tristantini, D., Ismawati, A., Pradana, B. T., & Gabriel, J. (2016). Pengujian Aktivitas Antioksidan Menggunakan Metode DPPH pada Daun Tanjung (*Mimusops elengi* L.). *Universitas Indonesia*, 2.
- Wahyuningtyas, S. E. P., Permana, I. D. G. M., & Wiadnyani, A. A. I. (2017). Pengaruh Jenis Pelarut Terhadap Kandungan Senyawa Kurkumin Dan Aktivitas Antioksidan Ekstrak Kunyit (*Curcuma domestica* Val.). *Itepa*, 6(2), 61–70. <https://ojs.unud.ac.id/index.php/itepa/article/view/36950/22387>
- Wijaya, H., Novitasari, & Jubaidah, S. (2018). *Perbandingan Metode Ekstraksi Terhadap Rendemen Ekstrak Daun Rambai Laut (*Sonneratia caseolaris* L. Engl)*. 4(1), 79–83.
- Yuliani, N. N., Sambara, J., & Mau, M. A. (2016). Uji Aktivitas Antioksidan Fraksi Etilasetat Ekstrak Etanol Rimpang Jahe Merah (*Zingiber officinale* var. *Rubrum*) Dengan Metode DPPH(1,1-Diphenyl-2-Picrylhydrazyl). *Jurnal Info Kesehatan*, 14, 1092–1111.
- Yurleni. (2018). Penggunaan Beberapa Metode Ekstraksi Pada Rimpang Curcuma Untuk Memperoleh Komponen Aktif Secara Kualitatif (Use of Multiple Methods of Extraction on Curcuma Rhizomes To Acquire Qualitative Active Component). *Biospecies*, 11(1).